

# New data on the Arachnofauna (Arachnidae Araneae Opiliones) of Enna and surrounding areas (Sicily, Italy) with some first reports for Sicily

Antonino Dentici<sup>1</sup> & Franco Ciro Amata<sup>2</sup>

<sup>1</sup>Via Enrico Cialdini 2, 90124 Palermo, Italy; e-mail: antonino.dentic.1991@gmail.com; Orcid: 0000-0003-0632-1633

<sup>2</sup>Piazza S. Antonino 1, 94018 Troina (Enna), Italy; e-mail: amata.francociro@gmail.com

## ABSTRACT

In this paper we present faunistic data on Arachnid collected, with sampling on sight during 2019 and 2020, in Enna (Sicily, Italy) and surroundings areas. The Arachnofauna of this area of the Island is poorly studied, to date only 16 species of spiders are reported for this municipality. 180 taxa of spiders are known in total now, 15 of these (1 first report in Italy) and 6 genera are new for Sicily - *Priatha nana* (Simon, 1868) (Filistatidae), *Micaria coarctata* (Lucas, 1846), *Micaria dives* (Lucas, 1846), *Poecilochroa loricata* Kristscher, 1996 (Gnaphosidae) (new for Italy), *Agroeca* sp., *Scotina celans* (Blackwall, 1841) (Lycosidae), *Bolyphantes luteolus* (Blackwall, 1833), *Centromerus arcarius* (O.Pickard-Cambridge, 1873), *Lepthyphantes minutus* (Blackwall, 1833), *Linyphia mimonti* Simon, 1885, *Megalepthyphantes collinus* (L. Koch, 1872), *Walckenaeria corniculans* (O. Pickard-Cambridge, 1875), *Walckenaeria monoceros* (Wider, 1834) (Linyphiidae), *Synageles venator* (Lucas, 1836) (Salticidae), *Enoplognatha afrodite* Hippa et Oksala, 1983 (Theridiidae) - and 3 species of opilions. Additional biological and taxonomic notes on these spiders are also provided.

## KEY WORDS

Araneofauna; Spiders; Opilions; new records; Sicily.

Received 02.11.2023; accepted 08.01.2024; published online 30.06.2024

Proceedings of 6th International Congress on Biodiversity “*Biodiversity and the new scenarios on alien species, climate, environment and energy*” - Trapani (Italy, Sicily) 2–3 September 2022

## INTRODUCTION

In the naturalistic bibliography there are nospecific works on arachnofauna on the area of Enna (Sicily). To date, there were only 16 known spiders species, belonging to 15 genera and 9 families (Al-icata, 1964 & 1966; Brignoli, 1968, 1977; Cantarella, 1974, 1980; Dentici & Amata, 2018; Hansen, 2005; Lugetti & Tongiorgi, 1969).

The few species known for this area and the recent news reports for Sicily (Dentici & Amata, 2021; Dentici & Galasso, 2021; Dentici et al., 2021, 2022; Dentici, 2022) have prompted us to deepen

the study, trying to produce a more complete list of spider species observable in this area as reported in the following work.

## MATERIAL AND METHODS

All samples were collected on sight on plants, under the rocks or in their web. All samples were photographed in nature with a Canon EOS 7D Mark II with a EF 100mm Macro with 20 mm extension rings and 1.4 canon multiplier (F.C.Amata: [www.amatafoto.it](http://www.amatafoto.it)).

All the samples were taken in laboratory, were treated with ethyl acetate and observed with binoculars for identification;

They are preserved in A. Dentici collection (Palermo, Italy). The specimens were stored in centrifuge tubes of different sizes, depending on the size of the sample, and were fixed in 75% ethanol (Levi, 1977). The samples are stored in the collection of one author (A. Dentici) and the “legit” is indicated for each sample. Each locality and/or collection site is in the original language (Italian).

The classification, taxonomic order and nomenclatural arrangement for spiders follow Roberts (1995), Trotta (2004), Pantini & Isaia (2019), and World Spider Catalog (2023).

Other sources have also been used for the correct identification and are, when used, signed in the species remarks.

## RESULTS

In total, up to date, 180 taxa, belonging to 130 genera and 33 families, are listed in the study area; among these, 15 species and 6 genera are reported for the first time for Sicily (one of which is also new for Italy).

The species of opiliones sampled are 3 belonging to 3 genera and 3 different families.

In the checklist below the new records are marked by (\*), of which the collection data, additional biological, literature, chorotype and distribution information are also provided.

### *Checklist*

Ordo ARANEAE Clerck, 1757

Familia AGELENIDAE C.L. Koch, 1837

*Agelena* Walckenaer, 1805

*Agelena labyrinthica* (Clerck, 1757)

*Aterigena* Bolzern, Hänggi et Burckhardt, 2010

*Aterigena aliquoi* (Brignoli, 1971)

*Eratigena* Bolzern, Burckhardt et Hänggi, 2013

*Eratigena agrestis* (Walckenaer, 1802)

*Lycosoides* Lucas, 1846

*Lycosoides coarctata* (Dufour, 1831)  
*Lycosoides flavomaculata* Lucas, 1846

*Tegenaria* Latreille, 1804  
*Tegenaria dalmatica* Kulczyński, 1906  
*Tegenaria domestica* (Clerck, 1757)  
*Tegenaria parietina* (Fourcroy, 1785)

*Textrix* Sundevall, 1833  
*Textrix caudata* Olivier, 1789

Familia AMAUROBIIDAE Thorell, 1870

*Amaurobius* C.L. Koch, 1837  
*Amaurobius erberi* (Keyserling, 1863)

Familia ANYPHAENIDAE Bertkau, 1878

*Anyphaena* Sundevall, 1833  
*Anyphaena sabina* L. Koch, 1866

Familia ARANEIDAE Clerck, 1757

*Aculeipera* Chamberlin et Ivie, 1942  
*Aculeipera armida* (Audouin, 1826)

*Agalenatae* Archer, 1951  
*Agalenatae redii* (Scopoli, 1763)

*Araneus* Clerck, 1757  
*Araneus angulatus* Clerck, 1757  
*Araneus diadematus* Clerck, 1757

*Araniella* Chamberlin et Ivie, 1942  
*Araniella cucurbitina* (Clerck, 1757)

*Argiope* Audouin, 1826  
*Argiope bruennichi* (Scopoli, 1772)  
*Argiope lobata* (Pallas, 1772)

*Gibbaranea* Archer, 1951  
*Gibbaranea bituberculata* (Walckenaer, 1802)

*Hypsosinga* Ausserer, 1871  
*Hypsosinga sanguinea* (C.L. Koch, 1844)

*Larinoides* Caporiacco, 1934  
*Larinoides cornutus* (Clerck, 1757)

*Lipocrea* Thorell, 1878

<i>Lipocrea epeiroides</i> (O. Pickard-Cambridge, 1872)	<i>Dysdera lagreciae</i> Alicata, 1964
<i>Mangora</i> O. Pickard-Cambridge, 1889	<i>Harpactea</i> Bristowe, 1939
<i>Mangora acalypha</i> (Walckenaer, 1802)	<i>Harpactea sicula</i> Alicata, 1966
<i>Neoscona</i> Simon, 1864	<i>Parachtes</i> Alicata, 1964
<i>Neoscona byzanthina</i> (Pavesi, 1876)	<i>Parachtes siculus</i> (Caporiacco, 1949)
<i>Neoscona subfuscata</i> (C.L. Koch, 1837)	Familia FILISTATIDAE Ausserer, 1867
<i>Nuctenea</i> Simon, 1864	<i>Filistata</i> Latreille, 1810
<i>Nuctenea umbratica</i> (Clerck, 1757)	<i>Filistata insidiatrix</i> (Forsskål, 1775)
<i>Zilla</i> C.L. Koch, 1836	<i>Pritha</i> Lehtinen, 1967
<i>Zilla diodia</i> (Walckenaer, 1802)	* <i>Pritha nana</i> (Simon, 1868)
<i>Zygiella</i> F.O. Pickard-Cambridge, 1902	Familia GNAPHOSIDAE Pocock, 1898
<i>Zygiella x-notata</i> (Clerck, 1757)	<i>Aphantaulax</i> Simon, 1878
Familia CHEIRACANTHIIDAE Wagner, 1887	<i>Aphantaulax trifasciata</i> (O. Pickard-Cambridge, 1872)
<i>Cheiracanthium</i> C.L. Koch, 1839	<i>Civizelotes</i> Senglet, 2012
<i>Cheiracanthium angulitarse</i> Simon, 1878	<i>Civizelotes</i> sp.
<i>Cheiracanthium erraticum</i> (Walckenaer, 1802)	<i>Drassodes</i> Westring, 1851
<i>Cheiracanthium mildei</i> L. Koch, 1864	<i>Drassodes lapidosus</i> (Walckenaer, 1802)
Familia CLUBIONIDAE Wagner, 1887	<i>Drassodes lutescens</i> (C.L. Koch, 1839)
<i>Clubiona</i> Latreille, 1804	<i>Gnaphosa</i> Latreille, 1804
<i>Clubiona corticalis</i> (Walckenaer, 1802)	<i>Gnaphosa lucifuga</i> (Walckenaer, 1802)
<i>Porrhoclubiona</i> Lohmander, 1944	<i>Haplodrassus</i> Chamberlin, 1922
<i>Porrhoclubiona leucaspis</i> (Simon, 1932)	<i>Haplodrassus kulczynskii</i> Lohmander, 1942
<i>Porrhoclubiona vegeta</i> (Simon, 1918)	<i>Haplodrassus rufipes</i> (Lucas, 1846)
Familia DICTYNIDAE O. Pickard-Cambridge, 1871	<i>Haplodrassus signifer</i> (C.L. Koch, 1839)
<i>Argenna</i> Thorell, 1870	<i>Leptodrassus</i> Simon, 1878
<i>Argenna</i> sp.	<i>Leptodrassus</i> sp.
<i>Brigittea</i> Lehtinen, 1967	<i>Marinarozelotes</i> Ponomarev, 2020
<i>Brigittea civica</i> (Lucas, 1850)	<i>Marinarozelotes cf. barbatus</i> (L. Koch, 1866)
<i>Nigma</i> Lehtinen, 1967	<i>Marinarozelotes bardiae</i> (Caporiacco, 1928)
<i>Nigma puella</i> (Simon, 1870)	<i>Marinarozelotes fuscipes</i> (L. Koch, 1866)
<i>Nigma walckenaeri</i> (Roewer, 1951)	<i>Micaria</i> Westring, 1851
Familia DYSDERIDAE C.L.Koch, 1837	* <i>Micaria coarctata</i> (Lucas, 1846)
<i>Dysdera</i> Latreille, 1804	* <i>Micaria dives</i> (Lucas, 1846)
<i>Dysdera crocata</i> C.L. Koch, 1838	<i>Nomisia</i> Dalmas, 1921

- Nomisia exornata* (C.L. Koch, 1839)
- Poecilochroa Westring, 1874*
- Poecilochroa albomaculata* (Lucas, 1846)
- \**Poecilochroa loricata* Kritscher, 1996
- Scotophaeus Simon, 1893*
- Scotophaeus blackwallii* (Thorell, 1871)
- Scotophaeus validus* (Lucas, 1846)
- Setaphis Simon, 1893*
- Setaphis carmeli* (O. Pickard-Cambridge, 1872)
- Zelotes Gistel, 1848*
- Zelotes atrocaeruleus* (Simon, 1878)
- Zelotes femellus* (L. Koch, 1866)
- Familia LINYPHIIDAE Blackwall, 1859
- Agyneta Hull, 1911*
- Agyneta rurestris* (C.L. Koch, 1836)
- \**Bolyphantes* C. L. Koch, 1837
- \**Bolyphantes luteolus* (Blackwall, 1833)
- Centromerus Dahl, 1886*
- \**Centromerus arcanus* (O. Pickard-Cambridge, 1873)
- Frontinellina* van Helsdingen, 1969
- Frontinellina fruterorum* (C.L. Koch, 1834)
- Gnathonarium Karsc, 1881*
- Gnathonarium dentatum* (Wider, 1834)
- \**Leptophantes* Menge, 1866
- \**Leptophantes minutus* (Blackwall, 1833)
- Linyphia Latreille, 1804*
- \**Linyphia mimonti* Simon, 1885
- Linyphia triangularis* (Clerck, 1757)
- \**Megaleptophantes* Wunderlich, 1994
- \**Megaleptophantes collinus* (L.Koch, 1872)
- Microlinyphia Gerhardt, 1928*
- Microlinyphia pusilla* (Sundevall, 1830)
- Neriene Blackwall, 1833*
- Neriene clathrata* (Sundevall, 1830)
- Pelecopsis Simon, 1864*
- Pelecopsis bucephala* (O. Pickard-Cambridge, 1875)
- Tenuiphantes Saaristo et Tavasevitch, 1996*
- Tenuiphantes herbicola* (Simon, 1884)
- Tenuiphantes tenuis* (Blackwall, 1852)
- \**Walckenaeria* Blackwall, 1833
- \**Walckenaeria corniculans* (O. Pickard-Cambridge, 1875)
- \**Walckenaeria monoceros* (Wider, 1834)
- Familia LIOCRANIDAE Simon, 1897
- \**Agroeca Westring, 1861*
- \**Agroeca* sp.
- Mesiotelus Simon, 1897*
- Mesiotelus mauritanicus* Simon, 1909
- \**Scotina* Menge, 1873
- \**Scotina celans* (Blackwall, 1841)
- Familia LYCOSIDAE Sundevall, 1833
- Alopecosa Simon, 1885*
- Alopecosa albofasciata* (Brullé, 1832)
- Hogna Simon, 1885*
- Hogna radiata* (Latreille, 1817)
- Lycosa Latreille, 1804*
- Lycosa tarantula* (Linnaeus, 1758)
- Pardosa* C.L.Koch, 1847
- Pardosa proxima* (C.L. Koch, 1847)
- Trabea* Simon, 1876
- Trabea paradoxa* Simon, 1876
- MIMETIDAE Simon, 1881
- Mimetus* Hentz, 1832
- Mimetus laevigatus* (Keyserling, 1863)
- Familia MITURGIDAE Simon, 1886
- Prochora* Simon, 1886
- Prochora lycosiformis* (O. Pickard-Cambridge, 1872)

- Zora* C.L. Koch, 1847  
*Zora manicata* Simon, 1878
- Familia OECOBIIDAE Blackwall, 1862
- Oecobius* Lucas, 1846  
*Oecobius maculatus* Simon, 1870
- Familia OXYOPIDAE Thorell, 1870
- Oxyopes* Latreille, 1870  
*Oxyopes heterophthalmus* (Latreille, 1804)  
*Oxyopes lineatus* Latreille, 1806  
*Oxyopes ramosus* (Martini et Goeze, 1778)
- Familia PALPIMANIDAE Thorell, 1870
- Palpimanus* Dufour, 1820  
*Palpimanus gibbulus* Dufour, 1820
- Familia PHIODROMIDAE Thorell, 1870
- Philodromus* Walckenaer, 1826  
*Philodromus aureolus* (Clerck, 1757)  
*Philodromus dispar* Walckenaer, 1826  
*Philodromus emarginatus* (Schrank, 1803)
- Pulchellodromus* Wunderlich, 2012  
*Pulchellodromus bistigma* (Simon, 1870)
- Rhysodromus* Schick, 1965  
*Rhysodromus* sp.
- Thanatus* C.L. Koch, 1837  
*Thanatus vulgaris* Simon, 1870
- Familia PHOLCIDAE C.L. Koch, 1850
- Holocnemus* Simon, 1875  
*Holocnemus pluchei* (Scopoli, 1763)
- Familia PHRUROLITHIDAE Banks, 1892
- Phrurolithus* C.L. Koch, 1839  
*Phrurolithus festivus* (C.L. Koch, 1835)
- Familia PISAURIDAE Simon, 1890
- Pisaura* Simon, 1886  
*Pisaura mirabilis* (Clerck, 1757)
- Familia SALTICIDAE Blackwall, 1841
- Ballus* C.L. Koch, 1850  
*Ballus armadillo* (Simon, 1817)  
*Ballus rufipes* (Simon, 1868)
- Cyrba* Simon, 1876  
*Cyrba algerina* (Lucas, 1846)
- Euophrys* C.L. Koch, 1834  
*Euophrys rufibarbis* (Simon, 1868)
- Evarcha* Simon, 1902  
*Evarcha jucunda* (Lucas, 1846)
- Heliophanus* C.L. Koch, 1833  
*Heliophanus apiatus* Simon, 1868  
*Heliophanus melinus* L. Koch, 1867
- Icius* Simon, 1876  
*Icius hamatus* (C.L. Koch, 1846)
- Leptorchestes* Thorell, 1870  
*Leptorchestes mutilloides* (Lucas, 1846)
- Menemerus* Simon, 1868  
*Menemerus semilimbatus* (Hahn, 1829)
- Pellenes* Simon, 1876  
*Pellenes brevis* (Simon, 1868)  
*Pellenes seriatus* (Thorell, 1875)
- Philaeus* Thorell, 1869  
*Philaeus chrysops* (Poda, 1761)
- Phlegra* Simon, 1876  
*Phlegra bresnieri* (Lucas, 1846)
- Pseudeuophrys* Dahl, 1912  
*Pseudeuophrys lanigera* (Simon, 1871)
- Pseudicius* Simon, 1885  
*Pseudicius encarpatus* (Walckenaer, 1802)
- Salticus* Latreille, 1804  
*Salticus scenicus* (Clerck, 1757)  
*Salticus zebraneus* (C.L. Koch, 1837)
- Synageles* Simon, 1876  
*Synageles albotrimaculatus* (Lucas, 1846)  
*\*Synageles venator* (Lucas, 1836)

Familia SCYTODIDAE Blackwall, 1864

*Scytodes* Latreille, 1804

*Scytodes thoracica* (Latreille, 1802)

Familia SEGESTRIIDAE

*Segestria* Simon, 1893

*Segestria florentina* (Rossi, 1790)

*Segestria senoculata* (Linnaeus, 1758)

Familia SICARIIDAE Keyserling, 1880

*Loxosceles* Heineken et Lowe, 1832

*Loxosceles rufescens* (Dufour, 1820)

Familia SPARASSIDAE Bertkau, 1872

*Micrommata* Latreille, 1804

*Micrommata ligurina* (C.L.Koch, 1845)

*Olios* Walckenaer, 1837

*Olios argelasius* (Walckenaer, 1806)

*Tibellus* Simon, 1875

*Tibellus* cfr. *maritimus* (Menge, 1875)

Familia TETRAGNATHIDAE Menge, 1866

*Metellina* Chamberlin et Ivie, 1941

*Metellina mengei* (Blackwall, 1870)

*Metellina merianae* (Scopoli, 1763)

*Pachygnatha* Sundevall, 1823

*Pachygnatha degeeri* Sundevall, 1830

*Tetragnatha* Latreille, 1804

*Tetragnatha extensa* (Linnaeus, 1758)

*Tetragnatha obtusa* C.L. Koch, 1837

Familia THERIDIIDAE Sundevall, 1833

*Anelosimus* Simon, 1891

*Anelosimus vittatus* (C.L. Koch, 1836)

*Argyrodes* Simon, 1864

*Argyrodes argyrodes* (Walckenaer, 1834)

*Asagena* Sundevall, 1833

*Asagena phalerata* (Panzer, 1801)

*Enoplognatha* Pavesi, 1880

\**Enoplognatha afrodite* Hippa et Oksala, 1983

*Enoplognatha mandibularis* (Lucas, 1846)

*Episinus* Walckenaer, 1809

*Episinus angulatus* (Blackwall, 1836)

*Episinus maculipes* Cavanna, 1876

*Euryopis* Menge, 1868

*Euryopis episinoidea* (Walckenaer, 1847)

*Kochiura* Archer, 1950

*Kochiura aulica* (C.L. Koch, 1838)

*Neottiura* Menge, 1868

*Neottiura uncinata* (Lucas, 1846)

*Paidiscura* Archer, 1950

*Paidiscura* sp.

*Platnickina* Koçak et Kemal, 2008

*Platnickina tincta* (Walckenaer, 1802)

*Simitidion* Wunderlich, 1992

*Simitidion simile* (C.L. Koch, 1836)

*Steatoda* Sundevall, 1833

*Steatoda albomaculata* (De Geer, 1778)

*Steatoda bipunctata* (Linnaeus, 1758)

*Steatoda grossa* (C.L. Koch, 1838)

*Steatoda paykulliana* (Walckenaer, 1806)

*Steatoda triangulosa* (Walckenaer, 1802)

*Theridion* Walckenaer, 1805

*Theridion melanurum* Hahn, 1831

*Theridion mystaceum* L. Koch, 1870

*Theridion pictum* (Walckenaer, 1802)

*Theridion varians* Hahn, 1833

Familia THOMISIDAE Sundevall, 1833

*Bassaniodes* Pocock, 1903

*Bassaniodes bufo* (Dufour, 1820)

*Heriaeus* Simon, 1875

*Heriaeus hirtus* (Latreille, 1819)

*Leptorchestes* Thorell, 1870

- Leptorcheses mtilloides* (Lucas, 1846)

*Misumena* Latreille, 1804  
*Misumena vatia* (Clerck, 1757)

*Monaeses* Thorell, 1869  
*Monaeses paradoxus* (Lucas, 1846)

*Ozyptila* Simon, 1864  
*Ozyptila atomaria* (Panzer, 1801)  
*Ozyptila confluens* (C.L. Koch, 1845)

*Runcinia* Simon, 1875  
*Runcinia grammica* (C.L. Koch, 1837)

*Synema* Simon, 1864  
*Synema globosum* (Fabricius, 1775)

*Thomisus* Walckenaer, 1805  
*Thomisus onustus* Walckenaer, 1805

*Xysticus* C.L. Koch, 1835  
*Xysticus acerbus* Thorell, 1872  
*Xysticus audax* (Schrank, 1803)  
*Xysticus kochi* Thorell, 1872  
*Xysticus nubilus* Simon, 1875

Familia TITANOECIDAE Lehtinen, 1967

*Titanoeca* Thorell, 1870  
*Titanoeca tristis* L. Koch, 1872

Familia ULOBORIDAE Thorell, 1869

*Uloborus* Latreille, 1806  
*Uloborus plumipes* Lucas, 1846  
*Uloborus walckenaerius* Latreille, 1806

Familia ZODARIIDAE Thorell, 1881

*Zodarion* Walckenaer, 1826  
*Zodarion elegans* (Simon, 1873)

Familia ZOROPSIDAE Bertkau, 1882

*Zoropsis* Simon, 1878  
*Zoropsis spinimana* (Dufour, 1820)

Ordo OPILIONES Sundevall, 1833

Familia NEMASTOMATIDAE Simon, 1872

*Histicostoma Kratochvil, 1958*  
*Histicostoma argenteolumulatum* (Canestrini, 1875)

Familia PHALANGIIDAE Latreille, 1802

*Opilio* Herbst, 1798  
*Opilio canestrinii* (Thorell, 1876)

Familia TROGULIDAE Sundevall, 1833

*Anelasmococephalus* Simon, 1879  
*Anelasmococephalus calcaneatus* Martens et Chemici, 1988

## OBSERVATIONS

***Pritha nana*** (Simon, 1868) (Figs. 1, 2)

MATERIAL EXAMINED. ITALY • 1 male and 6 females; Sicily, Troina (Enna); 15 Nov. 2020; legit F.C. Amata; live specimens.

DISTRIBUTION. Mediterannean, India.

REMARKS. New species for Sicily. The specimens were collected under rocks, often found in numbers greater than two under the same rock.

***Micaria coarctata*** (Lucas, 1846) (Fig. 3)

MATERIAL EXAMINED. ITALY • 1 male; Sicily, Troina (Enna); 2 Feb. 2021; legit F.C. Amata; live specimen.

DISTRIBUTION. Mediterranean, Eastern Europe, Caucasus, Russia (Europe to Far East), Iran, Kazakhstan, Central Asia.

REMARKS. New species for Sicily. The specimen was collected under rocks.

***Micaria dives*** (Lucas, 1846) (Fig. 4)

MATERIAL EXAMINED. ITALY • 1 male and 1 female; Sicily, Troina (Enna); 2 Feb. 2021; legit F.C. Amata; live specimens.

DISTRIBUTION. Europe, Turkey, Israel, Caucasus, Russia (Europe to Far East), Iran, Central Asia, India, China, Korea, Japan.

REMARKS. New species for Sicily. The specimens were collected under rocks.

***Poecilochroa loricata* Kristscher, 1996**

MATERIAL EXAMINED. ITALY • 1 male and 1 female; Sicily, Troina (Enna); 2 Feb. 2021; legit F.C. Amata; live specimens.

DISTRIBUTION. Malta.

REMARKS. New species for Sicily and first observation for Italy. To date the species was only known from Malta (Kristscher, 1996; Pflieger et al., 2017), the taxonomic identification follows Bauer (2022).

The specimens were collected under rocks.

***Bolyphantes luteolus* (Blackwall, 1833)**

MATERIAL EXAMINED. ITALY • 1 male; Sicily, Troina (Enna); 12 May 2021; legit F.C. Amata; live specimen.

DISTRIBUTION. Europe, Russia (Europe to South Siberia), China.

REMARKS. New species and new genus for Sicily. The specimen was sampled on its web, built on low vegetation.

***Centromerus arcanus* (O. Pickard-Cambridge, 1873)**

MATERIAL EXAMINED. ITALY • 1 female; Sicily, Troina (Enna); 23 Dec. 2020; legit F.C. Amata; live specimen.

DISTRIBUTION. Greenland, Europe, Russia (Europe to Middle Siberia).

REMARKS. New species for Sicily. For a more certain identification, the specimen was compared with some of the species *C. serratus* (O. Pickard-Cambridge, 1875), already known for Sicily.



Figure 1. Male *Pritha nana*. Figure 2. Female *Pritha nana*.

Figure 3. *Micaria coarctata*. Figure 4. *Micaria dives*.

***Lepthyphantes minutus* (Blackwall, 1833)**

MATERIAL EXAMINED. ITALY • 1 male and 1 female; Sicily, Troina (Enna); 23 Dec. 2020; legit F.C. Amata; live specimens.

DISTRIBUTION. Europe.

REMARKS. New species and new genus for Sicily.

The specimen was collected on its web, built on low vegetation.

***Linyphia mimonti* Simon, 1885**

MATERIAL EXAMINED. ITALY • 3 females; Sicily, Troina (Enna); 9 Feb. 2021; legit F.C. Amata; live specimens.

DISTRIBUTION. Italy, Albania, Greece (incl. Crete), Lebanon, Israel.

REMARKS. New species for Sicily.

The specimen was collected on its web, built on low vegetation.

***Megalepthyphantes collinus* (L. Koch, 1872)**

MATERIAL EXAMINED. ITALY • 1 male; Sicily, Troina (Enna); 23 Dec. 2020; legit F.C. Amata; live specimen.

DISTRIBUTION. Europe, Turkey.

REMARKS. New genus and new species for Sicily.

The specimens were collected wandering on low vegetation.

***Walckenaeria corniculans* (O. Pickard-Cambridge, 1875)**

MATERIAL EXAMINED. ITALY • 1 male and 1



Figure 5. *Poecilochroa loricata*. Figure 6. *Bolyphantes luteolus*.  
Figure 7. *Centromerus arcanus*. Figure 8. *Lepthyphantes minutus*.

female; Sicily, Troina (Enna); 6 Feb. 2021; legit F.C. Amata; live specimens.

DISTRIBUTION. Europe, North Africa, Turkey.

REMARKS. New genus and new species for Sicily.

The specimens were collected on low vegetation.

***Walckenaeria monoceros* (Wider, 1834)**

MATERIAL EXAMINED. ITALY • 1 male and 1 female; Sicily, Troina (Enna); 6 Feb. 2021; legit F.C. Amata; live specimens.

DISTRIBUTION. Europe, Azerbaijan, Central Asia.

REMARKS. New genus and new species for Sicily.

The specimens were collected under a rock.

***Agroeca* sp.**

MATERIAL EXAMINED. ITALY • 2 nymphs; Sicily, Troina (Enna); 3 Jan. 2021; legit F.C. Amata; live specimens.

DISTRIBUTION (of the genus). Subcosmopolitan.

REMARKS. New genus for Sicily. Having managed to sample only two juvenile specimens, the authors limit themselves to pointing out the new genus for the island. With further collection campaigns in other periods they will clarify the species to which the specimens belong.

***Scotina celans* (Blackwall, 1841)**

MATERIAL EXAMINED. ITALY • 1 female; Sicily, Troina (Enna); 20 May 2021; legit F.C. Amata; live specimen.

DISTRIBUTION. Europe, Algeria.



Figure 9. *Linyphia mimonti*. Figure 10. *Megalepthyphantes collinus*.  
Figure 11. *Walckenaeria corniculans*. Figure 12. *Walckenaeria monoceros*.



Figure 13. *Agroeca* sp. Figure 14. *Scotina celans*.  
Figure 15. *Synageles venator*. Figure 16. *Enoplognatha afrodite*.

REMARKS. New species and new genus for Sicily.

The specimen was sampled while wandering on the ground.

#### *Synageles venator* (Lucas, 1836)

MATERIAL EXAMINED. ITALY • 1 female; Sicily, Troina (Enna); 16 Jul. 2021; legit F.C. Amata; live specimen.

DISTRIBUTION. Europe, Turkey, Caucasus, Russia (Europe to Far East), Kazakhstan, Central Asia, China, Japan. Introduced to Canada.

REMARKS. New species for Sicily. The specimen was sampled wandering on the ground.

#### *Enoplognatha afrodite* Hippa et Oksala, 1983

MATERIAL EXAMINED. ITALY • 1 male; Sicily,

Troina (Enna); 14 Jun. 2021; legit F.C. Amata; live specimen.

DISTRIBUTION. Southern Europe.

REMARKS. New species for Sicily. The specimen was sampled in its web, built on the branches of an olive tree.

#### ACKNOWLEDGEMENTS

We want to thank our families for their constant support.

#### REFERENCES

- Alicata P., 1964. Le *Dysdera* del gruppo andreinii della fauna italiana (Araneae, Dysderidae). Atti dell'Acca-

- demia Gioenia di Scienze Naturali in Catania, 6: 67–86.
- Alicata P., 1966. Le *Harpactea* (Araneae, Dysderidae) della fauna italiana e considerazioni sulla loro origine. Atti dell'Accademia Gioenia di Scienze Naturali in Catania, 18: 190–221.
- Bauer T., 2022. Remarks on the type material of *Poecilochroa loricata* Krötscher, 1996 (Araneae: Gnaphosidae). Zootaxa, 5169: 97–100. <https://doi.org/10.11646/zootaxa.5169.1.10>
- Brignoli P.M., 1968. Su alcuni Araneidae e Theridiidae di Sicilia (Araneae). Atti dell'Accademia Gioenia di Scienze Naturali in Catania serie 6, 20: 85–104.
- Brignoli P.M., 1977. Ragni d'Italia XXVII. Nuovi dati su Agelenidae, Argyronetidae, Hahniidae, Oxyopidae e Pisauridae, cavernicoli ed epigei (Araneae). Quaderni del Museo di Speleologia "V. Rivera" L'Aquila, 2: 3–118.
- Cantarella T., 1974. Contributo alla conoscenza degli *Heiliophanus* (Arachnida, Araneae, Salticidae) di Sicilia. Animalia, 1: 157–173.
- Cantarella T., 1980. Contributo alla conoscenza dei Salticidi (Arachnida, Araneae) di Sicilia. Animalia, 7: 55–68.
- Chemini C., 1995. Arachnida Scorpionida, Palpigradi, Solifugae, Opiliones. In: Minelli A., Ruffo S. & La Posta S. (Eds.), Checklist delle specie della fauna italiana, 2: 8 S. Calderini, Bologna.
- Dentici A. & Amata F.C., 2018. New faunistic data for the Sicilian Aracnofauna (Arachnida Araneae). Biodiversity Journal, 9: 271–276. <https://doi.org/10.31396/Biodiv.Jour.2018.9.3.271.276>
- Dentici A. & Amata F.C., 2021. *Prodidomus amaranthinus* (Lucas, 1846) (Araneae Gnaphosidae) new for the Italian araneofauna. Biodiversity Journal, 12: 357–359. <https://doi.org/10.31396/Biodiv.Jour.2021.12.2.357.359>
- Dentici A., Galasso P. & Barbera A., 2021. First record of *Trichoncus aurantiipes* Simon, 1884 (Araneae Linyphiidae) in Italy. Biodiversity Journal, 12: 125–128. <https://doi.org/10.31396/Biodiv.Jour.2021.12.1.125.128>
- Dentici A. & Galasso P., 2021. Checklist of spiders and Opiliones (Arachnida Araneae, Opiliones) of Sicilian southwest swamp lakes "Pantano Cuba", "Pantano Longarini" and "Pantano Bruno" (Italy). Fifth contribution to knowledge of Sicilian spider fauna Biodiversity Journal, 21: 695–703. <https://doi.org/10.31396/Biodiv.Jour.2021.12.3.695.703>
- Dentici A., Barbera A., Ditta A. & Surdo S., 2022. On some new reports on the spider fauna of Italy and Sicily (Arachnida Araneae). Biodiversity Journal, 13: 399–408. <https://doi.org/10.31396/Biodiv.Jour.2022.13.2.399.408>
- Dentici A., 2022. Araneofauna (Arachnida Araneae) of Mount Pellegrino (Sicily, Italy). Fourth contribution to knowledge of the Sicilian spider fauna. Biodiversity Journal, 13: 641–650. <https://doi.org/10.31396/Biodiv.Jour.2022.13.3.641.650>
- Hansen H., 2005. Arachnida Araneae Salticidae + CDROM. In: Ruffo S. & Stoch F. (eds) 2005, Checklist e distribuzione della fauna italiana. Memorie del Museo civico di Storia naturale di Verona, 2 Serie, Sezione Scienze della Vita, 16: 77–78.
- Levi H.W., 1966. The Care of Alcoholic Collections of Small Invertebrates. Systematic Zoology, 15: 183–188.
- Lugetti G. & Tongiorgi P., 1969. Ricerche sul genere *Alopecosa* Simon (Araneae, Lycosidae). Atti della Società Toscana di Scienze Naturali residente in Pisa. Memorie, Serie B, 76: 1–100.
- Krötzer E., 1996. Ein Beitrag zur Kenntnis der Spinnen-Fauna der Maltesischen Inseln (Chelicerata: Araneae). Annalen des Naturhistorischen Museums in Wien, Serie B, 98: 117–156.
- Marcellino I., 1970. Su alcuni Opilioni (Arachnida) della Sicilia sud-orientale e centrale. Bollettino delle sedute dell'Accademia Gioenia di Scienze Naturali, 10: 283–308.
- Marcellino I., 1973. Opilioni delle isole Eolie ed Egadi. Lavori della Società Italiana di Biogeografia, 3: 327–339.
- Marcellino I., 1974. Nuovi dati sugli opilioni (Arachnida) di Sicilia e di altre isole del Mediterraneo. Animalia, 1: 185–200.
- Marcellino I., 1975. Considérations biogéographiques sur les Opilions de Sicile. In: Vlijm L. (Ed.), Proceedings of the 6th International Arachnological Congress. Amsterdam: Nederlandse Entomologische Vereniging, pp. 222–226.
- Martens J. & Chemini C., 1988. Die Gattung *Anelasmoccephalus* Simon 1879. Biogeographie, Artgrenzen und Biospezies-Konzept (Opiliones: Troglidae). Zoologische Jahrbücher Abteilung für Systematik, Geographie und Biologie der Tiere, 115: 1–48.
- Nentwig W., Blick T., Bosmans R., Gloor D., Hänggi A. & Kropf C., 2023. Spiders of Europe. Version 4.2023. Online at <https://www.araneae.nmbe.ch>, accessed on 16.Apr.2023. <https://doi.org/10.24436/1>
- Pantini P. & Isaia M., 2019. Araneae.it: the online Catalog of Italian spiders with addenda on other Arachnid Orders occurring in Italy (Arachnida: Araneae, Opiliones, Palpigradi, Pseudoscorpionida, Scorpiones, Solifugae). Fragmenta Entomologica, 51: 127–152. Online at [www.araneae.it](http://www.araneae.it), accessed on 16.Apr.2023.
- Pflieger W.A., Schönhofen A., Niedbala W., Vella P., Sciberras A. & Vella A., 2017. New records of mites

- (Acari) and harvestmen (Opiliones) from Malta with a preliminary checklist of Maltese Arachnida. *Soil Organisms*, 89: 85–110.
- Roberts M.J., 1995. Spiders of Britain and Northern Europe. Harper Collins, London, 383 pp.
- Trotta A., 2004. Introduzione ai Ragni italiani (Arachnida Araneae). *Memorie della Società Entomologica Italiana*, 83: 3–178.
- World Spider Catalog, 2023. World Spider Catalog. Version 23.0. Natural History Museum Bern, online at <http://wsc.nmbe.ch>, accessed on 16.Apr.2023.  
<https://doi.org/10.24436/2>