

Checklist of Norwegian spiders (Arachnida: Araneae), including Svalbard and Jan Mayen

Kjetil Aakra & Erling Hauge

Aakra, K. & Hauge, E. 2003. Checklist of Norwegian spiders (Arachnida: Araneae), including Svalbard and Jan Mayen. Norw. J. Entomol. 50, 109–129.

A new checklist of Norwegian spiders, including the Svalbard archipelago and the island of Jan Mayen, is presented along with taxonomic amendments and updates to the previous checklist. A total of 562 species are currently known from the Norwegian Mainland, 17 from Svalbard and five from Jan Mayen.

Keywords: Araneae, Checklist, Norway.

Kjetil Aakra, Midt-Troms Museum, Box 82, N-9059, Storsteinnes, Norway. kjetil.aakra@midt-troms.museum.no

Erling Hauge, Zoological Museum, University of Bergen, Norway.

INTRODUCTION

Since the publication of the last Norwegian checklist (Hauge 1989), there has been tremendous changes in the systematics and taxonomy of spiders. Also, many new species of spiders have been reported from Norway since then. A preliminary updated checklist with distribution maps has been available on the Internet (Aakra & Hauge 2002) and a new and updated checklist based upon the net presentation is given below. Separate checklists of spiders from the Svalbard archipelago and the remote Mid-Atlantic island of Jan Mayen are also included.

In addition to the checklists a bibliography of literature of Norwegian spiders is given, again separately for the Norwegian Mainland and the Svalbard archipelago/Jan Mayen. These bibliographies contain all known papers where records of species from the respective regions are provided.

METHODS

The present list follow Platnick (2003) in regard to taxonomy and nomenclature, with one noteable exception. We agree with Saaristo (1973) and

Saaristo & Koponen (1998) in that *Meioneta* Hull, 1920 is a junior synonym of *Agyrleta* Hull, 1911. Although this has been accepted by several authors the former genus continues to be used in the aforementioned catalogue. Accepting the arguments of Saaristo (1973) and Roberts (1987), all members of «*Meioneta*» are here listed under *Agyrleta*.

Several other taxonomic changes have been incorporated into this new checklist compared to Hauge (1989). The most important are,

- *Meta* and *Metellina* transferred to the Tetragnathidae (Metidae not recognised).
- *Zygiella* transferred to the Araneidae (Piel & Nutt 1997, Scharff & Coddington 1997).
- *Leptophantes* split into several new genera (Saaristo & Tanasevitch 1996, 1999).
- *Argyroneta aquatica* transferred to the Cybaeidae (Grothendieck & Kraus 1994, also see Platnick 2003).
- *Cheiracanthium* spp. transferred to the Miturgidae (Ramírez et al. 1997).
- *Cicurina cicur* (Fabricius, 1793) and *Masti-*

gusa arietina (Thorell, 1871) transferred from the Agelenidae to the Dictynidae (see Lethinen 1967 and Platnick 2003).

- *Cryphoeca silvicola* transferred from the Agelenidae to the Hahniidae (see Platnick 2003).
- The family name Eusparassidae (and Heteropodidae) has been changed to Sparassidae (Jäger 1999).

The first author has recently revised some of the numerous linyphiid species described by Embrik Strand (see Aakra 2002) that were included in the previous checklist (Hauge 1989). The following taxonomic changes have been made,

- *Ceratinella oculatissima* Strand, 1901 = *C. wideri* (Thorell, 1871) (first and only record of this species in Norway)
- *Cnephalocotes dentiger* Strand, 1902 = *Silometopus reussi* (Thopprell, 1871)
- *Cnephalocotes ophthalmicus* Strand, 1901 = *Silometopus ambiguus* (O.P.-Cambridge, 1905) (older name suppressed for lack of usage)
- *Pseudogonatum fuscomarginatum* Strand, 1901 = *Zornella cultrigera* (L. Koch. 1879).

Other synonyms now included in the list are,

Araneidae

- *Araneus sinister* (Thorell, 1873) = *A. saevus* (L. Koch, 1872) (see Tullgren, 1952).

Gnaphosidae

- *Gnaphosa intermedia* Holm, 1939 = *G. sticta* Kulczyn'ski, 1908 (Ovtsharenko & Marusik 1988).
- *Zelotes serotinus* (L. Koch, 1866) = *Z. longipes* (L. Koch, 1866) (see Tullgren 1946).

Linyphiidae

- *Centromerus granulosus* (L. Koch, 1879) = *Macrargus multesimus* (O. P.-Cambridge, 1875) (see Holm, 1945).
- *Certainopsis pectinata* (Tullgren, 1955) = *Troxochrota scabra* Kulczynski, 1894 (see Lethinen et al. 1979).

- *Diplocentria replicata* Holm, 1950 = *Wabasso quaestio* (Chamberlin, 1948) (see Millidge 1984).
- *Diplostyla colletti* (Strand, 1899) = *Kaestneria pullata* (O. P.-Cambridge, 1863) (see Eskov & Marusik 1994).
- *Leptyphantes zebrinus* (Menge, 1866) = *Improphanthes decolor* (Westring, 1861) (see Kronestedt 1975).
- *Meioneta beata* (O. P-Cambridge, 1906) = *Agyneta affinis* (Kulczynski, 1898) (see Wunderlich 1973).
- *Oedothorax recurvus* (Strand, 1901) = *Zornella cultrigera* (L. Koch, 1879) (see Holm, 1944).
- *Silometopus laesus* (L. Koch, 1879) = *S. reussi* (Thorell, 1871) (see Denis 1949).

Lycosidae

- *Pirata piccolo* Dahl, 1908 = *P. insularis* Emerton, 1885 (see Hackman 1954).

Thomisidae

- *Xysticus augur* Strand, 1900 = *X. cristatus* (Clerck, 1757) (Tambs-Lyche 1942).
- *Xysticus austerus* L. Koch, 1879 = *X. obscurus* Collett, 1877 (Holm 1973).
- *Xysticus norvegicus* Strand, 1900 = *X. bifasciatus* C. L. Koch, 1837 (see Tambs-Lyche 1942).
- *Xysticus sexangulatus* Strand, 1900 = *X. cristatus* (Clerck, 1757) (Tambs-Lyche 1942).

Salticidae

- *Attulus cinereus* (Westring, 1861) = *Sitticus distinguendus* (Simon, 1868) (see Platnick 2003).
- *Ballus depressus* (Walckenaer, 1802) = *Ballus chalybeius* (Walckenaer, 1802) (see Platnick 2003).

Strand (1900, 1904) described a comparatively large number of *Gnaphosa* spp. (*G. porrecta*, *G. pseudolapponica*, *G. limbata*, *G. nordlandica* and

G. norvegica) that were also included in the previous checklist. Preliminary examination of the types of these by the first author have revealed that they all belong to common and widespread species, or in the case of *G. limbata* (the type of which is lost) must be considered a nomen dubium (Aakra in prep.). They are therefore omitted from the present list. Formal synonymization will be provided in a forthcoming paper.

The Norwegian record of *Dysdera crocota* Cl. L. Koch, 1835 (Dysderidae) mentioned by Bristowe (1939) is highly unlikely and the species has been omitted from the checklist.

The species *Theridion melanurum* Hahn, 1831 has also been removed from the list. Although specimen(s) could not be located in Norwegian collections it is highly likely that this species has been confused with the similar *T. mystaceum* L. Koch, 1870 which has been found in Norway (see Aakra 2000).

The lycosid *Pardosa arenicola* (O.P.-Cambridge, 1875) has been removed from the current checklist. This name probably refers to the subspecies *P. agrestis purbeckensis* F. O. P.-Cambridge, 1895, the taxonomic status of which is rather uncertain. It probably warrants species status (T. Blick pers. comm.). Specimens resembling *P. a. purbeckensis* has been found in both the Oslojord area and in the vicinity of Trondheim (Aakra unpub. data) but the status of this taxon will have to await further study. For the time being only the subspecies is listed from Norway.

Also omitted is *Pardosa thorelli* (Collett, 1876), even if a specimen has not been found and examined. Given the paucity of material since the origi-

nal description we find it best to treat this species as a nomen dubium until the type is located. However, surviving original drawings of the epigyne of the Holotype indicates that *P. thorelli* belongs to a species of *Acantholycosa*, as also mentioned by Tambs-Lyche (1940) and Holm (1950). A formal synonymization will be presented in a later paper.

Kronestedt (1990) split the species *Alopecosa aculeata* (Clerck 1757) into two distinct species, i.e. the parent species *A. aculeata* and the long overlooked *A. taeniata* (C. L. Koch, 1835). The latter was reported from Norway by Kronestedt (1990) and has since been found in other Norwegian material (Hauge unpub. data).

Diplocephalus semiglobosus (Westring, 1861) is a junior synonym (T. Kronestedt pers. comm.) and has been deleted from the present list.

The salticid *Neon reticulatus* (Blackwall, 1853) was inadvertently omitted from the last checklist.

The latest addition to the list is by Aakra & Olsen (2003) which reported *Ostearius melanopygius* (O.P.-Cambridge, 1879) (Linyphiidae) and *Uloborus plumipes* Lucas, 1846 (Uloboridae).

Other details regarding nomenclatorial changes (transfers to other genera, etc.) compared to Hauge (1989) are given in the checklist where relevant. All families and species are presented in alphabetical order. New species discovered since the previous checklist are marked with *.

The total number and number of species from each family recorded from Norway is shown in table 1.

Table 1. Total number of species and number of species in each family.

Family	No. of species	Family	No. of species	Family	No. of species	Family	No. of species
Agelenidae	4	Hahniidae	7	Oxyopidae	1	Theridiidae	37
Amaurobiidae	3	Linyphiidae	248	Philodromidae	16	Thomisidae	20
Anyphaenidae	1	Liocranidae	10	Pholcidae	2	Titanococcidae	1
Araneidae	33	Lycosidae	50	Pisauridae	2	Uloboridae	2
Clubionidae	18	Mimetidae	1	Salticidae	30	Zoridae	2
Cybaeidae	1	Miturgidae	3	Segestriidae	1		
Dictynidae	8	Nesticidae	1	Sparassidae	1		
Gnaphosidae	45	Oonopidae	1	Tetragnathidae	13		
Total species: 562							

CHECKLIST-NORWEGIAN MAINLAND

Agelenidae

- Agelena labyrinthica* (Clerck, 1757)
Tegenaria atrica C. L. Koch, 1843
T. domestica (Clerck, 1757)
Textrix denticulata (Olivier, 1789)

Amaurobiidae

- Amaurobius fenestralis* (Stroem, 1768)
A. similis (Blackwall, 1861)
Arctobius agelenoides Emerton, 1919*

Anyphaenidae

- Anyphaena accentuata* (Walckenaer, 1802)

Araneidae

- Aculepeira ceropegia* (Walckenaer, 1802)
Agelenata redii (Scopoli, 1763)
Araneus alsine (Walckenaer, 1802)
A. angulatus Clerck, 1757
A. diadematus Clerck, 1757
A. marmoreus Clerck, 1757
A. nordmanni (Thorell, 1870)
A. quadratus Clerck, 1757
A. saevus (L. Koch, 1872)
A. sturmi (Hahn, 1833)-Formerly *Atea s.* (see Levi 1991)
Araniella alpica (L. Koch, 1869)
A. cucurbitina (Clerck, 1757)
A. displicata (Hentz, 1847)
A. opistographa (Kulczynski, 1905)
Cercidia prominens (Westring, 1851)
Cyclosa conica (Pallas, 1772)
Gibbaranea bituberculata (Walckenaer, 1802)
G. omoeda (Thorell, 1870)
Hypsosinga albovittata (Westring, 1851)
H. pygmaea (Sundevall, 1831)
H. sanguinea (C. L. Koch, 1844)
Larinoides cornutus (Clerck, 1757)
L. patagiatus (Clerck 1757)

L. sclopetarius (Clerck, 1757)

Mangora acalypha (Walckenaer 1802)

Neoscona adianta (Walckenaer, 1802)

Nuctenea silvicoltrix (C. L. Koch, 1844)-Formerly *Araneus s.*

N. umbratica (Clerck, 1757)

Singa hamata (Clerck, 1757)

S. nitidula C. L. Koch, 1844*

Zygiella atrica (C. L. Koch, 1845)

Z. stroemi (Thorell, 1870)

Z. x-notata (Clerck, 1757)

Clubionidae

Clubiona brevipes Blackwall, 1841

C. caerulescens L. Koch, 1867

C. comta C. L. Koch, 1839

C. diversa O. P.-Cambridge, 1862

C. frisia Wunderlich & Schuett, 1995*-Norwegian specimens previously reported as *C. similis* L. Koch, 1867 belong to this species (Aakra unpub. data, also see Wunderlich & Schuett 1995).

C. frutetorum L. Koch, 1866

C. germanica Thorell, 1871

C. kulczynskii Lessert, 1905

C. lutescens Westring, 1851

C. neglecta O. P.-Cambridge, 1862

C. norvegica Strand, 1900

C. pallidula (Clerck, 1757)

C. phragmitis C. L. Koch, 1843

C. reclusa O. P.-Cambridge, 1863

C. stagnatilis Kulczynski, 1897

C. subsultans Thorell, 1875

C. terrestris Westring, 1851

C. trivialis L. Koch, 1843

Cybaeidae

Argyroneta aquatica (Clerck, 1757)

Dictynidae

Archaeodictyna consecuta (O. P.-Cambridge, 1872)-Formerly *Dictyna c.* (see Lethinen 1967).

- Argenna subnigra* (O. P.-Cambridge, 1861)
Cicurina cicur (Fabricius, 1793)
Dictyna arundinacea (Linnaeus, 1758)
D. latens (Fabricius, 1775)
D. pusilla Thorell, 1856
D. uncinata Thorell, 1856
Mastigusa arietina (Thorell, 1871)
- Gnaphosidae**
- Callilepis nocturna* (Linnaeus, 1758)
Drassodes cupreus (Blackwall, 1834)
D. lapidosus (Walckenaer, 1802)
D. pubescens (Thorell, 1856)
D. villosus (Thorell, 1856)
Drassyllus lutetianus (L. Koch, 1866)
D. praeficus (L. Koch, 1866)
D. pumilus (C. L. Koch, 1839)
D. pusillus (C. L. Koch, 1833)
Echemus angustifrons (Westring, 1861)*
Gnaphosa bicolor (Hahn, 1833)
G. lapponum (L. Koch, 1866)
G. leporina (L. Koch, 1866)
G. lucifuga (Walckenaer, 1802)
G. microps Holm, 1939
G. montana (L. Koch, 1866)
G. muscorum (L. Koch, 1866)
G. orites Chamberlin, 1922
G. sticta Holm, 1939
G. intermedia Holm, 1939
Haplodrassus cognatus (Westring, 1861)
H. minor (O. P.-Cambridge, 1879)
H. moderatus (Kulczynski, 1897)
H. signifer (C. L. Koch, 1839)
H. soerrenseni (Strand, 1900)
H. sylvestris (Blackwall, 1833)
H. umbratilis (L. Koch, 1866)
Micaria aenea Thorell, 1871
M. alpina L. Koch, 1872
M. formicaria (Sundevall, 1832)
M. fulgens (Walckenaer, 1802)

- M. nivosa* L. Koch, 1866
M. pulicaria (Sundevall, 1832)
M. silesiaca L. Koch, 1875
M. subopaca Westring, 1861
Poecilochroa variana (C. L. Koch, 1839)
Scotophaeus blackwalli (Thorell, 1873)
S. quadripunctatus (Linnaeus, 1758)
Zelotes clivicola (L. Koch, 1870)
Z. electus (C. L. Koch, 1839)
Z. latreillei (Simon, 1878)
Z. longipes (L. Koch, 1866)
Z. petrensis (C. L. Koch, 1839)
Z. puritanus Chamberlin, 1922*
Z. subterraneus (C. L. Koch, 1833)

Hahniidae

- Antistea elegans* (Blackwall, 1841)
Cryphoeca silvicola (C. L. Koch, 1834)
Hahnia helveola Simon, 1875
H. montana (Blackwall, 1841)
H. nava (Blackwall, 1841)
H. ononidum Simon, 1875
H. pusilla C. L. Koch, 1841

Linyphiidae

- Abacoproces saltuum* (L. Koch, 1872)
Abisko abiskoensis (Holm, 1945)*
Agnyphantes expunctus (O.P.-Cambridge, 1875)-
 Formerly *Leptophantes e.* (see Saaristo & Tanasevitch 1996 and Platnick 2003).
Agyneta affinis (Kulczynski, 1898)
A. cauta (O. P.-Cambridge, 1902)
A. conigera (O. P.-Cambridge, 1863)
A. decora (O. P.-Cambridge, 1870)
A. fuscipalpus (C.L.Koch, 1836)
A. gulosa (L.Koch, 1869)
A. innotabilis (O.P.-Cambridge, 1863)
A. mossica (Schikora 1993)*
A. nigripes (Simon, 1884)
A. olivacea (Emerton, 1882)
A. ramosa Jackson, 1912

- A. rurestris* (C.L.Koch, 1836)
- A. saxatilis* (Blackwall, 1844)
- A. similis* (Kulczynski, 1926)
- A. subtilis* (O. P-Cambridge, 1863)
- A. suecica* Holm, 1950-Probably a synonym of
A. arietans (O.P.-Cambridge, 1872) (Palmgren
1975).
- Allomenga scopigera* (Grube, 1859)
- A. vidua* (L.Koch, 1879)*
- Anguliphantes angulipalpis* (Westring, 1851)-Formerly *Lepthyphantes a.* (Saaristo & Tanasevitch 1996).
- Araeoncus crassiceps* (Westring, 1861)
- Asthenargus paganus* (Simon, 1884)
- Baryphyma trifrons* (O. P.-Cambridge, 1863)-Transferred to *Minyrioloides* by Tanasevitch (1990), but not accepted by Platnick (2003).
- Bathyphantes approximatus* (O.P.-Cambridge, 1871)
- B. gracilis* (Blackwall, 1841)
- B. nigrinus* (Westring, 1851)
- B. parvulus* (Westring, 1851)
- B. setiger* F.O. P-Cambridge, 1894
- B. similis* Kulczynski, 1894-Formerly *B. norvegicus* Strand, 1902 (Holm 1987, contra Holm 1944 (?), see Hauge 1989)
- Bolephthyphantes index* (Thorell, 1856)-Formerly *Bolyphantes i.* (see Saaristo & Tanasevitch 1996, 2000 and Platnick 2003)
- Bolyphantes alticeps* (Sundevall, 1833)
- B. luteolus* (Blackwall, 1833)
- Caviphantes saxetorum* (Hull, 1916)*
- Centromerita bicolor* (Blackwall, 1833)-Transfer to *Centromerus* (Saaristo & Tanasevitch 1996) not accepted by Platnick (2003).
- C. concinna* (Thorell, 1875)-Transfer to *Centromerus* (Saaristo & Tanasevitch 1996) not accepted by Platnick (2003).
- Centromerus brevivulvatus* Dahl, 1912-Formerly *C. aequalis* (C. L. Koch, 1841)(see Platnick 2003)
- C. arcarius* (O.P.-Cambridge, 1873)
- C. dilutus* (O. P-Cambridge, 1875)
- C. incillum* (L.Koch, 1881)
- C. pabulator* (O. P-Cambridge, 1875)
- C. prudens* (O. P-Cambridge, 1873)
- C. sylvaticus* (Blackwall, 1841)
- Ceratinella brevipes* (Westring, 1851)
- C. brevis* (Wider, 1834)
- C. scabrosa* (O. P-Cambridge, 1871)
- Ceratinella wideri* (Thorell, 1871)*-Formerly *C. oculatissima* (Strand, 1901) (see Palmgren 1976 and Aakra 2002).
- Cnephalocotes obscurus* (Blackwall, 1834)
- Collinsia holmgreni* (Thorell, 1871)
- C. inerrans* (L.Koch, 1879)-Formerly *Milleriana i.* (O.P.-Cambridge, 1885) (see Marusik et al. 1993 and Platnick 2003).
- C. spetsbergensis* (Thorell, 1871)
- Decipiphantes decipiens* (L.Koch, 1879)-Formerly *Lepthyphantes d.* (see Saaristo & Tanasevitch 1996).
- Dicymbium nigrum* (Blackwall 1834)-Specimens have not been checked for *D. n. brevisetosum* Locket, 1962.
- D. tibiale* (Blackwall, 1836)
- Diplocentria bidentata* (Emerton, 1882)
- D. rectangulata* (Emerton, 1915)-Formerly *Microrcentria pusilla* (Schenkel, 1925)
- Diplocephalus cristatus* (Blackwall, 1833)
- D. latifrons* (O.P.-Cambridge, 1863)
- D. permixtus* (O.P.-Cambridge, 1871)
- D. picinus* (Blackwall, 1841)
- Diplostyla concolor* (Wider, 1834)
- Dismodicus bifrons* (Blackwall, 1841)
- D. elevatus* (C.L.Koch, 1838)
- Drapetisca socialis* (Sundevall, 1833)
- Drepanotylus uncatus* (O.P.-Cambridge, 1873)
- Entelecara acuminata* (Wider, 1834)
- E. congenera* (O.P.-Cambridge, 1879)
- E. erythropus* (Westring, 1851)
- E. flavipes* (Blackwall, 1834)*

- E. media* Kulczynski, 1887
Erigone arctica (White, 1852)
E. atra Blackwall, 1833
E. dentigera O.P.-Cambridge, 1874—Formerly *E. capra* Simon, 1884 (see Eskov 1994).
E. dentipalpis (Wider, 1834)
E. longipalpis (Sundevall, 1830)
E. promiscua (O.P.-Cambridge, 1872)
E. psychrophila Thorell, 1872
E. tirolensis L.Koch, 1872
Erigonella hiemalis (Blackwall, 1841)
E. ignobilis (O.P.-Cambridge, 1871)
Evansia merens O.P.-Cambridge, 1900
Floronia bucculenta (Clerck, 1757)
Gnathonarium dentatum (Wider, 1834)
Gonatium paradoxum (L.Koch, 1869)
G. rubellum (Blackwall, 1841)
G. rubens (Blackwall, 1833)
Gongylidiellum latebricola (O.P.-Cambridge, 1871)
G. murcidum Simon, 1884
G. vivum (O.P.-Cambridge, 1875)
Gongylidium rufipes (Linnaeus, 1758)
Halorates reprobus (O.P.-Cambridge, 1879)
Helophora insignis (Blackwall, 1841)
Hilaira excisa (O.P.-Cambridge, 1871)
H. frigida (Thorell, 1872)
H. herniosa (Thorell, 1875)
H. nubigena Hull, 1911
H. pervicax Hull, 1908
Hypomma bituberculatum (Wider, 1834)
H. cornutum (Blackwall, 1833)
Hypselistes jacksoni (O.P.-Cambridge, 1902)
Improphanes complicatus (Emerton, 1882)—Formerly *Leptyphantes c.* (see Saaristo & Tanasevitch 1996).
I. decolor (Westring, 1861)—Formerly *Leptyphantes d.* (Saaristo & Tanasevitch 1996).
Incestophantes crucifer (Menge, 1866)—Transferred from *Bolyphantes* to *Leptyphantes* by Thaler et al. (1994), transferred to *Inceto-*
phantes by Saaristo & Tanasevitch (2000).
I. kochiellus (Strand, 1900)—Formerly *Leptyphantes k.* (see Saaristo & Tanasevitch 1996).
Jacksonella falconeri (Jackson, 1908)
Kaestneria dorsalis (Wider, 1834)
K. pullata (O.P.-Cambridge, 1863)
Labulla thoracica (Wider, 1834)
Lasiargus hirsutus (Menge, 1869)
Leptyphantes antronensis Schenkel, 1933
L. ericaeus (Blackwall, 1853)
L. leprosus (Ohlert, 1867)
L. minutus (Blackwall, 1833)
L. pallidus (O.P.-Cambridge, 1871)
Leptorhoptrum robustum (Westring, 1851)
Leptothrix hardyi (Blackwall, 1850)
Linyphia hortensis Sundevall, 1830
L. tenuipalpis Simon, 1884*
L. triangularis (Clerck, 1757)
Lophomma punctatum (Blackwall, 1841)
Macrargus boreus Holm, 1968
M. carpenteri (O.P.-Cambridge, 1894)
M. multesimus (O.P.-Cambridge, 1875)
M. rufus (Wider, 1834)
Maro lehtineni Saaristo, 1971
M. lepidus Casimir, 1961*
M. minutus O.P.-Cambridge, 1906
M. sublestus Falconer, 1915
Maso sundevalli (Westring, 1851)
Mecynargus borealis (Jackson, 1930)
M. monticola (Holm, 1943)
M. morulus (O.P.-Cambridge, 1873)
M. paetus (O.P.-Cambridge, 1875)
M. sphagnicola (Holm, 1939)
Megaleptyphantes nebulosus (Sundevall, 1830)—Formerly *Leptyphantes n.* (Saaristoa & Tanasevitch 1996).
Metopobactrus prominulus (O. P.-Cambridge, 1872)
Micrargus apertus (O. P.-Cambridge, 1871)
M. herbigradus (Blackwall, 1854)

- | | |
|--|--|
| <i>M. subaequalis</i> (Westring, 1851) | <i>P. radicicola</i> (L.Koch, 1872) |
| <i>Microctenonyx subitaneus</i> (O.P.-Cambridge, 1875) | <i>Peponocranium ludicum</i> (O.P.-Cambridge, 1861) |
| <i>Microlinyphia pusilla</i> (Sundevall, 1830) | <i>Pityohyphantes phrygianus</i> (C.L.Koch, 1836) |
| <i>Microneta viaria</i> (Blackwall, 1841) | <i>Pocadicnemis pumila</i> (Blackwall, 1841) |
| <i>Minicia marginella</i> (Wider, 1834) | <i>Poeciloneta variegata</i> (Blackwall, 1841)-Formerly <i>P. globosa</i> (Wider, 1834). |
| <i>Minyriolus pusillus</i> (Wider, 1834) | |
| <i>Mioxena blanda</i> (Simon, 1884) | <i>Porrhomma campbelli</i> F.O.P.-Cambridge, 1894 |
| <i>Moebelia penicillata</i> (Westring, 1851) | <i>P. convexum</i> (Westring, 1851) |
| <i>Monocephalus castaneipes</i> (Simon, 1884) | <i>P. lativelum</i> Tretzel, 1956 |
| <i>Mughiphantes suffusus</i> (Strand, 1901)-Formerly <i>Leptphyphantes s.</i> (Saaristo & Tanasevitch 1999). | <i>P. microphthalmum</i> (O.P.-Cambridge, 1871)* |
| <i>M. whymperi</i> (F.O.P.-Cambridge, 1894)-Formerly <i>Leptphyphantes w.</i> (Saaristo & Tanasevitch 1996) | <i>P. montanum</i> Jackson, 1913 |
| <i>Neriene clathrata</i> (Sundevall, 1830) | <i>P. oblitum</i> (O. P.-Cxambridge, 1871) |
| <i>N. montana</i> (Clerck, 1757) | <i>P. pallidum</i> Jackson, 1913 |
| <i>N. peltata</i> (Wider, 1834) | <i>P. pygmaeum</i> (Blackwall, 1834) |
| <i>N. radiata</i> (Walckenaer, 1841) | <i>Satilatlas britteni</i> (Jackson, 1913) |
| <i>Notioscopus sarcinatus</i> (O.P.-Cambridge, 1872) | <i>Saaristoa abnormis</i> (Blackwall, 1841) |
| <i>Obscuriphantes obscurus</i> (Blackwall, 1841)-Formerly <i>Leptphyphantes o.</i> (Saaristo & Tanasevitch 2000). | <i>S. firma</i> (O.P.-Cambridge, 1905)* |
| <i>Oedothorax agrestis</i> (Blackwall, 1853) | <i>Savignia frontata</i> Blackwall, 1833 |
| <i>O. apicatus</i> (Blackwall, 1850) | <i>Scandichrestus tenuis</i> (Holm, 1943)-Formerly <i>Typhochrestus t.</i> (Wunderlich 1995). |
| <i>O. fuscus</i> (Blackwall, 1834) | <i>Scotinotylus clavatus</i> (Schenkel, 1927)-Probably senior synonym of <i>S. sacer</i> (Crosby, 1929) (see Thaler 1970). |
| <i>O. gibbosus</i> (Blackwall, 1841) | <i>S. evansi</i> (O.P.-Cambridge, 1894) |
| <i>O. retusus</i> (Westring, 1851) | <i>Semljicola angulatus</i> (Holm, 1963)-Formerly <i>Eboria a.</i> (see Eskov & Marusik 1994). |
| <i>Oreonetides vaginatus</i> (Thorell, 1872) | <i>S. faustus</i> (O.P.-Cambridge, 1900)-Formerly <i>Latthorax f.</i> (see Saaristo & Eskov 1996). |
| <i>Oryphantes angulatus</i> (O.P.-Cambridge, 1881)-Formerly <i>Leptphyphantes a.</i> (Saaristo & Tanasevitch 1996) | <i>S. latus</i> (Holm, 1939)-Formerly <i>Latthorax l.</i> (see Saaristo & Eskov 1996). |
| <i>Ostearius melanopygius</i> (O..-Cambridge, 1879)-Synantropic | <i>S. lapponicus</i> (Holm, 1939)* |
| <i>Panamomops mengei</i> Simon, 1926 | <i>Silometopus ambiguus</i> (O.P.-Cambridge, 1905) |
| <i>Parapelecopsis nemoralis</i> (Blackwall, 1841)-Formerly <i>Pelecopsis n.</i> (Wunderlich 1992). | <i>S. elegans</i> (O.P.-Cambridge, 1872) |
| <i>Pelecopsis elongata</i> (Wider, 1834) | <i>S. incurvatus</i> (O.P.-Cambridge, 1873) |
| <i>P. mengei</i> (Simon, 1884) | <i>S. reussi</i> (Thorell, 1871) |
| <i>P. parallela</i> (Wider, 1834) | <i>Sintula corniger</i> (Blackwall, 1856) |

- Ceratinopsis* s. (see Wunderlich 1970 and Märsik & Tanasevitch 1998).
- Syedra gracilis* (Menge, 1869)
- Tallusia experta* (O.P.-Cambridge, 1871)
- Tapinocyba insecta* (L.Koch, 1869)
- T. pallens* (O.P.-Cambridge, 1872)
- Tapinocyboides pygmaeus* (Menge, 1869)
- Tapinopa longidens* (Wider, 1834)
- Taranucnus setosus* (O.P.-Cambridge, 1863)
- Tenuiphantes alacris* (Blackwall, 1853)-Formerly *Leptyphantes a.* (Saaristo & Tanasevitch 1996).
- T. cristatus* (Menge, 1866)-Formerly *Leptyphantes c.* (see Saaristo & Tanasevitch 1996).
- T. flavipes* (Blackwall, 1854)-Formerly *Leptyphantes f.* (see Sdaaristo & Tanasevitch 1996).
- T. mengei* (Kulczynski, 1887)-Formerly *Leptyphantes m.* (Saaristo & Tanasevitch 1996).
- T. nigriventris* (L.Koch, 1879)-Formerly *Leptyphantes n.* (Saaristo & Tanasevitch 1996).
- T. tenebricola* (Wider, 1834)-Formerly *Leptyphantes t.* (see Saaristo & Tanasevitch 1996).
- T. tenuis* (Blackwall, 1852)-Formerly *Leptyphantes t.* (Saaristo & Tanasevitch 1996).
- T. zimmermanni* (Bertkau, 1890)-Formerly *Leptyphantes z.* (Saaristo & Tanasevitch 1996).
- Thyreosthenius biovatus* (O.P.-Cambridge, 1875)
- T. parasiticus* (Westring, 1851)
- Tiso aestivus* (L.Koch, 1872)
- T. vagans* (Blackwall, 1834)
- Tmeticus affinis* (Blackwall, 1855)
- T. nigriceps* (Kulczynski, 1916)
- Trichoncus vasconicus* Denis, 1944
- Troxochrota scabra* Kulczynski, 1894
- Troxochrus nasutus* Schenkel, 1925
- T. scabriculus* (Westring, 1851)
- Typhochrestus digitatus* (O.P.-Cambridge, 1872)
- T. sylviae* Hauge, 1968
- Wabasso quaestio* (Chamberlin, 1948)
- Walckenaeria acuminata* Blackwall, 1833
- W. antica* (Wider, 1834)
- W. atrotibialis* (O.P.-Cambridge, 1878)
- W. capito* (Westring, 1861)
- W. clavicornis* (Emerton, 1882)
- W. cucullata* (C.L.Koch, 1836)
- W. cuspidata* (Blackwall, 1833)
- W. dysderoides* (Wider, 1834)
- W. inflexa* (Westring, 1861) (?)-Doubtful species (L. J. Jonsson pers. comm.), but accepted by Platnick (2003).
- W. karpinskii* (O.P.-Cambridge, 1873)
- W. kochi* (O. P.-Cambridge, 1872)
- W. mitrata* (Menge, 1868)
- W. monoceros* (Wider, 1834)
- W. nodosa* O. P.-Cambridge, 1873
- W. nudipalpis* (Westring, 1851)
- W. obtusa* Blackwall, 1836
- W. unicornis* O. P.-Cambridge, 1861
- W. vigilax* (Blackwall, 1853)
- Zornella cultrigera* (L.Koch, 1879)
- ### Liocranidae
- Agroeca brunnea* (Blackwall, 1833)
- A. cuprea* Menge, 1873*
- A. proxima* (O. P.-Cambridge, 1871)
- Apostenus fuscus* Westring, 1851
- Liocranum rupicola* (Walckenaer, 1830)
- Phrurolithus festivus* (C. L. Koch, 1835)
- P. minimus* C. L. Koch, 1839
- Scotina celans* (Blackwall, 1841)
- S. gracilipes* (Blackwall, 1859)
- S. palliardi* (L. Koch, 1881)
- ### Lycosidae
- Acantholycosa lignaria* (Clerck, 1757)
- Acantholycosa norvegica* (Thorell, 1872)
- Alopecosa aculeata* (Clerck, 1757)
- A. barbipes* (Sundevall, 1833)
- A. cuneata* (Clerck, 1757)
- A. fabrilis* (Clerck, 1757)

- A. inquilina* (Clerck, 1757)
A. pinetorum (Thorell, 1856)
A. pulverulenta (Clerck, 1757)
A. taeniata (C. L. Koch, 1835)*
A. trabalis (Clerck, 1757)
Arctosa alpigena (Doleschall, 1852)-Formerly
 Tricca a. (see Dondale & Redner 1983).
A. cinerea (Fabricius, 1777)
A. leopardus (Sundevall, 1833)
A. lutetiana (Simon, 1876)*
A. perita (Latreille, 1799)
A. stigmosa (Thorell, 1875)*
Hygrolycosa rubrofasciata (Ohlert, 1865)
Pardosa agrestis (O. P.-Cambridge, 1875)
P. agrestis purbeckensis F. O. P.-Cambridge,
 1895*-Formerly *P. arenicola* (O. P.-Cam-
 bridge, 1875)
P. agricola (Thorell, 1856)
P. amentata (Clerck, 1757)
P. atrata (Thorell, 1873)
P. eiseni (Thorell, 1875)
P. fulvipes (Collett, 1876)
P. hyperborea (Thorell, 1872)
P. lapponica (Thorell, 1872)
P. lasciva L. Koch, 1879*
P. lugubris (Walckenaer, 1802) senso latto-
 Material has not yet been checked for speci-
 mens of *P. alacris* (C. L. Koch, 1833) or *P.
saltans* Töpfer-Hofman , 2000 (see Töpfer-
 Hofman et al. 2000).
P. monticola (Clerck, 1757)
P. nigriceps (Thorell, 1856)-Material has not yet
 been checked for specimens of *P. maisa* Hippa
 & Mannila, 1982 (see Hippa & Manilla 1982).
P. paludicola (Clerck, 1757)
P. palustris (Linnaeus, 1758)
P. prativaga (C. L. Koch, 1870)
P. pullata (Clerck, 1757)
P. riparia (C. L. Koch, 1833)
P. schenkeli Lessert, 1904
P. septentrionalis (Westring, 1861)

- P. sphagnicola* (Dahl, 1908)
P. trailli (O. P.-Cambridge, 1873)
Pirata hygrophilus Thorell, 1872
P. insularis Emerton, 1885
P. piraticus (Clerck, 1757)
P. piscatorius (Clerck, 1757)
P. uliginosus (Thorell, 1856)
Trochosa ruricola (De Geer, 1778)
T. spinipalpis (F. O. P.-Cambridge, 1895)
T. terricola Thorell, 1856.
Xerolycosa miniata (C. L. Koch, 1834).
X. nemoralis (Westring, 1861)

Mimetidae

- Ero furcata* (Villers, 1789)

Miturgidae

- Cheiracanthium erraticum* (Walckenaer, 1802)
C. oncognathum Thorell, 1871
C. virescens (Sundevall, 1833)

Nesticidae

- Nesticus cellularis* (Clerck, 1757)

Oonopidae

- Oonops pulcher* Templeton, 1835

Oxyopidae

- Oxyopes ramosus* (Martini & Goeze, 1879)*-
Doubtful record, only reported by Collett (1876),
specimen(s) apparently lost.

Philodromidae

- Philodromus aureolus* (Clerck, 1757)
P. cespitum (Walckenaer, 1802)
P. collinus C. L. Koch, 1835
P. dispar Walckenaer, 1826
P. emarginatus (Schrank, 1803)
P. fuscomarginatus De Geer, 1778
P. histrio (Latreille, 1819)*
P. margaritatus (Clerck, 1757)
P. rufus Walckenaer, 1825

Thanatus arcticus Thorell, 1872

T. arenarius Thorell, 1872

T. atratus Simon, 1875

T. formicinus (Clerck, 1757)

T. striatus C. L. Koch, 1845

Tibellus maritimus (Menge, 1875)

T. oblongus (Walckenaer, 1802)

Pholcidae

Psilochorus simoni (Berland, 1911)

Pisauridae

Dolomedes fimbriatus (Clerck, 1757)

Pisaura mirabilis (Clerck, 1757)

Salticidae

Aelurillus v-insignitus (Clerck, 1757)

Ballus chalybeius (Walckenaer, 1802)

Bianor aurocinctus (Ohlert, 1865)

Dendryphantes hastatus (Clerck, 1757)

D. rufus (Sundevall, 1832)

Euophrys frontalis (Walckenaer, 1802)*

Evarcha arcuata (Clerck, 1757)

E. falcata (Clerck, 1757)

Heliophanus cupreus (Walckenaer, 1802)

H. dampfi Schenkel, 1923

H. dubius C. L. Koch, 1835

H. flavigipes (Hahn, 1832)

Marpissa muscosa (Clerck, 1757)

Myrmarachne formicaria (De Geer, 1778)*

Neon reticulatus (Blackwall, 1853)

Pellenes tripunctatus (Walckenaer, 1802)

Phlegra fasciata (Hahn, 1826)

Pseudeuophrys erratica (Walckenaer, 1826)-

Formerly *Euophrys e.* (see Zabka 1997)

Salticus cingulatus (Panzer, 1797)

S. scenicus (Clerck, 1757)

S. zebraneus (C. L. Koch, 1837)

Sitticus distinguendus (Simon, 1868)

S. floridula (C. L. Koch, 1837)

S. inexpectus Logunov & Kronestedt, 1997* -specimens previously reported as *S. rupicola* (C. L. Koch, 1837) probably belong to this species (see Kronestedt & Logunov 1997).

S. pubescens (Fabricius, 1775)

S. saltator (Simon, 1868)-Formerly *Attulus s.* (see Platnick 2003)

S. terebratus (Clerck, 1757)

Talavera aequipes (O. P.-Cambridge, 1871)* -Formerly *Euophrys a.* (see Logunov 1992)

T. petrensis (C. L. Koch, 1837)

T. thorelli (Kulczynski, 1891)*

Segestriidae

Segestria senoculata (Linnaeus, 1758)

Sparassidae

Micrommata virescens (Clerck, 1757)

Tetragnathidae

Meta menardi (Latrelle, 1804)

Metellina mengei (Blackwall, 1869)

M. merianae (Scopoli, 1763)

M. segmentata (Clerck, 1757)

Pachygnatha clercki Sundevall, 1823

P. degeeri Sundevall, 1830

P. listeri Sundevall, 1830

Tetragnatha dearmita Thorell, 1873

T. extensa (Linnaeus, 1758)

T. montana Simon, 1874

T. obtusa C. L. Koch, 1837

T. pinicola L. Koch, 1870

T. striata L. Koch, 1862

Theridiidae

Achaearanea lunata (Clerck, 1757)

A. ohlerti Thorell, 1870-Fomerly *Theridion o.* (see Heimer & Nentwig 1991).

Achaearanea riparia (Blackwall, 1834)*

A. tepidariorum (C. L. Koch, 1841)-Probably synanthropic

Anelosimus vittatus (C. L. Koch, 1836)

- Crustulina guttata* (Wider, 1834)
Dipoena inornata (O. P.-Cambridge, 1861)
D. melanogaster (C. L. Koch, 1837)
D. torva (Thorell, 1871)*
Enoplognatha ovata (Clerck, 1757)
E. thoracica (Hahn, 1833)
Episinus angulatus (Blackwall, 1853)
E. truncatus Latreille, 1809
Euryopis flavomaculata (C. L. Koch, 1836)
Lasaeola tristis (Hahn, 1833)-Formerly *Dipoena t.*
Neottiura bimaculata (Linnaeus, 1767)-Formerly
 Theridion bimaculatum (see Platnick 2003)
Paidiscura pallens (Blackwall, 1834)-Formerly
 Theridion p. (see Wunderlich 1987)
Pholcomma gibbum (Westring, 1851)
Robertus arundineti (O. P.-Cambridge, 1871)
R. lividus (Blackwall, 1836)
R. lyrifer Holm, 1939
R. neglectus (O. P.-Cambridge, 1871)
R. scoticus Jackson, 1914
Rugathodes bellicosus (Simon, 1873)-Formerly
 Theridion b. (see Kronestedt 1993).
Simitidion simile (C. L. Koch, 1836)*-Formerly
 Theridion s. (see Wunderlich 1992).
Steatoda albomaculata (De Geer, 1778)
S. bipunctata (Linnaeus, 1758)
S. phalerata (Panzer, 1801)
Theonoe minutissima (O. P.-Cambridge, 1879)
Theridion impressum L. Koch, 1881
T. montanum Emerton, 1882
T. mystaceum L. Koch, 1870*-Previously listed
 as *T. melanurum* Hahn, 1831 (see Aakra 2000).
T. pictum (Walckenaer, 1802)
T. pinastri L. Koch, 1862*
T. sisyphium (Clerck, 1757)
T. tinctum (Walckenaer, 1802)
T. varians Hahn, 1833
- Thomisidae**
Coriarachne depressa (C. L. Koch, 1837)
- Diaeа dorsata* (Fabricius, 1777).
Misumena vatia (Clerck, 1757)
Ozyptila arctica Kulczynski, 1908
O. atomaria (Panzer, 1801)
O. praticola (C. L. Koch, 1837)
O. rauda Simon, 1875
O. trux (Blackwall, 1846)
Xysticus albidus Greese, 1909
X. audax (Schrank, 1803)
X. bifasciatus C. L. Koch, 1837
X. cristatus (Clerck, 1757)
X. erraticus (Blackwall, 1834)
X. kochi Thorell, 1872
X. lanio C. L. Koch, 1835
X. lineatus (Westring, 1851)
X. luctuosus (Blakcwall, 1836)
X. obscurus Collett, 1876
X. sabulosus (Hahn, 1832)
X. ulmi (Hahn, 1832)
- Titanoecidae**
Titanoeeca nivalis Simon, 1874
- Uloboridae**
Hyptiotes paradoxus (C. L. Koch, 1834)
Uloborus plumipes Lucas, 1846* -Synantropic
- Zoridae**
Zora nemoralis (Blackwall, 1861)
Z. spinimana (Sundevall, 1833)
- CHECKLIST-SVALBARD ARCHIPELAGO**
- Gnaphosidae**
Micaria eltoni (Jackson 1922)
- Hahniidae**
Hahnia helveola Simon, 1875-Probably synan-
thropic

Linyphiidae

- Agylneta nigripes* (Simon, 1884)-Formerly *Meioneta* n. (see Saaristo & Tanasevitch 1996).
- Collinsia holmgreni* (Thorell, 1871)
- C. spetsbergensis* (Thorell, 1872)
- C. thulensis* (Jackson, 1934)
- Erigone arctica palaeearctica* Braendegård, 1934
- E. psychrophila* Thorell, 1872
- E. tirolensis* L.Koch, 1872
- Hilaira glacialis* (Thorell, 1871)
- Improphanes complicatus* (Emerton, 1882)-Formerly *Lepthyphantes c.* (Saaristo & Tanasevitch 1996).
- Mecynargus borealis* (Jackson, 1930)
- Mughiphantes sobrius* (Thorell, 1872)-Formerly *Lepthyphantes s.* (Saaristo & Tanasevitch 1996).
- Oreonetides vaginatus* (Thorell, 1872)
- Tapinocyba insecta* (L.Koch, 1869)-Probably synanthropic
- Walckenaeria karpinskii* (O.P.-Cambridge, 1873)

Philodromidae

- Thanatus formicinus* (Clerck, 1757)*-Synanthropic.

CHECKLIST-JAN MAYEN**Linyphiidae**

- Agylneta nigripes* (Simon, 1884)-Formerly *Meioneta* n. (see Saaristo & Tanasevitch 1996).
- Collinsia holmgreni* (Thorell, 1871)
- Erigone tirolensis* L.Koch, 1872
- Hilaira frigida* (Thorell, 1872)
- Walckenaeria clavicornis* (Emerton, 1882)

BIBLIOGRAPHY-NORWEGIAN MAINLAND

- Aagaard, K. & Dolmen, D. 1996. Limnofauna Norvegica. 309 pp. Katalog over norsk ferskvannsfauna. Tapir forlag., Trondheim. 1996.
- Aakra, K. 1998. Epigaeic spiders of the island of Askøy, western Norway. Cand. Scient Thesis, Zoological Museum, University of Bergen, June 1998. 148 pp.
- Aakra, K. 2000a. New records of spiders (Araneae) from Norway with notes on epigynal characters of *Philodromus fuscomarginatus* (De Geer) (Philodromidae) and *Araneus sturmii* (Hahn) (Araneidae). Norw. J. Entomol. 47, 77-88.
- Aakra, K. 2000b. *Agyneta mossica* (Schikora, 1993) (Araneae, Linyphiidae) in Norway. Norw. J. Entomol. 47, 95 – 99.
- Aakra, K. 2000c. Noteworthy records of spiders (Araneae) from central regions of Norway. Norw. J. Entomol. 47, 153-162.
- Aakra, K. 2002. The riparian spider fauna (Araneae) of the river Gaula, Central Norway, implications for conservation efforts. European Arachnology 2000 (S. Toft & N. Scharff eds.). Proceedings of the 19th European Colloquium of Arachnology, Århus, 17- 22. July 2000, 243-252.
- Aakra, K. 2002. Taxonomic notes on some Norwegian linyphiid spiders described by E. Strand (Araneae, Linyphiidae). Bull. Br. Arachnol. Soc 12, 267-269.
- Aakra, K. & Dolmen, D. 2003. Distribution and ecology of the water spider, *Argyroneta aquatica* (Clerck) (Araneae, Cybaeidae), in Norway. Norw. J. Entomol. 50, 11-16.
- Aakra, K. & Djursvoll, P. 2000. Terrestre Arthropoder fra Herlandsnesjane Naturreservat på Osterøy, Hordaland. Fylkesmannen i Hordaland, Miljøvern-avdelingen. MVA-Rapport 3/2000, 1 – 27.
- Aakra, K. & Hauge, E. 2000. Provisional List of Rare and Potentially Threatened Spiders (Arachnida, Araneae) in Norway including their Proposed Red List Status. NINA Fagrapport 42, 1 – 38.
- Aakra, K. & Hauge, E. 2002. Araneae Norvegiae. Checklist and distribution maps of Norwegian spiders with Taxonomic, zoogeographic and ecological notes. <http://www.museumsnett.no/midtromsmuseum/forskningssider/norspider/index.htm>
- Aakra, K., Bretten, A. & Frengen, O. 2000. Spiders (Araneae) new to Norway. Norw. J. Entomol. 47, 149-152.
- Aakra, K., Hanssen, O. & Ødegaard, F. 2000. A collection of spiders (Araneae) from palsa bogs in the vicinity of Vardø, northern Norway. Fauna norv. 20, 43-47.
- Aakra, K. & Hauge, E. 2001. *Semljicola lapponicus* (Holm, 1939)(Araneae), new to Norway. Norw. J. Entomol. 48, 288.
- Aakra, K., Hansen, L. O. & Rinden, H. 2002. Three species of spiders (Araneae) new to Norway. Norw. J. Entomol. 49, 59-61.
- Aakra, K. & Olsen, K. M. 2003. Two introduced species of spiders (Araneae) new to Norway and the first

- record of *Uloborus plumipes* Lucas (Uloboridae) from the Faroes. Norw. J. Entomol. 48, 104.
- Alvseike, E. 1991. *Euophrys aequipes* (O. P.-Cambridge, 1871) (Araneae, Salticidae) new to Norway. Fauna Norv. Ser. B.38, 90.
- Andersen, A. 1990. Spiders in Norwegian spring barely fields and the effects of two insecticides. Norwegian Journal of Agricultural Sciences, 4, 226 – 271.
- Andersen, J. & Hanssen, O. 1994. Invertebratfaunaen på elvebredder-ett oversett element. 1. Biller (Coleoptera) ved Gaula i Sør-Trøndelag. NINA Oppdragsmelding 326, 1-23.
- Andersen, T. & Hauge, E. 1995. Pitfall catches of spiders (Araneae) from proposed Nature Reserves on Tjøme, Vestfold, SE Norway. Fauna Norv. Ser. B 42, 1-10.
- Andersen, T., Bakke, B. & Klausen, F. E. 1980. Spiders (Araneae) from five damp localities on Østerøy, Western Norway. Fauna norv. Ser. B 27, 53 – 59.
- Arneklev, J. V. 1982. Nye funn av vannedderkoppen (*Argyroneta aquatica*) i Nordland. Fauna 35, 140 – 141.
- Ashmole, N. P. & Planterose, B. R. 1979. Spides from arctic Fennoscandia. Fauna Norv. Ser. B, 26 – 30.
- Bretten, A. 1995. Effekter av vegetasjonsmanipulering på edderkoppfaunaen i lavalpin vier/dvergbjørkhei på Dovrefjell. Hovedfagsoppgave i økologi. Zooloisk institutt, Universitetet i Trondheim, Trondheim.
- Bristowe, W. S. 1939. The comity of spiders. 228 pp. Ray Society, London.
- Collett, R. 1876. Oversigt over Norges Araneida I. Forh. VidenskSelsk. Krist. 1875, 225-259.
- Collett, R. 1877. Oversigt over Norges Araneida II. Forh. VidenskSelsk. Krist. 1876, 1-27.
- Cooke, J. A. L. 1967. Spiders from Norway. Entomologists mon. Mag. 103, 12 – 13.
- Dolmen, D. 1977. Vannedderkoppen, *Argyroneta aquatica* Clerck, i Trøndelagsområdet. Fauna 30, 121 – 126.
- Dolmen, D. 1990. Ferskvannbiologiske og hydrografiske undersøkelser av Verneplan- IV-vassdrag i Trøndelag 1989. Universitetet i Trondheim. Vitenskapsmuseet. Rapport Zoologisk Serie 1990-6, 1 – 72.
- Dolmen, D. 1991. Ferskvannbiologiske og hydrografiske undersøkelser av 20 vassdrag i Møre og Romsdal 1988. Verneplan IV. Universitetet i Trondheim. Vitenskapsmuseet. Rapport Zoologisk Serie 1989-3, 1 – 105.
- Dolmen, D. 1996. Invertebrat- og amfibiefaunaen i dammer rundt Fjergen og i Tevdalen, Meråker. Vitenskapsmuseet, Rapport Zoologisk Serie 1996-1, 1 – 28.
- Duffey, E. 1995. The distribution, status and habitats of *Dolomedes fimbriatus* (Clerck) and *D. plantarius* (Clerck) in Europe. In, V. Ruzicka (Editor). Proc. 15th Eur. Colloq. Arachnol. Institute of Entomology, Ceske Budejovice, 1995, pp. 54 – 65.
- Ellefsen, G. E. & Hauge, E. 1986. Spiders (Araneae) from the Eidanger peninsula, Grenland, Telemark, SE Norway. Fauna norv. Ser. B. 33-39.
- Fabricius, J. C. 1779. Reise nach Norwegen, mit Be- merkungen aus der Naturgeschichte und Oekonomie. Hamburg.
- Folvik, E. 1992. The spider fauna of coastal sand dunes in southwestern Norway. Unpub. Cand. scient. thesis. Museum of Zoology, University of Bergen, Norway. August 1992. 65 pp.
- Folvik, E. 1994. *Porrhomma microphthalmum* (O. P.- Cambridge, 1891) (Araneae, Linyphiidae) recorded new to Norway. Fauna norv. Ser. B, 91 – 92.
- Greve, L & Hauge, E. 1989. Insekts- og edderkoppfaunaen på myrer i Hordaland. Rapport terrestrisk økologi nr. 49. Zoologisk museum, Universitet i Bergen, desember 1989. 35 pp.
- Haftorn, S. 1954. Contribution to the food biology of tits especially about storing of surplus food. Part I. The crested tit (*Parus c. cristatus* L.). Kgl. N. VidenskSel. Skr. 1953.
- Hanssen, O. & Hansen, L. O. 1998. Verneverdige insekt-habitarer. Oslofjordområdet. NINA Oppdragsmelding 546,1 – 132.
- Hauge, E. 1968. *Typhocrestus sylviae* n.sp. and *Panamomops mengei* Simon (Araneae) in Norway. Norsk ent. Tidsskr. 15, 65-69.
- Hauge, E. 1969. Six species of spiders (Araneae) new to Norway. Norsk ent. Tidsskr. 16, 1 – 8.
- Hauge, E. 1971a. Araneae from Northern Norway. Norsk ent. Tidsskr. 18, 25 – 28.
- Hauge, E. 1971b. Notes on Norwegian spiders I. Norsk ent. Tidsskr. 18, 137 – 138.
- Hauge, E. 1971c. Notes on Norwegian spiders II. Norsk ent. Tidsskr. 18, 138.
- Hauge, E. 1972a. New records of spiders from Norway. Norsk ent. Tidsskr. 19, 17 – 18.
- Hauge, E. 1972b. Notes on Norwegian spiders III. Norsk ent. Tidsskr. 19, 111.
- Hauge, E. 1972c. Spiders and Harvestmen from Møre and Romsdal and Trøndelag, Norway. Norsk ent. Tidsskr. 19, 117 – 121.
- Hauge, E. 1974a. Notes on Norwegian spiders IV. Norsk ent. Tidsskr. 21, 111 – 112.
- Hauge, E. 1974b. Notes on little known spiders in Norway. Norsk ent. Tidsskr. 21, 159 – 160.
- Hauge, E. 1975. Three species of spiders (Araneae) new to Norway. Norw. J. Entomol. 22, 81.
- Hauge, E. 1976a. Notes on eight species of spiders

- (Araneae) from the Saltfjellet area, Nordland. Norw. J. Entomol. 23, 45 – 46.
- Hauge, E. 1976b. Araneae from Finnmark, Norway. Norw. J. Entomol. 23, 121–125.
- Hauge, E. 1976c. Spiders (Araneae) of a West Norwegian *Calluna*-heath. Lindåsprosjektet, Rapp. 20, 1 – 98.
- Hauge, E. 1977. The spider fauna of two forest habitats in northern Norway. Astarte 10, 93–101.
- Hauge, E. 1979. A contribution to the knowledge to the spider fauna of Norway. Fauna norv. Ser. B, 26, 96.
- Hauge, E. 1980. A contribution to the knowledge to the spider fauna of Norway. Fauna norv. Ser. B, 27, 68 – 71.
- Hauge, E. 1986a. Spiders (Araneae) from Vestfold, South-east Norway. Fauna norv. Ser. B 33, 104 – 105.
- Hauge, E. 1986b. Notes on new and rare spiders in Norway. Fauna norv. Ser. B. 33 105–107.
- Hauge, E. 1987a. *Trichoncus vasconicus* Denis (Araneae, Linyphiidae) new to Norway. Fauna norv. Ser. B 34, 93.
- Hauge, E. 1987b. Additions to the knowledge of the Norwegian spider fauna (Araneae). Fauna norv. Ser. B 34, 94 – 95.
- Hauge, E. 1989. An annotated check-list of Norwegian spiders (Araneae). Insecta Norvegiae 4, 1–40.
- Hauge, E. 1993. Faunistical and zoogeographical notes on some Norwegian spider species (Araneae). Fauna norv. Ser. B, 40, 90 – 91.
- Hauge, E. 1998. Notes on the spatial distribution of some spider species (Araneae) in a north Norwegian birch forest. Fauna norv. Ser. A 19, 41 – 46.
- Hauge, E. 2000. Spiders (Araneae) from square samples and pitfall traps in coastal heathland, western Norway. Habitat preference, phenology and distribution. Fauna norv. 20, 31–42.
- Hauge, E., Bruvoll, A. & Solhøy, T. 2000. Epigaeic spiders (Araneae) from three habitats at Stryn, Nordfjord, Western Norway. Fauna norv. 20, 19–26.
- Hauge, E., Bruvoll, A. & Solhøy, T. 1991. Spiders (Araneae) from islands of Øygarden, West Norway. Species associations, with ecological and zoogeographical remarks. Fauna norv. Ser. B 38, 11–26.
- Hauge, E., Ellingsen, R. & Fremmersvik, M. 2002. Spiders (Araneae) from three human-influenced habitats in the Golsfjellet mountain area, central southern Norway. Fauna norv. 22, 9–14.
- Hauge, E. & Furunes, K. A. 1976. New localities for three species of spiders (Araneae) in Norway. Norw. J. Entomol. 25, 207 – 220.
- Hauge, E. & Hansen, L. O. 1991. Spiders (Araneae) from six small islands in the middle Oslofjord, SE Norway. Fauna Norv. Ser. B 38, 45–52.
- Hauge, E. & Hansen, L. O. 2000. Noteworthy records of spiders (Araneae) from the Oslofjord area, SE Norway. Norw. J. Entomol. 47, 73 – 75.
- Hauge E., Hägvar, S. & Østbye, E. 1978. Pit-fall catches of surface-active arthropods in some high mountain habitats at Finse, south Norway. III. The species of Araneida. Norw. J. Entomol. 25, 207–220.
- Hauge, E., Kauri, H. & Solhøy, T. 1998. Spiders (Araneae) from Stigstuv Hardangervidda (Norway). Fauna of Hardangervidda 20, 1 – 17.
- Hauge, E. & Kvamme, T. 1983. Spiders from forest-fire areas in southeast Norway. Fauna norv. Ser. B 30, 39–45.
- Hauge E., Meidell, B. A. & Solhøy, T. 1975. Edeløvskog på Vestlandet. I – III. Miljøverndepartementet, Rapp., 1 – 277.
- Hauge E., Meidell, B. A. & Solhøy, T. 1976. *Maro minutus* (Cbr.) (Araneae) new to Norway. Norw. J. Entomol. 23, 90.
- Hauge, E. & Midtgård, F. 1986. Spiders (Araneae) in Malaise traps from two islands in the Oslofjord, Norway. Fauna Norv. Ser. B 33, 98–102.
- Hauge, E. & Nielsen, T. R. 2001. Widening the knowledge of *Dipoena torva* (Thorell, 1875) (Araneae, Theridiidae) in Norway. Norw. J. Entomol. 48, 268.
- Hauge, E. & Otesen, P. S. 2002. Spiders (Araneae) in low alpine areas at Finse, Southern Norway. Norw. J. Entomol. 49, 17–138.
- Hauge, E. & Refseth, D. 1979. The spider fauna of 5 alpine and subalpine habitats in the Jotunheimen area, southern Norway. Fauna norv. ser. B 26, 84–90.
- Hauge, E., Syvertsen, P. O. & Rønning, A.-H. 1998. *Allomengea vidua* (L. Koch, 1879) (Araneae, Linyphiidae) in Norway, Fauna norv. Ser. B 45, 104 – 105.
- Hauge, E. & Thingstad, P. G. 1999. Spiders (Araneae) from the Kvitbergvatnet area in Saltdal, northern Norway. Norw. J. Entomol. 46, 49–56.
- Hauge, E. & Wiger, R. 1980. The spider fauna (Araneae) from 12 habitats in the Vassfaret region, south-eastern Norway. Fauna norv. Ser. B 27, 60–67.
- Helsdingen, P. J. van, Thaler, K. & deltchev, K. 1977. The *tenuis* group of Menge (Araneae, Linyphiidae). Tijdschr. Ent. 120, 1 – 69.
- Hippa, H., Koponen, S. & Mannila, R. 1984. Invertebrates of Scandinavian caves I. Araneae, Opiliones and Pseudoscorpionida (Arachnida). Ann. ent. Fenn. 50, 23 – 29.
- Holm, Å. 1944. Revision einiger norwegischen Spinnearten und Bemerkungen über deren Vorkommen in Schweden. Ent. Tidskr. 65, 122 – 134.
- Holm, Å. 1960. Notes on arctic spiders. Ark. Zool, Ser.

2. 12, 511 – 514.
- Holm, Å. 1967. Spiders (Araneae) from West-Greenland. Meddr. Grønland 184, 1 – 99.
- Holm, Å. 1973. On the spiders collected during the Swedish expeditions to Novaya Zemlya and Yenisey in 1875 and 1876. Zoologica Scr. 2, 71–110.
- Holm, Å. 1987. Några för Sverige nya spindelarter (Araneae). Ent. Tidskr. 108, 159 – 165.
- Holm, Å. & Kronestedt, T. 1970. A taxonomic study of the wolf spiders of the *Pardosa pullata*-group (Aranea, Lycosidae). Acta ent. bohem slov. 67, 408 – 428.
- Hågvar, S. & Hegstad, A. 1969. A sample of spiders (Araneida) from Svalbard. Norsk Polarinstitutt Årbok 1969, 218 – 220.
- Hågvar, E. B. & Hågvar, S. 1975. Studies on the invertebrate fauna on branches of spruce (*Picea abies* L.) during winter. Norw. J. Entomol. 22, 23 – 30.
- Jackson, A. R. 1932. Results of the Oxford expedition to lapland in 1930. Araneae and Opiliones. Proc. zool. Soc. Lond. 1932, I, 92 – 112.
- Johannessen, O. H. 1968. *Micaria decorata* Tullgren 1942 (Clubionidae, Araneae) new to Norway. Norsk ent. Tidsskr. 15, 37 – 39.
- Kauri, H. 1966. En kolleksjon Araneae and Opiliones fra Sogn. Norsk ent. Tidsskr. 13, 394 – 395.
- Klausen, F.E. 1972a. *Physocyclus simoni* Berland (Araneae, Pholcidae) new to Norway. Norsk. ent. Tidskr. 19, 123 – 125.
- Klausen, F.E. 1972b. Notes on the distribution of *Halorates reprobus* (O. P-Cambridge) (Araneae, Linyphiidae) in Norway. Norsk ent. Tidsskr. 19, 172.
- Klausen, F.E. 1974. Spiders new to Norway. Norsk ent. Tidsskr. 21, 191–194.
- Klausen, F.E. 1975. *Oonops pulcher* Templeton, 1835 (Araneae) new to Norway. Norw. J. Entomol. 22, 84.
- Klausen, F. E. 1978. En sammenligning av edderkoppfaunaen på to øyer og nærliggende fastland i Hordaland. Unpubl. Cand. Scient. Thesis. Univ. Berg. Zool. Mus. Spring 1978. 164 pp.
- Klausen, F. E. & Andersen, T. J. 1990. *Echemus angustifrons* (Westring, 1862) (Araneae, Gnaphosidae) a new spider for Norway. Fauna norv. Ser. B 37, 45–46.
- Koponen, S. 2002. Ground-living spiders in bogs in northern Europe. J. Arachnol. 30, 262–267.
- Krogerus, R. 1960. Ökologische studien über nordische Moorarthropoden. Societas Scientiarum Fennica. Commentationes Biologica XXI, 3–328.
- Kronestedt, T. 1983. Spindlar på Ölands store alvar. Ent. Tidskr. 104, 183–212.
- Kronestedt, T. 1990. Separation of two species stand-
ing as *Alopecosa aculeata* (Clerck) by morphological, behavioural and ecological characters, with remarks on related species in the *pulverulenta* group (Araneae, Lycosidae). Zoologica Scr. 19, 203–225.
- Logunov, D. V. & Kronestedt, T. 1997 A new Palearctic species of the genus *Sitticus* Simon, with notes on related species in the *floricola* group (Araneae, Salticidae). Bull. Br. arachnol. Soc. 10, 225–233.
- Martens, T. 1982. Registrering av landlevende evertebrater på Kårstø og Ognøy i Rogaland. Laboratorium for ferskvannsøkologi og innlandsfiske, Zoologisk Museum, Universitetet i Bergen. Rapport Nr. 45. 89 pp.
- Moen, A. & Moksnes, A. 1970. Forrdalsområdet – nordtrøndersk naturperle som trues av kraftutbygging. Norsk natur 6, 130 – 141.
- Moksnes, A. 1970. Trekk fra dyrelivet i Stjørdalsbygdene. Liv og lagnad i Stjørdalsbygdene, Vol. 1, pt. 1, 98 – 139.
- Moksnes, A. 1975. Dyrelivet langs vassdraget. pp. 21 – 22 i Moksnes, A. (ed.), Verneverdiene i Stjørdalsvassdraget. Nord-Trøndelag Naturvern og Sør-Trøndelag Naturvern. Trondheim. 31 pp.
- Otto, C. & Svensson, B. 1982. Structure of communities of ground-living spiders along altitudinal gradients. Holarct. Ecol. 5, 35 – 47.
- Palmgren, P. 1943. Die Spinnenfauna Finnlands. II. Pisauridae, Oxyopidae, Salticidae, Clubionidae, Anyphaenidae, Sparassidae, Ctenidae, Drassidae. Acta. Zool. Fenn. 36, 1–112.
- Palmgren, P. 1964. Några spindelprov (Arachnoidea) från Nordmarka vid Oslo. Norsk ent. Tidsskr. 12, 176 – 178.
- Palmgren, P. 1977. Die Spinnenfauna Finnlands und Ostfennoskandiens VIII. Argyronetidae, Agelenidae, Hahnidae, Dictynidae, Amaurobiidae, Titanocidae, Segestriidae, Pholcidae and Sicariidae. Fauna Fennica 30, 1–50.
- Paulsen Thingstad, G. 1982. Aktivitetsmønster og klassifisering av den markelvende edderkoppfaunaen i en subalpin fjellbjørkeskog i Sør-Trøndelag. Cand. Scient Thesis, University of Trondheim, fall 1982. 151 pp.
- Pedersen, H. C., Bretten, A., Dalen, T., Hanssen, O., Munkeby Smith, E. & Wilmann, B. 1996. Viltstell-tiltak for lirype, brenning og kutting av heivegetasjon. NINA Oppdragsmelding 283, 1 – 22.
- Platnick, N. 1975. A revision of the Holarctic spider genus *Callilepis* (Araneae, Gnaphosidae). Amer. Mus. Novitates 2573, 1 – 32.
- Pommeresche, R. 1999. Diversitet, samfunnstrukturer og habitatspreferanser hos epigiske edderkopper i ulike vegetasjonstyper innen Geitaknottane naturreservat, indre Hordaland. Cand. scient. the-

- sis. University of Bergen, spring 1999. 70 pp.
- Pommeresche, R. 2002a. Spiders (Araneae) in organically managed ley and pasture, Tingvoll farm, Norway. Norw. J. Entomol. 51–58.
- Pommeresche, R. 2002b. Spiders species and communities in bog and forest habitats in Geitaknottane nature Reserve, Western Norway. European Arachnology 2000 (S. Toft & N. Scharff eds.). Proceedings of the 19th European Colloquium of Arachnology, Århus, 17–22. July 2000, 199–205.
- Schenkel, E. 1928. Über einige Spinnen aus Lappland. Rev. suisse Zool. 35.
- Schikora, H.-B. 1995. Intraspecific variation in taxonomic characters, and notes on distribution and habitats of *Meioneta mossica* Schikora and *M. Saxatilis* (Blackwall), two closely related spiders from northern and central Europe (Araneae, Linyphiidae). Bull. Br. Arachnol. Soc. 10, 65–74.
- Simon, E. 1887. Liste des Arachnides recueillis en 1881, 1884, et 1885 par MM. J. de Guerne et C. Rabot, en Laponie (Norvege, Finlande et Russie). Bull. Soc. zool. Fr., 12, 456.
- Smith, F. P. 1912. In, Strand, E. Neue Beiträge zur Arthropodenfauna Norwegens. XIV. Nyt Mag. Naturvidensk. B. 50.
- Solem, J. O. & Hauge, E. 1973. Araneae and Opiliones in light traps at Målsjøen Sør-Trøndelag. Norsk ent. Tidsskr. 20, 275–279.
- Storm, V. 1898. Iagttagelser over Arachnider i Trondhjems omegn. K. norske Vidensk. Selsk. Skr. 1898 (7), 1–10.
- Strand, E. 1896–97. Zurt Naturgeschichte der Araneiden Norwegens. Ber. Naturw. Vereins Regensburg VI., 1–5.
- Strand, E. 1898a. Einige fundorte für Araneiden im südlischen Norwegen. Verh. zool.-bot. Ges. Wien 48, 401–404.
- Strand, E. 1898b. Oversigt over de Skandinaviske arter av slekten *Lycosa* (Lat.). Ent. Tidskr. 19, 145–158.
- Strand, E. 1899. Araneae Halingdaliae. Arch. Math. Naturv. 21, 1–23.
- Strand, E. 1900a. Zur Kenntnis der Arachniden Norwegens. K. norske Vidensk. Selsk. Skr. 1900, 1–46.
- Strand, E. 1900b. Fortegnelser over en del av Sparre-Schneider i det arktiske Norge insamlede Araneider. Tromsø Mus. Årsberetn. 23, 9–12.
- Strand, E. 1900c. Arachnologisches. Nyt. Mag. Naturvid. 38, 95–102.
- Strand, E. 1900d. Drei neue Xysticusarten. Zool. Anz. B. XXIII. 1900.
- Strand, E. 1901a. Bemerkungen über Norwegische Laterigraden nebst Beschreibungen drei neuer oder wenig bekannter arten. Abh. naturforsch. Ges. Görlitz 23, 1–15.
- Strand, E. 1901b. Theridiiden aus dem nördlichen Norwegen. Arch. Math. Naturv. 24, 1–66.
- Strand, E. 1902. Theridiiden aus dem westlichen Norwegen. Bergens Mus. Årb. 6, 1–23.
- Strand, E. 1904a. Die Dictyniden, Dysderiden, Drassiden, Clubioniden, und Ageleniden der Collett'schen Spinnensammlung. Forh. VidenskSelsk Krist. 1904, 1–16.
- Strand, E. 1904b. Theridiidae, Argiopidae und Mimetidae aus der Collet'schen Spinnensammlung. K. norske Vidensk. Selsk. Skr. 1903 107, 1–9.
- Strand, E. 1905. Araneae in, E. Strand, Coleoptera, Hymenoptera, Lepidoptera und Araneae. Report of the Second Norwegian Arctic Expedition in the «Fram» 1898–1902. No. 3, 1–30.
- Strand, E. 1906. Die arktischen Araneae, Opiliones und Chernetes. Fauna Arctica 4, 431–478.
- Strand, E. 1912. Neue Beiträge zur Arthropodenfauna Norwegens nebst gelegentlichen Bemerkungen über deutsche Arten. XIII–XIV. Nyt Mag. Naturv. 50, 200–222.
- Strand, E. 1915. Arachnologica varia I–IX. Archiv für Naturgeschichte A 11, 112–123.
- Strand, E. 1927. Catalogue des arachnides que j'ai décrits jusqu'à l'année 1926. Archiv für Naturgesichte B. 91.
- Strand, E. 1928. Spinnen aus Hallingdal in Südlischen Norwegen. Arch. Naturgesch. 92A, 86–9.
- Strand, E. 1941a. Über H. Tambs-Lyche's sogenannte «Revision» von V. Storm's Spinnensammlung aus der Umgebung von Trondheim, nebst einigen allgemeinen Bemerkungen über Literatur, «Spezialisten» und Sammlungen. Folia Zoologica et Hydrobiologica. XI. No. 1, 82–95.
- Strand, E. 1941b. Kritik der Arbeit des Herrn Hans Tambs-Lyche über die norwegischen *Pardosa*-[recte, *Lycosa*-] Arten. Folia Zoologica et Hydrobiologica. XI. No. 1, 95–109.
- Stroem, H. 1765. Beskrivelser over norske insekter. 1. Trondhjemske Selsk. Skrift., III.
- Stroem, H. 1768. Beskrivelse over norske insekter. 2. Trondhjemsk. Selsk. Skr. 4, 313–371 (Araneae, pp. 362–363).
- Tambs-Lyche, H. 1940. Die Norwegischen Spinnen der Gattung *Pardosa* Koch. Avh. norske VidensAkad. Oslo 1939, 1–59.
- Tambs-Lyche, H. 1941. Revision von Storm's Spinnensammlung aus der Umgebung von Trondheim. Kgl. norske Vid. Seisk. Forh. 13, 191–194.
- Tambs-Lyche, H. 1942. Notes on Norwegian spiders. Norsk. ent. Tidskr. 6, 107–114.
- Tambs-Lyche, H. 1955. Araneae from the Norwegian Finmarken =pedition 1924. Norsk ent. Tidsskr. 9,

- 224–226.
- Tambs-Lyche, H. 1964. A semi-marine spider *Halorates reprobus* (O.P.-Cambridge) in Norway. *Sarsia* 17, 17–19.
- Tambs-Lyche, H. 1967. Notes on the distribution of some arctic spiders. *Astarte* 28, 1–13.
- Thorell, T. 1870–73. Remarks on synonyms of European spiders. 654 pp.
- Thorell, T. 1875. Descriptions on several European and Nord-African spiders.
- Tveit, L. & Hauge, E. 1982. Notes on *Micrargus herbigradus* (Blackwall) and *M. apertus* (O.P.-Cambridge) in Norway (Araneae). *Fauna norv. Ser. B* 30, 34–38.
- Tveit, L. & Hauge, E. 1983. The spider fauna of Kristiansand and Setesdalen, S. Norway. *Fauna norv. Ser. B* 31, 23–45.
- Tullgren, A. 1942. Bidrag till kännedomen om den svenska spindelfaunaen. I. *Ent. Tidskr.* 63, 217–234.
- Tømmerås, B. Å. & Breistein, J. 1995. Fragmentation experiments in boreal forest. Purpose, methods and data from the field season 1994. NINA Oppdragsmelding 342 (in Norwegian), 1–43.
- Tømmerås, B. Å., Wilmann, B., Ødegaard, F., Gjerhaug, J. O., Breistein, J., Abildsnes, J., Prestø, T., Aakra, K. & Krogstad, S. 2000. Effekter av fragmentering på biodiversitet i granskog. NINA Fagrappart 40, 1–89.
- Waaler, P. F. 1966a. *Theridion bellicosum* Simon (Araneae, Theridiidae) new to Norway. *Norsk ent. Tidsskr.* 13, 183–184.
- Waaler, P. F. 1966b. *Porrhomma campbelli* F. O. P. Cambridge (Araneae, Linyphiidae) in Norway. *Norsk ent. Tidsskr.* 13, 407.
- Waaler, P. F. 1967. A collection of spiders from Son, Norway. *Norsk ent. Tidsskr.* 14, 91–93.
- Waaler, P. F. 1970a. Spiders (Araneae) from Syd-Varanger, North Norway with a note on gyandromorph *Cornicularia*. *Rhizocrinus* 1, 1–9.
- Waaler, P. F. 1970b. *Hyptiotes paradoxus* (C. L. Koch) in Norway. *Norsk ent. Tidsskr.* 17, 103–104.
- Waaler, P. F. 1971a. Spiders new to Norway. *Norsk ent. Tidsskr.* 18, 17–24.
- Waaler, P. F. 1971b. Spiders (Araneae) in mosses from Son, Norway. *Norsk ent. Tidsskr.* 18, 95–98.
- Waaler, P. F. 1971. *Edderkoppene*, pp. 126–142 in Frislid, R. & Semb-Johanson, A. (eds.). *Norges Dyr* 4. J. W. Cappelen, Oslo. 429 pp.
- Waaler, P. F. 1972. Spiders (Araneae) from Ringsaker, Norway. *Norsk ent. Tidsskr.* 19, 49–57.
- Westring, N. Förteckning öfver de till närvarande tid Kände, i Sverige förekommande Spindlarter, utgörande ett antal af 253, deraf 132 äro nya för svenska Faunan. Göteborg. Kongl. Vet. Handl. 2, 25–62.
- Westring, N. 1861. Aranæae Suecicae descriptæ. Göteborg.
- Wunderlich, J. 1973. Zwei für Norwegen neue Spinnentiere, *Nemastoma bimaculatum* und *Theridion montanum* (Arachnida, Opiliones, Nemastomatidae und Aranæae, Theridiidae). *Senckenb. biol.* 54, 177.
- Østbye, E., Lauritzen, S.-E., Fjellberg, A., Hauge, E., Leinaas, H. P., Ottesen, P. & Solhøy, T. 1987. Invertebrates of Norwegian caves I, Gastropoda, Oligochaeta, Aranæae, Acari, Amphipoda, Collembola, Coleoptera, Lepidoptera and Diptera
- ## BIBLIOGRAPHY-SVALBARD ARCHIPELAGO AND JAN MAYEN
- Bengtson, S.-A., Fjellberg, A. & Solhøy, T. 1974. Abundance of tundra arthropods in Spitsbergen. *Ent. Scand.* 5, 137–142; Copenhagen.
- Bengtson, S.-A. Fjellberg, A. 1975. Summer food of the purple sandpiper (*Calidris maritima*) in Spitsbergen. *Astarte* 8, 1–6.
- Bristowe, W. S. 1922. The Insects and Arachnids of Jan Mayen.-*Proc. Cambridge Phil. Soc.* 21, 38–43.
- Bristowe, W. S. 1925. The fauna of the arctic island of Jan Mayen and its probable origin. *Ann. Mag. Nat. Hist.* 9 (15), 480–485; London.
- Bristowe, W. S. 1933. The spiders of the Bear Island.- *Norsk ent. Tidskrift* 3 (3) (1935), 149–154.
- Bristowe, W. S. 1948. Spiders from the Arctic Island of Jan Mayen. *Proc. Zool. Soc. London* 118, 223–225.
- Coulson, S. J. A review of the terrestrial and freshwater invertebrate fauna of the High Arctic archipelago of Svalbard. *Norw. J. Entomol.* 47, 41–63.
- Hinz, W. 1976. Zur Ökologie der Tundra Zentral-spitbergen.-*Norsk Polarinst.* 163, 1–47; Oslo.
- Holm, Å. 1937. Zur Kenntnis der Spinnenfauna Spitzbergens und der Bären Insel. *Ark. Zool.* 29 A (18), 1–13; Stockholm.
- Holm, Å. 1958. The spiders of the Isfjord Region of Spitsbergen.-*Zool. Bidr. Uppsala* 33, 29–67, Pl. I.
- Huhta, V. 1967. Some notes on the spider fauna of Spitsbergen. *Ann. Ent. Fenn.* 33 (1), 27; Helsinki.
- Jackson, A. R. 1922. On the spiders captured by Mr. C.S. Elton at Spitsbergen and Bear Island in 1921. Results of the Oxford University expedition to Spitsbergen. 5. *Ann. Mag. nat. Hist.* (9) 9, 163–169; London.
- Jackson, A. R. 1924. On the spiders of Spitsbergen. Results of the Merton College expedition to Spitsbergen in 1923. *Ann. Mag. nat. Hist.* 9 (13), 77–79; London.

- Jackson, A. Randell 1925. On the third collection of spiders captured by Mr. C.S. Elton in Spitsbergen. Results of the Oxford University Arctic Expedition, 1924. Ann. Mag. nat. Hist. 9 (15), 536-537; London.
- Jackson, A. R. 1934. On a collection of spiders made in 1928 by Dr. Sig Thor in Spitsbergen. Norsk ent. Tidsskrift 3 (5), 332-354; Oslo.
- Koponen, S. 1980. Spider fauna in the Adventfjorden area, Spitsbergen. Rep. Kevo Subarctic Res. Stat. 16, 13-16; Turku.
- Kulczynski, V. 1902. Zoologische Ergebnisse der russischen Expedition nach Spitzbergen. Araneae et Oribatidae. Ann. Mus. Zool. Acad. Imp. Sci. St. Petersburg 7 (3), 335-354, pl. VII.
- Roewer, C. F. 1930. Arachnoidea. In Thor, S. (1930), Beiträge zur Kenntnis der Invertebratenfauna von Svalbard. Skrifter om Svalbard og Ishavet 27, 55-56; Oslo.
- Strand, Embrik 1915. Arachnologica varia. I-IX. ... VIII. Spinnen von Spitzbergen. Arch. Naturgeschichte 81 A (11), 112-123; Berlin.
- Strand, E. 1941. Kritik der Sig Thor'schen «Beiträge» zur Kenntnis der Invertebraten-Fauna von Svalbard. Folia zool. hydrobiol. 11 (1), 1-11; Riga.
- Summerhaves, V.S., C.S. Elton (1923 & 1928). Contributions to the ecology of Spitsbergen and Bear Island. Further contributions to the ecology of Spitsbergen.-J. Ecolgy 11, 214-286; 16, 193-268.
- Thaler, K. 1975. Über einige Spinnen aus Spitzbergen (Arachnida, Aranei). Mitt. schweiz. ent. Ges. 48 (3/4), 437-442; Zürich.
- Tambs-Lyche, H. 1967. Notes on the distribution of some Arctic spiders. Astarte 28, 1 – 13.
- Toft, S. 1979. Spiders from Jan Mayen. Fauna norvegica Ser B 26, 24-25; Oslo.
- record of *Uloborus plumipes* Lucas (Uloboridae) from the Faroes. Norw. J. Entomol. 48, 104.
- Bristowe, W. S. 1939. The comity of spiders. Ray Society, London, 228 pp.
- Denis, J. 1949. Notes sur les érigonides. XVI. Essai sur la détermination des femelles d'érigonides. Bull. Soc. Hist. nat. Toulouse 83, 129-158.
- Dondale, C. D. & J. H. Redner. 1983. Revision of the wolf spiders of the genus *Arctosa* C. L. Koch in North and Central America (Araneae, Lycosidae). J. Arachnol. 11, 1-30.
- Eskov, K. Y. 1994. Catalogue of the linyphiid spiders of northern Asia (Arachnida, Araneae, Linyphiidae). Pensoft Publishers, Sofia, 144 pp.
- Eskov, K. Y. & Marusik, Y. M. 1994. New data on the taxonomy and faunistics of North Asian linyphiid spiders (Aranei Linyphiidae). Arthropoda Selecta 2(4), 41-79.
- Grothendieck, K. & Kraus, O. 1994. Die Wasserspinne *Argyroneta aquatica*, Verwandtschaft und Spezialisierung (Arachnida, Araneae, Agelenidae). Verh. naturwiss. Ver. Hamburg (N. F.) 34, 259 – 273.
- Hackman, W. 1954. The spiders of Newfoundland. Acta zool. fenn. 79, 1-99.
- Hauge, E. 1989. An annotated check-list of Norwegian spiders (Araneae). Insecta Norvegiae 4, 1-40.
- Heimer, S. & Nentwig, W. 1991. Spinnen Mitteleuropas, Ein Bestimmungsbuch. 543 pp. Verlag Paul Parey, Berlin.
- Hippa, H. & Mannila, R. 1982. *Pardosa maisa* n. sp. (Araneae, Lycosidae) from northern Europe. Bull. Br. arachnol. Soc. 5, 420-422.
- Holm, Å. 1944. Revision einiger norwegischen Spinnenarten und Bemerkungen über deren Vorkommen in Schweden. Ent. Tidskr. 65, 122 – 134.
- Holm, Å. 1945. Zur Kenntnis der Spinnenfauna des Torneträskgebietes. Ark. Zool. 36(A 15), 1-80.
- Holm, Å. 1950. Studien über die Spinnenfauna des Torneträskgebietes. Zool. Bidr. Upps. 29, 103-213.
- Lehtinen, P. T. 1967. Classification of the cribellate spiders and some allied families, with notes on the evolution of the suborder Araneomorpha. Ann. Zool. Fenn. 4, 199-468.
- Lehtinen, P.T., Koponen, S. & Saaristo, M. 1979. Studies on the spider fauna of the southwestern archipelago of Finland II. Memoranda Soc. Fauna Flora Fennica 55, 33-52.
- Levi, H. W. 1991. The Neotropical and Mexican species of the orb-weaver genera *Araneus*, *Dubiepeira*, and *Aculepeira* (Araneae, Araneidae). Bull. Mus. comp. Zool. Harv. 152, 167-315.
- Logunov, D. V. 1992. Definition of the spider genus *Talavera* (Araneae, Salticidae), with a description of a new species. Bull. Inst. R. Sci. nat. Belg. (Ent.)

REFERENCES

- Aakra, K. 2000. New records of spiders (Araneae) from Norway with notes on epigynal characters of *Philodromus fuscomarginatus* (De Geer) (Philodromidae) and *Araneus sturmi* (Hahn) (Araneidae). Norw. J. Entomol. 47, 77-88.
- Aakra, K. 2002. Taxonomic notes on some Norwegian linyphiid spiders described by E. Strand (Araneae, Linyphiidae). Bull. Br. Arachnol. Soc 12, 267-269.
- Aakra, K. & Hauge, E. 2002. Araneae Norvegiae. Checklist and distribution maps of Norwegian spiders with Taxonomic, zoogeographic and ecological notes. <http://www.museumsnett.no/midttromsmuseum/forskningsssider/norspider/index.htm>
- Aakra, K & Olsen, K. M. 2003. Two introduced species of spiders (Araneae) new to Norway and the first

- 62, 75-82.
- Logunov, D. V. & Kronestedt, T. 1997 A new Palearctic species of the genus *Sitticus* Simon, with notes on related species in the *floricola* group (Araneae, Salticidae). Bull. Br. arachnol. Soc. 10, 225-233.
- Jäger, P. 1999. Sparassidae-the valid scientific name for the huntsman spiders (Arachnida, Araneae). Arachnol. Mitt. 17, 1-10.
- Kronestedt, T. 1975. A new species of the genus *Lepthyphantes* from Sweden (Araneae, Linyphiidae). Ent. Tidskr. 96, 162-166.
- Kronestedt, T. 1990. Separation of two species standing as *Alopecosa aculeata* (Clerck) by morphological, behavioural and ecological characters, with remarks on related species in the *pulverulenta* group (Araneae, Lycosidae). Zoologica Scr. 19, 203-225.
- Kronestedt, T. 1993. On *Rugathodes instabilis* (O. P. Cambridge) and *R. bellicosus* (Simon) (Araneae, Theridiidae). Ent. Tidskr. 114, 111 - 113.
- Marusik, Y. M., Eskov, K., Logunov, D. V. & Basarukin, A. M. 1993. A check-list of spiders (Arachnida Aranei) from Shakalin and Kurile islands. Arthropoda Selecta 2, 73 - 85.
- Marusik, Y. M. & Tanasevitch, A. V. 1998. Notes on the spider genus *Stylocetor* Simon, 1884 and some related genera, with description of two new species from Siberia (Aranei, Linyphiidae). Arthropoda Selecta 7, 153-159.
- Millidge, A. F. 1984. The erigonine spiders of North America. Part 7. Miscellaneous genera. J. Arachnol. 12, 121-169.
- Ovtsharenko, V. I. & Marusik Y. M. 1988. Spiders of the family Gnaphosidae (Aranei) of the north-east of the USSR (the Magadan Province). Ent. Obozr. 67, 204-217.
- Palmgren, P. 1975. Die Spinnenfauna Finnlands und Ostfennoscandiens VI, Linyphiidae 1. Fauna fenn. 28, 1-102.
- Palmgren, P. 1976. Die Spinnenfauna Finnlands und Ostfennoscandiens. VII. Linyphiidae 2. Fauna Fennica 29, 1- 126.
- Piel, W & Nutt, K. 1997. *Kaira* is a likely sister group to *Metepeira*, and *Zygilla* is an araneid (Araneae, Araneidae), evidence from mitochondrial DNA. J. Arachnol. 25, 262 - 268.
- Platnick, N. I. 2003. The world spider catalog, version 3.5. American Museum of Natural History, online at <http://research.amnh.org/entomology/spiders/catalog81-87/index.html>
- Ramírez, M. J., A. B. Bonaldo, & A. D. Brescovit. Revisión del género *Macerio* y comentarios sobre la ubicación de *Cheiracanthium*, *Tecution* y *Helebiona* (Araneae, Miturgidae, Eutichurinae). Iheringia (Zool.) 82, 43-66.
- Roberts, M. J. 1987. The spiders of Great Britain and Ireland, Volume 2, Linyphiidae and check list. Harley Books, Colchester, England.
- Roberts, M. J. 1998. Spinnen Gids. Uitgebreide beschrijving van ruim 500 Europese soorten. 390 pp. Baarn (NL) (Tirion Natuur).
- Saaristo, M. I. 1973. Taxonomical analysis of the type-species of *Agyneta*, *Anomalaria*, *Meioneta*, *Aprolagus*, and *Syedrula* (Araneae, Linyphiidae). Ann. Zool. Fenn. 10, 451-466.
- Saaristo, M. I. & Eskov, K. Y. 1996. Taxonomy and zoogeography of the hypoarctic erigonine spider genus *Semljicola* (Araneae, Linyphiidae). Acta Zool. Fennica 201, 47 - 69.
- Saaristo, M. I. & S. Koponen 1998. A review of northern Canadian spiders of the genus *Agyneta* (Araneae, Linyphiidae), with descriptions of two new species. Can. J. Zool. 76, 566-583.
- Saaristo, M. & A.V. Tanasevitch 1996. Redelimitation of subfamily Micronetinae Hull, 1920 and the genus *Lepthyphantes* Menge, 1866 with descriptions of some new genera (Aranei, Linyphiidae). Ber. nat.-med. Ver. Innsbruck 83, 163-186
- Saaristo, M.I., Tanasevitch A.V. 1999. Reclassification of the *mughi*-group of the genus *Lepthyphantes* Menge, 1866 (sensu lato) (Aranei, Linyphiidae, Micronetinae). Ber. nat.-med. Verein Innsbruck, 87, 390-398.
- Saaristo, M. I. & Tanasevitch, A.V. 2000. Systematics of the *Bolyphantes-Poeciloneta* genus-group of the subfamily Micronetinae Hull, 1920 (Arachnida, Araneae, Linyphiidae). Reichenbachia, 33 (32), 253-263.
- Scharff, N. & Coddington, J. A. 1997. A phylogenetic analysis of the orb-weaving spider family Araneidae (Arachnida, Araneae). Zool. J. Linn. Soc. 120, 355 - 434.
- Strand, E. 1900. Zur Kenntnis der Arachniden Norwegens. Kong. norsk. Vid. Selsk. Skrift. 1900 (2), 1-46.
- Strand, E. 1904. Die dictyniden, dysderiden, drassiden, clubioniden und ageleniden der Collett'schen Spinnensammlung. Forh. Vid. Selsk. Christian 1904 (5), 1-16.
- Tambs-Lyche, H. 1940. Die Norwegischen Spinnen der Gattung *Pardosa* Koch. Avh. norske VidensAkad. Oslo 1939, 1 - 59.
- Tambs-Lyche, H. 1942. Notes on Norwegian spiders. Norsk ent. Tidsskr. 6, 107-114.
- Tanasevitch, A. V. The spider family Linyphiidae in the fauna of the Caucasus (Arachnida, Aranei). Pp. 5-114 in B. R. Striganova (ed.), Fauna nazemnykh bespozvonochnykh Kavkaza. Moscow, Akaedemia

Nauk.

- Thaler, K. 1970. Über einige wenig bekannte Zwerbspinnen aus den Alpen (Arachnida, Araneae, Erigonidae). Ber. nat. -med. Ver. Innsbruck 58, 255–276.
- Thaler, K., P. van Helsdingen & C. Deltshev. 1994. Vikariante Verbreitung im Artenkomplex von *Lepthyphantes annulatus* in Europa und ihre Deutung (Araneae, Linyphiidae). Zool. Anz. 232, 111–127.
- Töpfer-Hofman, G., Cordes, D. & Helversen, Otto von, 2000. Cryptic species and behavioural isolation in the *Pardosa lugubris* group (Araneae, Lycosidae), with description of two new species. Bull. Br. arachnol. Soc. 11, 257 – 274.
- Tullgren, A. 1946. Svenska spindelfauna, 3. Egentliga spindlar. Araneae. Fam. 5-7. Clubionidae, Zoridae och Gnaphosidae. Entomologiska Föreningen, Stockholm, 141 pp.
- Tullgren, A. 1952. Zur Kenntnis schwedischer Spinnen. I. Ent. Tidskr. 73, 151–177.
- Wunderlich, J. 1973. Zur Spinnefauna Deutschlands, XV. Witere seltene und bisher unbekannte Arten sowie Anmerkungen zur Taxonomie und Synonymie (Arachnida, Araneae). Senckenberg. Biol. 54, 405 – 428.
- Wunderlich, J. 1987. Die Spinnen der Kanarischen Inseln und Madeiras, Adaptive Radiation, Biogeographie, Revisionen und Neubeschreibungen. Triops Verlag, Langen, West Germany.
- Wunderlich, J. 1992. Die Spinnen-Fauna Makaronesischen Inseln, Tazonomie, Ökologie, Biogeographie und Evolution. Beitr. Araneol. 1, 1–619.
- Wunderlich, J. 1995. Zur Taxonomie europäischer Gattungen der Zwerbspinnen (Arachnida, Araneae, Linyphiidae). Beitr. Araneol. 4, 643 – 654.
- Wunderlich, J. & Schuett, K. 1995. Beschreibung der bisher verkannten Sackspinnen-Art *Clubiona frisia* n. sp. aus Europa (Arachnida, Araneae, Clubionidae). Ent. Z. Frankf. a. M. 105, 10 – 17.
- Zabka, M. 1997. Salticidae, Pajaki skaczace (Arachnida, Araneae). Fauna Polski 19, 1–188.

Received 22 September 2003,
accepted 17 November 2003