

## Spiders of Poland



Details

Monday, 17 March 2014

### **CHECK-LIST OF SPIDERS (*ARANEAE*) OF POLAND**

1. DECEMBER 2008

Prepared by:

**Robert Rozwalka:** Department of Zoology, Maria Curie-Skłodowska University, Akademicka 19, 20-033 Lublin, [arachnologia@wp.pl](mailto:arachnologia@wp.pl)

**Marzena Stańska:** Department of Zoology, Siedlce University of Natural Sciences and Humanities, Prusa 12, 08-110 Siedlce, [stanska@uph.edu.pl](mailto:stanska@uph.edu.pl)

#### **ATYPIDAE**

- 1 *Atypus affinis* EICHWALD, 1830
- 2 *Atypus muralis* BERTKAU, 1890
- 3 *Atypus piceus* (SULZER, 1776)

#### **SCYTODIDAE**

- 4 *Scytodes thoracica* (LATREILLE, 1802)

#### **PHOLCIDAE**

- 5 *Pholcus opilionoides* (SCHRANK, 1781)
- 6 *Pholcus phalangioides* (FUSSLIN, 1775)
- 7 *Psilochorus simoni* (BERLAND, 1911)

#### **SEGESTRIIDAE**

- 8 *Segestria bavarica* C.L. KOCH, 1843
- 9 *Segestria senoculata* (LINNAEUS, 1758)

#### **DYSDERIDAE**

- 10 *Dasumia carpatica* (KULCZYŃSKI, 1882)
- 11 *Dysdera erythrina* (WALCKENAER, 1802)
- 12 *Harpactea hombergi* (SCOPOLI, 1763)
- 13 *Harpactea lepida* (C.L. KOCH, 1839)

- 14 *Harpactea rubicunda* (C.L. KOCH, 1839)

#### **OONOPIDAE**

- 15 *Oonops domesticus* DALMAS, 1916  
16 *Oonops pulcher* TEMPLETON, 1835

#### **MIMETIDAE**

- 17 *Ero aphana* (WALCKENAER, 1802)  
18 *Ero cambridgei* KULCZYŃSKI, 1911  
19 *Ero furcata* (VILLERS, 1789)  
20 *Ero tuberculata* (DE GEER, 1778)

#### **ERESIDAE**

- 21 *Eresus kollari* (ROSSI, 1846)

#### **ULOBORIDAE**

- 22 *Hyptiotes paradoxus* (C.L. KOCH, 1834)  
23 *Uloborus plumipes* LUCAS, 1846

#### **NESTICIDAE**

- 24 *Nesticus cellulanus* (CLERCK, 1758)

#### **THERIDIIDAE**

- 25 *Achaearana lunata* (CLERCK, 1758)  
26 *Achaearana ohlerti* (THORELL, 1870)  
27 *Achaearana riparia* (BLACKWALL, 1834)  
28 *Achaearana simulans* (THORELL, 1875)  
29 *Achaearana tabulata* LEVI, 1980  
30 *Achaearana tepidariorum* (C.L. KOCH, 1841)  
31 *Anelosimus pulchellus* (WALCKENAER, 1802)  
32 *Anelosimus vittatus* (C.L. KOCH, 1836)  
33 *Crustulina guttata* (WIDER, 1834)  
34 *Crustulina sticta* (O. P.-CAMBRIDGE, 1861)  
35 *Dipoena braccata* (C.L. KOCH, 1841)  
36 *Dipoena coracina* (C.L. KOCH, 1841)  
37 *Dipoena erythropus* (SIMON, 1881)  
38 *Dipoena inornata* (O. P.-CAMBRIDGE, 1861)  
39 *Dipoena melanogaster* (C.L. KOCH, 1837)  
40 *Dipoena nigroreticulata* (SIMON, 1879)  
41 *Dipoena prona* (MENGE, 1868)  
42 *Dipoena torva* (THORELL, 1875)  
43 *Enoplognatha caricis* (FICKERT, 1876)  
44 *Enoplognatha latimana* HIPPA & OKSALA, 1982  
45 *Enoplognatha mordax* (THORELL, 1875)  
46 *Enoplognatha oelandica* (THORELL, 1875)  
47 *Enoplognatha ovata* (CLERCK, 1758)  
48 *Enoplognatha thoracica* (HAHN, 1833)  
49 *Episinus angulatus* (BLACKWALL, 1836)  
50 *Episinus truncatus* LATREILLE, 1809  
51 *Euryopsis flavomaculata* (C.L. KOCH, 1836)  
52 *Euryopsis laeta* (WESTRING, 1861)  
53 *Euryopsis saukea* LEVI, 1951  
54 *Keijia tinctoria* (WALCKENAER, 1802)  
55 *Lasaeola tristis* (HAHN, 1833)

- 56 *Neottiura bimaculata* (LINNAEUS, 1758)
- 57 *Paidiscura pallens* (BLACKWALL, 1834)
- 58 *Pholcomma gibbum* (WESTRING, 1851)
- 59 *Robertus arundineti* (O. P.-CAMBRIDGE, 1871)
- 60 *Robertus insignis* O. P.-CAMBRIDGE, 1907
- 61 *Robertus lividus* (BLACKWALL, 1836)
- 62 *Robertus neglectus* (O. P.-CAMBRIDGE, 1871)
- 63 *Robertus scoticus* JACKSON, 1914
- 64 *Robertus truncorum* (L. KOCH, 1872)
- 65 *Robertus unguulatus* VOGELSANGER, 1944
- 66 *Rugathodes bellicosus* (SIMON, 1873)
- 67 *Rugathodes instabilis* (O. P.-CAMBRIDGE, 1871)
- 68 *Simitidion simile* (C.L. KOCH, 1836)
- 69 *Steatoda albomaculata* (DE GEER, 1778)
- 70 *Steatoda bipunctata* (LINNAEUS, 1758)
- 71 *Steatoda castanea* (CLERCK, 1758)
- 72 *Steatoda grossa* (C.L. KOCH, 1838)
- 73 *Steatoda phalerata* (PANZER, 1801)
- 74 *Theonoe minutissima* (O. P.-CAMBRIDGE, 1879)
- 75 *Theridion betteni* WIEHLE, 1960
- 76 *Theridion blackwalli* O. P.-CAMBRIDGE, 1871
- 77 *Theridion conigerum* (SIMON, 1914)
- 78 *Theridion familiare* O. P.-CAMBRIDGE, 1871
- 79 *Theridion impressum* L. KOCH, 1881
- 80 *Theridion melanurum* HAHN, 1831
- 81 *Theridion mystaceum* L. KOCH, 1870
- 82 *Theridion nigrovariegatum* SIMON, 1873
- 83 *Theridion palmgreni* MARUSIK & CELLARIUS, 1986
- 84 *Theridion pictum* (WALCKENAER, 1802)
- 85 *Theridion pinastri* L. KOCH, 1872
- 86 *Theridion sisyphium* (CLERCK, 1758)
- 87 *Theridion uhligi* MARTIN, 1974
- 88 *Theridion varians* HAHN, 1833

#### **THERIDIOSOMATIDAE**

- 89 *Theridiosoma gemmosum* (L. KOCH, 1877)

#### **MYSMENIDAE**

- 90 *Mysmenella jobi* (KRAUS, 1967)
- 91 *Trogloneta granulum* SIMON, 1922

#### **LINYPHIIDAE**

- 92 *Abacoproeces saltuum* (L. KOCH, 1872)
- 93 *Abiskoa abiskoensis* (HOLM, 1945)
- 94 *Acartauchenius scurrilis* (O. P.-CAMBRIDGE, 1872)
- 95 *Agyphantes expunctus* (O. P.-CAMBRIDGE, 1875)
- 96 *Agyphantes arietans* (O. P.-CAMBRIDGE, 1872)
- 97 *Agyphantes cauta* (O. P.-CAMBRIDGE, 1902)
- 98 *Agyphantes conigera* (O. P.-CAMBRIDGE, 1863)
- 99 *Agyphantes decora* (O. P.-CAMBRIDGE, 1871)
- 100 *Agyphantes ramosa* JACKSON, 1914
- 101 *Agyphantes subtilis* (O. P.-CAMBRIDGE, 1863)
- 102 *Allomengea scopigera* (GRUBE, 1859)
- 103 *Allomengea vidua* (L. KOCH, 1879)

- 104 *Anguliphantes angulipalpis* (WESTRING, 1851)
- 105 *Anguliphantes monticola* (KULCZYŃSKI, 1882)
- 106 *Anguliphantes tripartitus* (MILLER & SVATON 1978)
- 107 *Aphileta misera* (O. P.-CAMBRIDGE, 1882)
- 108 *Araeoncus crassiceps* (WESTRING, 1861)
- 109 *Araeoncus humilis* (BLACKWALL, 1841)
- 110 *Asthenargus helveticus* SCHENKEL, 1936
- 111 *Asthenargus paganus* (SIMON, 1884)
- 112 *Baryphyma gowerense* (LOCKET, 1968)
- 113 *Baryphyma pratense* (BLACKWALL, 1861)
- 114 *Baryphyma trifrons* (O. P.-CAMBRIDGE, 1863)
- 115 *Bathyphantes approximatus* (O. P.-CAMBRIDGE, 1871)
- 116 *Bathyphantes eumenis* (L. KOCH, 1879)
- 117 *Bathyphantes gracilis* (BLACKWALL, 1841)
- 118 *Bathyphantes nigrinus* (WESTRING, 1851)
- 119 *Bathyphantes parvulus* (WESTRING, 1851)
- 120 *Bathyphantes setiger* F. P.-CAMBRIDGE, 1894
- 121 *Bathyphantes similis* KULCZYŃSKI, 1894
- 122 *Bolephthyphantes index* (THORELL, 1856)
- 123 *Bolyphantes alticeps* (SUNDEVALL, 1833)
- 124 *Bolyphantes luteolus* (BLACKWALL, 1833)
- 125 *Carorita limnaea* (CROSBY & BISHOP, 1927)
- 126 *Caviphantes saxetorum* (HULL, 1916)
- 127 *Centromerita bicolor* (BLACKWALL, 1833)
- 128 *Centromerita concinna* (THORELL, 1875)
- 129 *Centromerus arcanus* (O. P.-CAMBRIDGE, 1873)
- 130 *Centromerus brevivulvatus* DAHL, 1912
- 131 *Centromerus cavernarum* (L. KOCH, 1872)
- 132 *Centromerus incilium* (L. KOCH, 1881)
- 133 *Centromerus levitarsis* (SIMON, 1884)
- 134 *Centromerus pabulator* (O. P.-CAMBRIDGE, 1875)
- 135 *Centromerus persimilis* (O. P.-CAMBRIDGE, 1912)
- 136 *Centromerus prudens* (O. P.-CAMBRIDGE, 1873)
- 137 *Centromerus sellarius* (SIMON, 1884)
- 138 *Centromerus semiater* (L. KOCH, 1879)
- 139 *Centromerus serratus* (O. P.-CAMBRIDGE, 1875)
- 140 *Centromerus silvicola* (KULCZYŃSKI, 1894)
- 141 *Centromerus sylvaticus* (BLACKWALL, 1841)
- 142 *Ceraticelus bulbosus* (EMERTON, 1882)
- 143 *Ceratinella brevipes* (WESTRING, 1851)
- 144 *Ceratinella brevis* (WIDER, 1834)
- 145 *Ceratinella maior* KULCZYŃSKI, 1894
- 146 *Ceratinella scabrosa* (O. P.-CAMBRIDGE, 1871)
- 147 *Ceratinella wideri* (THORELL, 1871)
- 148 *Cinetata gradata* (SIMON, 1881)
- 149 *Cnephalocotes obscurus* (BLACKWALL, 1834)
- 150 *Collinsia distincta* (SIMON, 1884)
- 151 *Collinsia inerrans* (O. P.-CAMBRIDGE, 1885)
- 152 *Dicymbium nigrum* (BLACKWALL, 1834)
- 152a *Dicymbium nigrum brevisetosum* LOCKET, 1962
- 153 *Dicymbium tibiale* (BLACKWALL, 1836)
- 154 *Diplocentria bidentata* (EMERTON, 1882)
- 155 *Diplocephalus cristatus* (BLACKWALL, 1833)

- 156 *Diplocephalus dentatus* TULLGREN, 1955
- 157 *Diplocephalus helleri* (L. KOCH, 1869)
- 158 *Diplocephalus latifrons* (O. P.-CAMBRIDGE, 1863)
- 159 *Diplocephalus permixtus* (O. P.-CAMBRIDGE, 1871)
- 160 *Diplocephalus picinus* (BLACKWALL, 1841)
- 161 *Diplocephalus protuberans* (O. P.-CAMBRIDGE, 1875)
- 162 *Diplostyla concolor* (WIDER, 1834)
- 163 *Dismodicus bifrons* (BLACKWALL, 1841)
- 164 *Dismodicus elevatus* (C.L. KOCH, 1838)
- 165 *Donacochara speciosa* (THORELL, 1875)
- 166 *Drapetisca socialis* (SUNDEVALL, 1832)
- 167 *Drepanotylus uncutus* (O. P.-CAMBRIDGE, 1873)
- 168 *Entelecara acuminata* (WIDER, 1834)
- 169 *Entelecara congenera* (O. P.-CAMBRIDGE, 1879)
- 170 *Entelecara errata* O. P.-CAMBRIDGE, 1913
- 171 *Entelecara erythropus* (WESTRING, 1851)
- 172 *Entelecara flavipes* (BLACKWALL, 1834)
- 173 *Entelecara media* KULCZYŃSKI, 1887
- 174 *Entelecara omissa* O. P.-CAMBRIDGE, 1902
  
- 175a *Erigone arctica maritima* KULCZYŃSKI, 1902
- 176 *Erigone atra* BLACKWALL, 1833
- 177 *Erigone dentigera* O. P.-CAMBRIDGE, 1874
- 178 *Erigone dentipalpis* (WIDER, 1834)
- 179 *Erigone longipalpis* (SUNDEVALL, 1829)
- 180 *Erigone tirolensis* L. KOCH, 1872
- 181 *Erigonella hiemalis* (BLACKWALL, 1833)
- 182 *Erigonella ignobilis* (O. P.-CAMBRIDGE, 1871)
- 183 *Erigonella subelevata* (L. KOCH, 1869) DD
- 184 *Erigonoplus globipes* (L. KOCH, 1872) VU
- 185 *Evansia merens* O. P.-CAMBRIDGE, 1900 EN
- 186 *Floronia bucculenta* (CLERCK, 1758)
- 187 *Formiphantes lephthyphantiformis* (STRAND, 1907)
- 188 *Frontinellina frutetorum* (C.L. KOCH, 1834)
- 189 *Glyphesis cottonae* (LA TOUCHE, 1944)
- 190 *Glyphesis servulus* (SIMON, 1881)
- 191 *Gnathonarium dentatum* (WIDER, 1834)
- 192 *Gonatium paradoxum* (L. KOCH, 1869)
- 193 *Gonatium rubellum* (BLACKWALL, 1841)
- 194 *Gonatium rubens* (BLACKWALL, 1833)
- 195 *Gongylidiellum latebricola* (O. P.-CAMBRIDGE, 1871)
- 196 *Gongylidiellum murcidum* SIMON, 1884
- 197 *Gongylidiellum vivum* (O. P.-CAMBRIDGE, 1875)
- 198 *Gongylidium rufipes* (LINNAEUS, 1758)
- 199 *Helophora insignis* (BLACKWALL, 1841)
- 200 *Hilaira excisa* (O. P.-CAMBRIDGE, 1871)
- 201 *Hilaira frigida* (THORELL, 1872)
- 202 *Hilaira nubigena* HULL, 1911
- 203 *Hilaira tatica* KULCZYŃSKI, 1915
- 204 *Hylyphantes graminicola* (SUNDEVALL, 1829)
- 205 *Hylyphantes nigrinus* (SIMON, 1881)
- 206 *Hypomma bituberculatum* (WIDER, 1834)
- 207 *Hypomma cornutum* (BLACKWALL, 1833)
- 208 *Hypomma fulvum* (BÖSENBERG, 1903)

- 209 *Hypselistes jacksoni* (O. P.-CAMBRIDGE, 1902)
- 210 *Hypsocephalus dahli* (LESSERT, 1909)
- 211 *Improphantes complicatus* (EMERTON, 1882)
- 212 *Improphantes decolor* (WESTRING, 1861)
- 213 *Improphantes nitidus* (THORELL, 1875)
- 214 *Incestophantes annulatus* (KULCZYŃSKI, 1882)
- 215 *Incestophantes crucifer* (MENGE, 1866)
- 216 *Jacksonella falconeri* (JACKSON, 1908)
- 217 *Kaestneria dorsalis* (WIDER, 1834)
- 218 *Kaestneria pullata* (O. P.-CAMBRIDGE, 1863)
- 219 *Kaestneria torrentum* (KULCZYŃSKI, 1882)
- 220 *Kratochviliella bicapitata* MILLER, 1938
- 221 *Labulla thoracica* (WIDER, 1834)
- 222 *Lasiargus hirsutus* (MENGE, 1869)
- 223 *Lepthyphantes keyserlingi* (AUSSERER, 1867)
- 224 *Lepthyphantes leprosus* (OHLERT, 1865)
- 225 *Lepthyphantes minutus* (BLACKWALL, 1833)
- 226 *Lepthyphantes nodifer* (SIMON 1884)
- 227 *Leptorhoptrum robustum* (WESTRING, 1851)
- 228 *Leptothrix hardyi* (BLACKWALL, 1850)
- 229 *Lessertia dentichelis* (SIMON, 1884)
- 230 *Linyphia hortensis* SUNDEVALL, 1829
- 231 *Linyphia triangularis* (CLERCK, 1758)
- 232 *Lophomma punctatum* (BLACKWALL, 1841)
- 233 *Macrargus carpenteri* (O. P.-CAMBRIDGE, 1894)
- 234 *Macrargus rufus* (WIDER, 1834)
- 235 *Mansuphantes arciger* (KULCZYŃSKI, 1882)
- 236 *Mansuphantes mansuetus* (THORELL, 1875)
- 237 *Maro minutus* O. P.-CAMBRIDGE, 1906
- 238 *Maro sublestus* FALCONER, 1915
- 239 *Maso gallicus* SIMON, 1884
- 240 *Maso sundevalli* (WESTRING, 1851)
- 241 *Mecopisthes silus* (O. P.-CAMBRIDGE, 1872)
- 242 *Mecynargus foveatus* (DAHL, 1912)
- 243 *Mecynargus longus* (KULCZYŃSKI, 1882)
- 244 *Mecynargus morulus* (O. P.-CAMBRIDGE, 1873)
- 245 *Megalepthyphantes nebulosus* (SUNDEVALL, 1829)
- 246 *Megalepthyphantes pseudocollinus* SAARISTO, 1997
- 247 *Meioneta affinis* (KULCZYŃSKI, 1898)
- 248 *Meioneta equestris* (L. KOCH, 1881)
- 249 *Meioneta fuscipalpis* (C.L. KOCH, 1836)
- 250 *Meioneta gulosa* (L. KOCH, 1869)
- 251 *Meioneta innotabilis* (O. P.-CAMBRIDGE, 1863)
- 252 *Meioneta mossica* SCHIKORA, 1993
- 253 *Meioneta rurestris* (C.L. KOCH, 1836)
- 254 *Meioneta saxatilis* (BLACKWALL, 1844)
- 255 *Meioneta mollis* (O. P.-CAMBRIDGE, 1871)
- 256 *Metapanamomops kaestneri* (WIEHLE, 1961)
- 257 *Metopobactrus prominulus* (O. P.-CAMBRIDGE, 1872)
- 258 *Micrargus apertus* (O. P.-CAMBRIDGE, 1870)
- 259 *Micrargus herbigradus* (BLACKWALL, 1854)
- 260 *Micrargus subaequalis* (WESTRING, 1851)
- 261 *Microctenonyx subitaneus* (O. P.-CAMBRIDGE, 1875)
- 262 *Microlinyphia impigra* (O. P.-CAMBRIDGE, 1906)

- 263 *Microlinyphia pusilla* (SUNDEVALL, 1829)
- 264 *Microneta viaria* (BLACKWALL, 1841)
- 265 *Midia midas* (SIMON, 1884)
- 266 *Minicia marginella* (WIDER, 1834)
- 267 *Minyriolus pusillus* (WIDER, 1834)
- 268 *Mioxena blanda* (SIMON, 1884)
- 269 *Moebelia penicillata* (WESTRING, 1851)
- 270 *Mughiphantes mughi* (FICKERT, 1875)
- 271 *Mughiphantes pulcher* (KULCZYŃSKI, 1882)
- 272 *Mughiphantes varians* (KULCZYŃSKI, 1882)
- 273 *Neriere clathrata* (SUNDEVALL, 1829)
- 274 *Neriere emphana* (WALCKENAER, 1842)
- 275 *Neriere furtiva* (O. P.-CAMBRIDGE, 1871)
- 276 *Neriere montana* (CLERCK, 1758)
- 277 *Neriere peltata* (WIDER, 1834)
- 278 *Neriere radiata* (WALCKENAER, 1842)
- 279 *Notioscopus sarcinatus* (O. P.-CAMBRIDGE, 1872)
- 280 *Obscuriphantes obscurus* (BLACKWALL, 1841)
- 281 *Oedothorax agrestis* (BLACKWALL, 1853)
- 282 *Oedothorax apicatus* (BLACKWALL, 1850)
- 283 *Oedothorax fuscus* (BLACKWALL, 1841)
- 284 *Oedothorax gibbifer* (KULCZYŃSKI, 1882)
- 285 *Oedothorax gibbosus* (BLACKWALL, 1841)
- 286 *Oedothorax retusus* (WESTRING, 1851)
- 287 *Oreonetides glacialis* (L. KOCH, 1872)
- 288 *Oreonetides vaginatus* (THORELL, 1872)
- 289 *Oryphantes angulatus* (O. P.-CAMBRIDGE, 1881)
- 290 *Ostearius melanopygius* (O. P.-CAMBRIDGE, 1879)
- 291 *Palliduphantes alutacius* (SIMON, 1884)
- 292 *Palliduphantes insignis* (O. P.-CAMBRIDGE, 1913)
- 293 *Palliduphantes milleri* (STAREGA, 1972)
- 294 *Palliduphantes pallidus* (O. P.-CAMBRIDGE, 1871)
- 295 *Palliduphantes pillichi* (KULCZYŃSKI, 1915)
- 296 *Panamomops mengei* SIMON, 1926
- 297 *Parapelecopsis nemoralis* (BLACKWALL, 1841)
- 298 *Pelecopsis elongata* (WIDER, 1834)
- 299 *Pelecopsis mengei* (SIMON, 1884)
- 300 *Pelecopsis parallela* (WIDER, 1834)
- 301 *Pelecopsis radiccicola* (L. KOCH, 1872)
- 302 *Peponocranium ludicrum* (O. P.-CAMBRIDGE, 1861)
- 303 *Peponocranium orbiculatum* (O. P.-CAMBRIDGE, 1882)
- 304 *Peponocranium praeceps* MILLER, 1943
- 305 *Pityohyphantes phrygianus* (C.L. KOCH, 1836)
- 306 *Pocadicnemis carpatica* (CHYZER, 1894)
- 307 *Pocadicnemis juncea* LOCKET & MILLIDGE, 1953
- 308 *Pocadicnemis pumila* (BLACKWALL, 1841)
- 309 *Poeciloneta variegata* (BLACKWALL, 1841)
- 310 *Porrhomma campbelli* F. P.-CAMBRIDGE, 1894
- 311 *Porrhomma convexum* (WESTRING, 1861)
- 312 *Porrhomma egeria* SIMON, 1884
- 313 *Porrhomma errans* (BLACKWALL, 1841)
- 314 *Porrhomma hebescens* (L. KOCH, 1879)
- 315 *Porrhomma microphthalmum* (O. P.-CAMBRIDGE, 1871)
- 316 *Porrhomma oblitum* (O. P.-CAMBRIDGE, 1871)

- 317 *Porrhomma pallidum* JACKSON, 1913
- 318 *Porrhomma pygmaeum* (BLACKWALL, 1834)
- 319 *Porrhomma rosenhaueri* (L. KOCH, 1872)
- 320 *Pseudomaro aenigmaticus* DENIS, 1966
- 321 *Saaristoa abnormis* (BLACKWALL, 1841)
- 322 *Saaristoa firma* (O. P.-CAMBRIDGE, 1905)
- 323 *Saloca diceros* (O. P.-CAMBRIDGE, 1871)
- 324 *Saloca kulczynskii* MILLER & KRATOCHVÍL, 1939
- 325 *Satilatlas britteni* (JACKSON, 1913)
- 326 *Sauron rayi* (SIMON, 1881)
- 327 *Savignya frontata* BLACKWALL, 1833
- 328 *Scotargus pilosus* SIMON, 1913
- 329 *Scotinotylus antennatus* (O. P.-CAMBRIDGE, 1875)
- 330 *Semljicola faustus* (O. P.-CAMBRIDGE, 1900)
- 331 *Silometopus acutus* HOLM, 1977
- 332 *Silometopus elegans* (O. P.-CAMBRIDGE, 1872)
- 333 *Silometopus incurvatus* (O. P.-CAMBRIDGE, 1873)
- 334 *Silometopus reussi* (THORELL, 1871)
- 335 *Sintula corniger* (BLACKWALL, 1856)
- 336 *Sisicus apertus* (HOLM, 1939)
- 337 *Stemonyphantes lineatus* (LINNAEUS, 1758)
- 338 *Styloctetor romanus* (O. P.-CAMBRIDGE, 1872)
- 339 *Styloctetor stativus* (SIMON, 1881)
- 340 *Syedra gracilis* (MENGE, 1869)
- 341 *Syedra myrmicarum* (KULCZYŃSKI, 1882)
- 342 *Tallusia experta* (O. P.-CAMBRIDGE, 1871)
- 343 *Tapinocyba affinis* (LESSERT, 1907)
- 344 *Tapinocyba biscissa* (O. P.-CAMBRIDGE, 1872)
- 345 *Tapinocyba insecta* (L. KOCH, 1869)
- 346 *Tapinocyba pallens* (O. P.-CAMBRIDGE, 1872)
- 347 *Tapinocyba praecox* (O. P.-CAMBRIDGE, 1873)
- 348 *Tapinocyboides pygmaeus* (MENGE, 1869)
- 349 *Tapinopa longidens* (WIDER, 1834)
- 350 *Taranucnus bihari* FAGE, 1931
- 351 *Taranucnus setosus* (O. P.-CAMBRIDGE, 1863)
- 352 *Tenuiphantes alacris* (BLACKWALL, 1853)
- 353 *Tenuiphantes cristatus* (MENGE, 1866)
- 354 *Tenuiphantes flavipes* (BLACKWALL, 1854)
- 355 *Tenuiphantes mengei* (KULCZYŃSKI, 1887)
- 356 *Tenuiphantes tenebricola* (WIDER, 1834)
- 357 *Tenuiphantes tenuis* (BLACKWALL, 1852)
- 358 *Tenuiphantes zimmermanni* (BERTKAU, 1890)
- 359 *Thyreosthenius biovatus* (O. P.-CAMBRIDGE, 1875)
- 360 *Thyreosthenius parasiticus* (WESTRING, 1851)
- 361 *Tiso vagans* (BLACKWALL, 1834)
- 362 *Tmeticus affinis* (BLACKWALL, 1855)
- 363 *Trematocephalus cristatus* (WIDER, 1834)
- 364 *Trichoncus affinis* KULCZYŃSKI, 1894
- 365 *Trichoncus vasconicus* DENIS, 1944
- 366 *Trichopterna cito* (O. P.-CAMBRIDGE, 1872)
- 367 *Trichopterna thorelli* (WESTRING, 1861)
- 368 *Troxochrus nasutus* SCHENKEL, 1925
- 369 *Troxochrus scabriculus* (WESTRING, 1851)
- 370 *Typhochrestus digitatus* (O. P.-CAMBRIDGE, 1872)



- 371 *Walckenaeria acuminata* BLACKWALL, 1833
- 372 *Walckenaeria alticeps* (DENIS, 1952)
- 373 *Walckenaeria antica* (WIDER, 1834)
- 374 *Walckenaeria atrotibialis* (O. P.-CAMBRIDGE, 1878)
- 375 *Walckenaeria capito* (WESTRING, 1861)
- 376 *Walckenaeria corniculans* (O. P.-CAMBRIDGE, 1875)
- 377 *Walckenaeria cucullata* (C.L. KOCH, 1836)
- 378 *Walckenaeria cuspidata* BLACKWALL, 1833
- 379 *Walckenaeria dysderoides* (WIDER, 1834)
- 380 *Walckenaeria furcillata* (MENGE, 1869)
- 381 *Walckenaeria kochi* (O. P.-CAMBRIDGE, 1872)
- 382 *Walckenaeria mitrata* (MENGE, 1868)
- 383 *Walckenaeria monoceros* (WIDER, 1834)
- 384 *Walckenaeria nodosa* O. P.-CAMBRIDGE, 1873
- 385 *Walckenaeria nudipalpis* (WESTRING, 1851)
- 386 *Walckenaeria obtusa* BLACKWALL, 1836
- 387 *Walckenaeria suspecta* (KULCZYŃSKI, 1882)
- 388 *Walckenaeria unicornis* O. P.-CAMBRIDGE, 1861
- 389 *Walckenaeria vigilax* (BLACKWALL, 1853)
- 390 *Zornella cultrigera* (L. KOCH, 1879)

#### **TETRAGNATHIDAE**

- 391 *Meta menardi* (LATREILLE, 1804)
- 392 *Metellina mengei* (BLACKWALL, 1869)
- 393 *Metellina merianae* (SCOPOLI, 1763)
- 394 *Metellina segmentata* (CLERCK, 1758)
- 395 *Pachygnatha clercki* SUNDEVALL, 1823
- 396 *Pachygnatha degeeri* SUNDEVALL, 1830
- 397 *Pachygnatha listeri* SUNDEVALL, 1830
- 398 *Tetragnatha dearmata* THORELL, 1873
- 399 *Tetragnatha extensa* (LINNAEUS, 1758)
- 400 *Tetragnatha montana* SIMON, 1874
- 401 *Tetragnatha nigrita* LENDL, 1886
- 402 *Tetragnatha obtusa* C.L. KOCH, 1837
- 403 *Tetragnatha pinicola* L. KOCH, 1870
- 404 *Tetragnatha reimoseri* (ROSCA, 1939)
- 405 *Tetragnatha striata* L. KOCH, 1862

#### **ARANEIDAE**

- 406 *Aculepeira ceropegia* (WALCKENAER, 1802)
- 407 *Agalenatea redii* (SCOPOLI, 1763)
- 408 *Araneus alsine* (WALCKENAER, 1802)
- 409 *Araneus angulatus* CLERCK, 1758
- 410 *Araneus diadematus* CLERCK, 1758
- 411 *Araneus marmoreus* CLERCK, 1758
- 412 *Araneus nordmanni* (THORELL, 1870)
- 413 *Araneus quadratus* CLERCK, 1758
- 414 *Araneus saevus* (L. KOCH, 1872)
- 415 *Araneus sturmi* (HAHN, 1831)
- 416 *Araneus triguttatus* (FABRICIUS, 1775)
- 417 *Araniella alpica* (L. KOCH, 1869)
- 418 *Araniella cucurbitina* (CLERCK, 1758)
- 419 *Araniella displicata* (HENTZ, 1847)
- 420 *Araniella inconspicua* (SIMON, 1874)

- 421 *Araniella opisthographa* (KULCZYŃSKI, 1905)
- 422 *Araniella proxima* (KULCZYŃSKI, 1885)
- 423 *Argiope bruennichi* (SCOPOLI, 1772)
- 424 *Cercidia prominens* (WESTRING, 1851)
- 425 *Cyclosa conica* (PALLAS, 1772)
- 426 *Cyclosa oculata* (WALCKENAER, 1802)
- 427 *Gibbaranea bituberculata* (WALCKENAER, 1802)
- 428 *Gibbaranea gibbosa* (WALCKENAER, 1802)
- 429 *Gibbaranea omoeda* (THORELL, 1870)
- 430 *Gibbaranea ullrichi* (HAHN, 1835)
- 431 *Hypsosinga albovittata* (WESTRING, 1851)
- 432 *Hypsosinga heri* (HAHN, 1831)
- 433 *Hypsosinga pygmaea* (SUNDEVALL, 1832)
- 434 *Hypsosinga sanguinea* (C.L. KOCH, 1844)
- 435 *Larinia bonneti* SPASSKY, 1939
- 436 *Larinia jeskovi* MARUSIK, 1986
- 437 *Larinioides cornutus* (CLERCK, 1758)
- 438 *Larinioides ixobolus* (THORELL, 1873)
- 439 *Larinioides patagiatus* (CLERCK, 1758)
- 440 *Larinioides sclopetarius* (CLERCK, 1758)
- 441 *Larinioides suspicax* (O. P.-CAMBRIDGE, 1876)
- 442 *Mangora acalypha* (WALCKENAER, 1802)
- 443 *Neoscona adianta* (WALCKENAER, 1802)
- 444 *Nuctenea silvicultrix* (C.L. KOCH, 1835)
- 445 *Nuctenea umbratica* (CLERCK, 1758)
- 446 *Singa hamata* (CLERCK, 1758)
- 447 *Singa nitidula* C.L. KOCH, 1844
- 448 *Zilla diodia* (WALCKENAER, 1802)
- 449 *Zygiella atrica* (C.L. KOCH, 1845)
- 450 *Zygiella montana* (C.L. KOCH, 1834)
- 451 *Zygiella stroemi* (THORELL, 1870)
- 452 *Zygiella x-notata* (CLERCK, 1758)

#### LYCOSIDAE

- 453 *Acantholycosa lignaria* (CLERCK, 1758)
  
- 454a *Acantholycosa norvegica sudetica* (KOCH, 1875)
- 455 *Alopecosa accentuata* (LATREILLE, 1817)
- 456 *Alopecosa aculeata* (CLERCK, 1758)
- 457 *Alopecosa cuneata* (CLERCK, 1758)
- 458 *Alopecosa cursor* (HAHN, 1831)
- 459 *Alopecosa fabrilis* (CLERCK, 1758)
- 460 *Alopecosa inquilina* (CLERCK, 1758)
- 461 *Alopecosa mariae* (DAHL, 1908)
- 462 *Alopecosa pinetorum* (THORELL, 1856)
- 463 *Alopecosa pulverulenta* (CLERCK, 1758)
- 464 *Alopecosa schmidti* (HAHN, 1835)
- 465 *Alopecosa solitaria* (HERMAN, 1876)
- 466 *Alopecosa striatipes* (C.L. KOCH, 1837)
- 467 *Alopecosa taeniata* (C.L. KOCH, 1835)
- 468 *Alopecosa trabalis* (CLERCK, 1758)
- 469 *Arctosa alpigena alpigena* (DOLESCHALL, 1852)
  
- 469a *Arctosa alpigena lamperti* DAHL, 1908

- 470 *Arctosa cinerea* (FABRICIUS, 1777)
- 471 *Arctosa figurata* (SIMON, 1876)
- 472 *Arctosa leopardus* (SUNDEVALL, 1833)
- 473 *Arctosa lutetiana* (SIMON, 1876)
- 474 *Arctosa maculata* (HAHN, 1822)
- 475 *Arctosa perita* (LATREILLE, 1799)
- 476 *Arctosa stigmosa* (THORELL, 1875)
- 477 *Aulonia albimana* (WALCKENAER, 1805)
- 478 *Hygrolycosa rubrofasciata* (OHLERT, 1865)
- 479 *Pardosa agrestis* (WESTRING, 1861)
- 480 *Pardosa agricola* (THORELL, 1856)
- 481 *Pardosa albatula* (ROEWER, 1951)
- 482 *Pardosa amentata* (CLERCK, 1758)
- 483 *Pardosa bifasciata* (C.L. KOCH, 1834)
- 484 *Pardosa evelinae* WUNDERLICH, 1984
- 485 *Pardosa ferruginea* (L. KOCH, 1870)
- 486 *Pardosa hortensis* (THORELL, 1872)
- 487 *Pardosa hyperborea* (THORELL, 1870)
- 488 *Pardosa lugubris* (WALCKENAER, 1802)
- 489 *Pardosa maisa* HIPPA & MANNILA, 1982
- 490 *Pardosa monticola* (CLERCK, 1758)
- 491 *Pardosa morosa* (L. KOCH, 1870)
- 492 *Pardosa nigra* (C.L. KOCH, 1834)
- 493 *Pardosa nigriceps* (THORELL, 1856)
- 494 *Pardosa paludicola* (CLERCK, 1758)
- 495 *Pardosa palustris* (LINNAEUS, 1758)
- 496 *Pardosa prativaga* (L. KOCH, 1870)
- 497 *Pardosa pullata* (CLERCK, 1758)
- 498 *Pardosa riparia* (C.L. KOCH, 1833)
- 499 *Pardosa saltans* TÖPFER-HOFMANN, 2000
- 500 *Pardosa saltuaria* (L. KOCH, 1870)
- 501 *Pardosa schenkeli* LESSERT, 1904
- 502 *Pardosa sordidata* (THORELL, 1875)
- 503 *Pardosa sphagnicola* (DAHL, 1908)
- 504 *Pardosa wagleri* (HAHN, 1822)
- 505 *Pirata hygrophilus* THORELL, 1872
- 506 *Pirata insularis* EMERTON, 1885
- 507 *Pirata knorri* (SCOPOLI, 1763)
- 508 *Pirata latitans* (BLACKWALL, 1841)
- 509 *Pirata piraticus* (CLERCK, 1758)
- 510 *Pirata piscatorius* (CLERCK, 1758)
- 511 *Pirata tenuitarsis* SIMON, 1876
- 512 *Pirata uliginosus* (THORELL, 1856)
- 513 *Trochosa robusta* (SIMON, 1876)
- 514 *Trochosa ruricola* (DE GEER, 1778)
- 515 *Trochosa spinipalpis* (F. P.-CAMBRIDGE, 1895)
- 516 *Trochosa terricola* THORELL, 1856
- 517 *Xerolycosa miniata* (C.L. KOCH, 1834)
- 518 *Xerolycosa nemoralis* (WESTRING, 1861)

#### **PISAUROIDAE**

- 519 *Dolomedes fimbriatus* (CLERCK, 1758)
- 520 *Dolomedes plantarius* (CLERCK, 1758)
- 521 *Pisaura mirabilis* (CLERCK, 1758)

**OXYOPIDAE**

- 522 *Oxyopes ramosus* (PANZER, 1804)

**ZORIDAE**

- 523 *Zora armillata* SIMON, 1878  
 524 *Zora distincta* KULCZYŃSKI, 1915  
 525 *Zora manicata* SIMON, 1878  
 526 *Zora nemoralis* (BLACKWALL, 1861)  
 527 *Zora silvestris* KULCZYŃSKI, 1897  
 528 *Zora spinimana* (SUNDEVALL, 1833)

**AGELENIDAE**

- 529 *Agelena gracilens* C.L. KOCH, 1841  
 530 *Agelena labyrinthica* (CLERCK, 1758)  
 531 *Histocona torpida* (C.L. KOCH, 1834)  
 532 *Malthonica campestris* (C. L. KOCH, 1834)  
 533 *Malthonica ferruginea* (PANZER, 1804)  
 534 *Malthonica silvestris* (L. KOCH, 1872)  
 535 *Tegenaria agrestis* (WALCKENAER, 1802)  
 536 *Tegenaria atrica* C.L. KOCH, 1843  
 537 *Tegenaria domestica* (CLERCK, 1758)  
 538 *Textrix denticulata* (OLIVIER, 1789)

**CYBAEIDAE**

- 539 *Argyroneta aquatica* (CLERCK, 1758)  
 540 *Cybaeus angustiarum* L. KOCH, 1868  
 541 *Cybaeus tetricus* (C.L. KOCH, 1839)

**HAHNIIDAE**

- 542 *Antistea elegans* (BLACKWALL, 1841)  
 543 *Cryphoeca carpatica* HERMAN, 1876  
 544 *Cryphoeca silvicola* (C.L. KOCH, 1834)  
 545 *Hahnica candida* SIMON, 1875  
 546 *Hahnica helveola* SIMON, 1875  
 547 *Hahnica nava* (BLACKWALL, 1841)  
 548 *Hahnica ononidum* SIMON, 1875  
 549 *Hahnica parva* KULCZYŃSKI, 1882  
 550 *Hahnica picta* KULCZYŃSKI, 1897  
 551 *Hahnica pusilla* C.L. KOCH, 1841

**DICTYNIDAE**

- 552 *Altella biuncata* (MILLER, 1949)  
 553 *Archaeodictyna ammophila* (MENGE, 1871)  
 554 *Archaeodictyna consecuta* (O. P.-CAMBRIDGE, 1872)  
 555 *Argenna albopunctata* (MENGE, 1869)  
 556 *Argenna subnigra* (O. P.-CAMBRIDGE, 1861)  
 557 *Brommella falcigera* (BALOGH, 1935)  
 558 *Cicurina cicur* (FABRICIUS, 1793)  
 559 *Dictyna arundinacea* (LINNAEUS, 1758)  
 560 *Dictyna latens* (FABRICIUS, 1775)  
 561 *Dictyna major* MENGE, 1869  
 562 *Dictyna pusilla* THORELL, 1856  
 563 *Dictyna uncinata* THORELL, 1856

- 564 *Emblyna brevidens* (KULCZYŃSKI, 1897)
- 565 *Lathys humilis* (BLACKWALL, 1855)
- 566 *Lathys stigmatisata* (MENGE, 1869)
- 567 *Mastigusa arietina* (THORELL, 1871)
- 568 *Mastigusa macrophthalma* (KULCZYŃSKI, 1897)
- 569 *Nigma flavescens* (WALCKENAER, 1830)
- 570 *Nigma walckenaeri* (ROEWER, 1951)

#### **AMAUROBIIDAE**

- 571 *Amaurobius fenestralis* (STRÖM, 1768)
- 572 *Amaurobius ferox* (WALCKENAER, 1830)
- 573 *Amaurobius jugorum* L. KOCH, 1868
- 574 *Amaurobius similis* (BLACKWALL, 1861)
- 575 *Callobius claustrarius* (HAHN, 1833)
- 576 *Coelotes atropos* (WALCKENAER, 1830)
- 577 *Eurocoelotes inermis* (L. KOCH, 1855)
- 578 *Coelotes terrestris* (WIDER, 1834)

#### **TITANOECIDAE**

- 579 *Titanoeca psammophila* WUNDERLICH, 1993
- 580 *Titanoeca quadriguttata* (HAHN, 1833)

#### **MITURGIDAE**

- 581 *Cheiracanthium campestre* LOHMANDER, 1944
- 582 *Cheiracanthium erraticum* (WALCKENAER, 1802)
- 583 *Cheiracanthium montanum* L. KOCH, 1878
- 584 *Cheiracanthium oncognathum* THORELL, 1871
- 585 *Cheiracanthium pennyi* O. P.-CAMBRIDGE, 1873
- 586 *Cheiracanthium punctorium* (VILLERS, 1789)
- 587 *Cheiracanthium virescens* (SUNDEVALL, 1833)

#### **ANYPHAENIDAE**

- 588 *Anyphaena accentuata* (WALCKENAER, 1802)

#### **LIOCRANIDAE**

- 589 *Agroeca brunnea* (BLACKWALL, 1833)
- 590 *Agroeca cuprea* MENGE, 1873
- 591 *Agroeca dentigera* KULCZYŃSKI, 1913
- 592 *Agroeca lusatica* (L. KOCH, 1875)
- 593 *Agroeca proxima* (O. P.-CAMBRIDGE, 1871)
- 594 *Apostenus fuscus* WESTRING, 1851
- 595 *Liocranoeca striata* (KULCZYŃSKI, 1882)
- 596 *Liocranum rupicola* (WALCKENAER, 1830)
- 597 *Liocranum rutilans* (THORELL, 1875)
- 598 *Scotina celans* (BLACKWALL, 1841)
- 599 *Scotina gracillipes* (BLACKWALL, 1859)
- 600 *Scotina palliardi* (L. KOCH, 1881)

#### **CLUBIONIDAE**

- 601 *Clubiona abdita* L. KOCH, 1876
- 602 *Clubiona alpicola* KULCZYŃSKI, 1882
- 603 *Clubiona brevipes* BLACKWALL, 1841
- 604 *Clubiona caerulescens* L. KOCH, 1867
- 605 *Clubiona comta* C.L. KOCH, 1839

- 606 *Clubiona corticalis* (WALCKENAER, 1802)
- 607 *Clubiona diversa* O. P.-CAMBRIDGE, 1862
- 608 *Clubiona frisia* WUNDERLICH & SCHÜTT, 1995
- 609 *Clubiona frutetorum* L. KOCH, 1866
- 610 *Clubiona genevensis* L. KOCH, 1866
- 611 *Clubiona germanica* THORELL, 1871
- 612 *Clubiona juvenis* SIMON, 1878
- 613 *Clubiona kulczynskii* LESSERT, 1905
- 614 *Clubiona lutescens* WESTRING, 1851
- 615 *Clubiona marmorata* L. KOCH, 1866
- 616 *Clubiona neglecta* O. P.-CAMBRIDGE, 1862
- 617 *Clubiona norvegica* STRAND, 1900
- 618 *Clubiona pallidula* (CLERCK, 1758)
- 619 *Clubiona phragmitis* C.L. KOCH, 1843
- 620 *Clubiona reclusa* O. P.-CAMBRIDGE, 1863
- 621 *Clubiona rosserae* LOCKET, 1953
- 622 *Clubiona similis* L. KOCH, 1867
- 623 *Clubiona stagnatilis* KULCZYŃSKI, 1897
- 624 *Clubiona subsultans* THORELL, 1875
- 625 *Clubiona subtilis* L. KOCH, 1867
- 626 *Clubiona terrestris* WESTRING, 1851
- 627 *Clubiona trivialis* C.L. KOCH, 1843

#### **CORINNIDAE**

- 628 *Phrurolithus festivus* (C.L. KOCH, 1835)
- 629 *Phrurolithus minimus* C.L. KOCH, 1839
- 630 *Phrurolithus pullatus* KULCZYŃSKI, 1897

#### **ZODARIIDAE**

- 631 *Zodarion germanicum* (C.L. KOCH, 1837)
- 632 *Zodarion rubidum* SIMON, 1914

#### **GNAPHOSIDAE**

- 633 *Berlandina cinerea* (MENGE, 1872)
- 634 *Callilepis nocturna* (LINNAEUS, 1758)
- 635 *Drassodes lapidosus* (WALCKENAER, 1802)
- 636 *Drassodes pubescens* (THORELL, 1856)
- 637 *Drassodes cupreus* (BLACKWALL, 1834)
- 638 *Drassyllus lutetianus* (L. KOCH, 1866)
- 639 *Drassyllus praeficus* (L. KOCH, 1866)
- 640 *Drassyllus pumilus* (C.L. KOCH, 1839)
- 641 *Drassyllus pusillus* (C.L. KOCH, 1833)
- 642 *Gnaphosa bicolor* (HAHN, 1833)
- 643 *Gnaphosa lucifuga* (WALCKENAER, 1802)
- 644 *Gnaphosa lugubris* (C.L. KOCH, 1839)
- 645 *Gnaphosa microps* HOLM, 1939
- 646 *Gnaphosa montana* (L. KOCH, 1866)
- 647 *Gnaphosa muscorum* (L. KOCH, 1866)
- 648 *Gnaphosa nigerrima* L. KOCH, 1878
- 649 *Gnaphosa opaca* HERMAN, 1879
- 650 *Haplodrassus cognatus* (WESTRING, 1861)
- 651 *Haplodrassus dalmatensis* (L. KOCH, 1866)
- 652 *Haplodrassus kulczynskii* LOHMANDER, 1942
- 653 *Haplodrassus minor* (O. P.-CAMBRIDGE, 1879)

- 654 Haplodrassus moderatus (KULCZYŃSKI, 1897)
- 655 Haplodrassus signifer (C.L. KOCH, 1839)
- 656 Haplodrassus soerenseni (STRAND, 1900)
- 657 Haplodrassus silvestris (BLACKWALL, 1833)
- 658 Haplodrassus umbratilis (L. KOCH, 1866)
- 659 Kishidaia conspicua (L. KOCH, 1866)
- 660 Micaria formicaria (SUNDEVALL, 1832)
- 661 Micaria fulgens (WALCKENAER, 1802)
- 662 Micaria nivosa L. KOCH, 1866
- 663 Micaria pulicaria (SUNDEVALL, 1832)
- 664 Micaria rossica THORELL, 1875
- 665 Micaria silesiaca L. KOCH, 1875
- 666 Micaria subopaca WESTRING, 1861
- 667 Phaeoedus braccatus (L. KOCH, 1866)
- 668 Poecilochroa variana (C.L. KOCH, 1839)
- 669 Scotophaeus blackwalli (THORELL, 1871)
- 670 Scotophaeus quadripunctatus (LINNAEUS, 1758)
- 671 Scotophaeus scutulatus (L. KOCH, 1866)
- 672 Sosticus loricatus (L. KOCH, 1866)
- 673 Trachyzelotes pedestris (C.L. KOCH, 1837)
- 674 Zelotes aeneus (SIMON, 1878)
- 675 Zelotes apricorum (L. KOCH, 1876)
- 676 Zelotes clivicola (L. KOCH, 1870)
- 677 Zelotes electus (C.L. KOCH, 1839)
- 678 Zelotes exiguus (MÜLLER & SCHENKEL, 1895)
- 679 Zelotes latreillei (SIMON, 1878)
- 680 Zelotes longipes (L. KOCH, 1866)
- 681 Zelotes petrensis (C.L. KOCH, 1839)
- 682 Zelotes puritanus CHAMBERLIN, 1922
- 683 Zelotes pygmaeus MILLER, 1943
- 684 Zelotes subterraneus (C.L. KOCH, 1833)

#### **SPARASSIDAE**

- 685 Micrommata roseum (CLERCK, 1758)

#### **PHILODROMIDAE**

- 686 Philodromus albidus KULCZYŃSKI, 1911
- 687 Philodromus aureolus (CLERCK, 1758)
- 688 Philodromus buxi SIMON, 1884
- 689 Philodromus cespitum (WALCKENAER, 1802)
- 690 Philodromus collinus C.L. KOCH, 1835
- 691 Philodromus corticinus (C.L. KOCH, 1837)
- 692 Philodromus dispar (WALCKENAER, 1826)
- 693 Philodromus emarginatus (SCHRANK, 1803)
- 694 Philodromus fallax SUNDEVALL, 1833
- 695 Philodromus fuscomarginatus (DE GEER, 1778)
- 696 Philodromus histrio (LATREILLE, 1819)
- 697 Philodromus longipalpis SIMON, 1870
- 698 Philodromus margaritatus (CLERCK, 1758)
- 699 Philodromus poecilus (THORELL, 1872)
- 700 Philodromus praedatus O. P.-CAMBRIDGE, 1871
- 701 Philodromus rufus WALCKENAER, 1826
- 702 Philodromus vagulus SIMON, 1875
- 703 Thanatus atratus SIMON, 1875

- 704 *Thanatus arenarius* L. KOCH, 1872
- 705 *Thanatus formicinus* (CLERCK, 1758)
- 706 *Thanatus pictus* L. KOCH, 1881
- 707 *Thanatus sabulosus* (MENGE, 1874)
- 708 *Thanatus striatus* C.L. KOCH, 1845
- 709 *Thanatus vulgaris* SIMON, 1870
- 710 *Tibellus maritimus* (MENGE, 1874)
- 711 *Tibellus oblongus* (WALCKENAER, 1802)

#### **THOMISIDAE**

- 712 *Coriarachne depressa* (C.L. KOCH, 1837)
- 713 *Cozyptila blackwalli* (SIMON, 1875)
- 714 *Diaea dorsata* (FABRICIUS, 1775)
- 715 *Ebrechtella tricuspida* (FABRICIUS, 1775)
- 716 *Heriaeus graminicola* (DOLESCHAL, 1852)
- 717 *Heriaeus mellottei* SIMON, 1886
- 718 *Misumena vatia* (CLERCK, 1758)
- 719 *Ozyptila atomaria* (PANZER, 1801)
- 720 *Ozyptila brevipes* (HAHN, 1826)
- 721 *Ozyptila claveata* (WALCKENAER, 1837)
- 722 *Ozyptila gertschi* KURATA, 1944
- 723 *Ozyptila praticola* (C.L. KOCH, 1837)
- 724 *Ozyptila pullata* (THORELL, 1875)
- 725 *Ozyptila rauda* SIMON, 1875
- 726 *Ozyptila scabricula* (WESTRING, 1851)
- 727 *Ozyptila trux* (BLACKWALL, 1846)
- 728 *Pistius truncatus* (PALLAS, 1772)
- 729 *Synema globosum* (FABRICIUS, 1775)
- 730 *Thomisus onustus* WALCKENAER, 1805
- 731 *Tmarus piger* (WALCKENAER, 1802)
- 732 *Xysticus acerbus* (THORELL, 1872)
- 733 *Xysticus alpicola* KULCZYŃSKI, 1882
- 734 *Xysticus audax* (SCHRANK, 1803)
- 735 *Xysticus bifasciatus* C.L. KOCH, 1837
- 736 *Xysticus cristatus* (CLERCK, 1758)
- 737 *Xysticus erraticus* (BLACKWALL, 1834)
- 738 *Xysticus ferrugineus* MENGE, 1876
- 739 *Xysticus gallicus* SIMON, 1875
- 740 *Xysticus kochi* THORELL, 1872
- 741 *Xysticus lanio* C.L. KOCH, 1835
- 742 *Xysticus lineatus* (WESTRING, 1851)
- 743 *Xysticus luctator* L. KOCH, 1870
- 744 *Xysticus luctuosus* (BLACKWALL, 1836)
- 745 *Xysticus ninnii* THORELL, 1872
- 746 *Xysticus obscurus* COLLETT, 1877
- 747 *Xysticus robustus* (HAHN, 1832)
- 748 *Xysticus sabulosus* (HAHN, 1832)
- 749 *Xysticus striatipes* L. KOCH, 1870
- 750 *Xysticus ulmi* (HAHN, 1831)

#### **SALTICIDAE**

- 751 *Aelurillus v-insignitus* (CLERCK, 1758)
- 752 *Asianellus festivus* (C.L. KOCH, 1834)
- 753 *Ballus chalybeius* (WALCKENAER, 1802)



- 754 *Carrhotus xanthogramma* (LATREILLE, 1819)
- 755 *Dendryphantes hastatus* (CLERCK, 1758)
- 756 *Dendryphantes rudis* (SUNDEVALL, 1832)
- 757 *Euophrys frontalis* (WALCKENAER, 1802)
- 758 *Evarcha arcuata* (CLERCK, 1758)
- 759 *Evarcha falcata* (CLERCK, 1758)
- 760 *Evarcha laetabunda* C.L. KOCH, 1848)
- 761 *Hasarius adansoni* (AUDOUIN, 1826)
- 762 *Heliophanus aeneus* (HAHN, 1831)
- 763 *Heliophanus auratus* C.L. KOCH, 1835
- 764 *Heliophanus cupreus* (WALCKENAER, 1802)
- 765 *Heliophanus dampfi* SCHENKEL, 1923
- 766 *Heliophanus dubius* C.L. KOCH, 1835
- 767 *Heliophanus flavipes* (HAHN, 1831)
- 768 *Heliophanus patagiatus* THORELL, 1875
- 769 *Icius hamatus* (C. L. KOCH, 1846)
- 770 *Leptorchestes berlinensis* (C.L. KOCH, 1846)
- 771 *Marpissa muscosa* (CLERCK, 1758)
- 772 *Marpissa pomatia* (WALCKENAER, 1802)
- 773 *Marpissa radiata* (GRUBE, 1859)
- 774 *Myrmarachne formicaria* (DE GEER, 1778)
- 775 *Neon levis* (SIMON, 1871)
- 776 *Neon reticulatus* (BLACKWALL, 1853)
- 777 *Neon valentulus* FALCONER, 1912
- 778 *Pellenes nigrociliatus* (SIMON, 1875)
- 779 *Pellenes tripunctatus* (WALCKENAER, 1802)
- 780 *Philaeus chrysops* (PODA, 1761)
- 781 *Phlegra fasciata* (HAHN, 1826)
- 782 *Pseudeuophrys erratica* (WALCKENAER, 1826)
- 783 *Pseudeuophrys lanigera* (SIMON, 1871)
- 784 *Pseudeuophrys obsoleta* (SIMON, 1868)
- 785 *Pseudicius encarpatus* (WALCKENAER, 1802)
- 786 *Salticus cingulatus* (PANZER, 1797)
- 787 *Salticus scenicus* (CLERCK, 1758)
- 788 *Salticus zebraneus* (C.L. KOCH, 1837)
- 789 *Sibianor aurocinctus* (OHLERT, 1865) )
- 790 *Sitticus caricis* (WESTRING, 1861)
- 791 *Sitticus distinguendus* (SIMON, 1868)
- 792 *Sitticus dzieduszyckii* (L. KOCH, 1870)
- 793 *Sitticus floricola* (C.L. KOCH, 1837)
- 794 *Sitticus inexpectus* LOGUNOV & KRONESTEDT, 1997
- 795 *Sitticus penicillatus* (SIMON, 1875)
- 796 *Sitticus pubescens* (FABRICIUS, 1775)
- 797 *Sitticus rupicola* (C.L. KOCH, 1837)
- 798 *Sitticus saltator* (O. P.-CAMBRIDGE, 1868)
- 799 *Sitticus saxicola* (C.L. KOCH, 1837)
- 800 *Sitticus terebratus* (CLERCK, 1758)
- 801 *Sitticus zimmermanni* (SIMON, 1877)
- 802 *Synageles hilarulus* (C.L. KOCH, 1851)
- 803 *Synageles venator* (LUCAS, 1836)
- 804 *Talavera aequipes* (O. P.-CAMBRIDGE, 1871)
- 805 *Talavera monticola* (KULCZYŃSKI, 1884)
- 806 *Talavera parvistyla* LOGUNOV & KRONESTEDT 2003
- 807 *Talavera petrensis* (C.L. KOCH, 1837)

808 *Talavera thorelli* (KULCZYŃSKI, 1891)

809 *Yllenus arenarius* MENGE, 1868