

Checklist of Spiders (Araneae) in Sweden

[Preliminary version February 2001]

TORBJÖRN KRONESTEDT

Department of Entomology, Swedish Museum of Natural History, Stockholm

Being a preliminary version, I gratefully ask for amendments, changes, additions or other improvements

Introduction

This is a preliminary version of a checklist of spider species reported from Sweden up to present. The records in the published literature (also in press) have been considered and the list presently includes the names of 704 species. A few names mentioned in the literature have been omitted because the nominal species as such are in need of revision, or the identification of the published material from Sweden is in need of verification. In 1955, Tullgren described some linyphiid (erigonine) species as new. Most of these names have been found to be junior synonyms, referring to species known from Sweden by other records. Two of these names (marked with a blue ●●), refer to species which are known from Sweden only by Tullgren's names though already known to be junior, not yet formally established synonyms. The presence of some more species in the country is known by personal communication but these will be included in a later version after they have been published in print. Records of casual introductions have been omitted. It is unavoidable that there are uncertainties about the presence of some of the species, and a list like this will always have a provisional character.

Though several publications have been devoted to spiders in Sweden during the last centuries, no inclusive account of all spiders species found throughout Sweden has appeared since the book by Westring in 1861, in which 308 species by then known were treated (a number of which were considered new to science). This book was soon followed by the more or less extensive "remarks" by Thorell (1870-73) to each of the species treated by Westring. Many of the specific names used by Westring, and a part of the ones introduced by Westring (a number of which already in 1851), have later been synonymized due to priority, or allocated into other genera.

In the present checklist, the order of the families, the assignment of genera to the respective family, and the names of the species are listed according to Platnick (2000).

An annotated checklist of spider species in Sweden (with notes on distribution etc.) is in preparation.

Photos in the list (all © T. Kronestedt) are clickable for larger size.

Species appearing in the 2000 Red List of Swedish Species (Gärdenfors 2000) are marked with a "●●".

Please refer to this version as: Kronestedt, T.: Checklist of spiders (Araneae) in Sweden. Version 2001-02-15.

References

Gärdenfors, U. (ed.) 2000. Rödlistade arter i Sverige 2000 - The 2000 Red List of Swedish Species. ArtDatabanken, SLU, Uppsala. 397 pp.

Platnick, N. I. 2000. The World Spider Catalogue. American Museum of Natural History, New York. URL: <http://research.amnh.org/entomology/index.html>

Thorell, T. 1870-73. Remarks on synonyms of European spiders. Upsala. 645 pp.

Tullgren, A. 1955. Zur Kenntnis schwedischer Erigoniden. - Ark. Zool. (2) 7: 295-389.

Westring, N. 1851. Förteckning öfver de till närvarande tid kände, i Sverige förekommande spindelarter, utgörande ett antal af 253, däraf 132 äro nya för den svenska faunan. - Göteb. K. Vet. Vitth.-Samh. Handl. 2: 25-62.

Westring, N. 1861. Araneae Svecicae. Gothoburgi. 615 pp.

Families represented in Sweden (with number of species in the species list).

Family	Number of species	Family	Number of species
ATYPIDAE	1	AGELENIDAE	5
PHOLCIDAE	2	CYBAEIDAE	1
SEGESTRIIDAE	2	HAHNIDAE	7
DYSDERIDAE	1	DICTYNIDAE	18
MIMETIDAE	2	AMAUROBIDAE	5
ERESIDAE	1	TITANOECIDAE	2
ULOBORIDAE	2	MITURGIDAE	7
NESTICIDAE	1	ANYPHAENIDAE	1
THERIDIIDAE	53	LIOCRANIDAE	13
THERIDIOSOMATIDAE	1	CLUBIONIDAE	21
LINYPHIIDAE	293	GNAPHOSIDAE	50
TETRAGNATHIDAE	14	ZORIDAE	6
ARANEIDAE	41	SPARASSIDAE	1
LYCOSIDAE	56	PHILODROMIDAE	19
PISAUROIDAE	3	THOMISIDAE	29
OXYOPIDAE	1	SALTICIDAE	45
		Total	704 species

Species List

ATYPIDAE

- *Atypus affinis* Eichwald, 1830

PHOLCIDAE

- *Pholcus phalangioides* (Fuesslin, 1775)
- *Psilochorus simoni* (Berland, 1911)

SEGESTRIIDAE

- *Segestria bavarica* C. L. Koch, 1843
- *Segestria senoculata* (Linnaeus, 1758)



Segestria senoculata female

DYSDERIDAE

- *Harpactea hombergi* (Scopoli, 1763)

MIMETIDAE

- *Ero cambridgei* Kulczynski, 1911
- *Ero furcata* (Villers, 1789)

ERESIDAE

- *Eresus sandaliatus* (Martini et Goeze, 1778)

ULOBORIDAE

- *Hyptiotes paradoxus* (C. L. Koch, 1834)
- *Uloborus plumipes* Lucas, 1846

*Hyptiotes paradoxus* female**NESTICIDAE**

- *Nesticus cellulanus* (Clerck, 1757)

THERIDIIDAE

- *Achaeearanea lunata* (Clerck, 1757)
- *Achaeearanea ohlerti* (Thorell, 1870)
- *Achaeearanea riparia* (Blackwall, 1834)
- *Achaeearanea simulans* (Thorell, 1875)
- *Achaeearanea tepidariorum* (C. L. Koch, 1841)
- *Anelosimus vittatus* (C. L. Koch, 1836)
- *Crustulina guttata* (Wider, 1834)
- *Crustulina sticta* (O. P.-Cambridge, 1861)
- *Dipoena braccata* (C. L. Koch, 1841)
- *Dipoena inornata* (O. P.-Cambridge, 1861)
- *Dipoena melanogaster* (C. L. Koch, 1837)
- *Dipoena prona* (Menge, 1868)
- *Dipoena torva* (Thorell, 1875)
- *Enoplognatha latimana* Hippa et Oksala, 1982
- *Enoplognatha mordax* (Thorell, 1875)
- *Enoplognatha oelandica* (Thorell, 1875)
- *Enoplognatha ovata* (Clerck, 1757)
- *Enoplognatha thoracica* (Hahn, 1833)
- *Episinus angulatus* (Blackwall, 1836)
- *Episinus truncatus* Latreille, 1809
- *Euryopis flavomaculata* (C. L. Koch, 1836)
- *Euryopis laeta* (Westring, 1861)
- *Lasaeola tristis* (Hahn, 1833)
- *Neottiura bimaculata* (Linnaeus, 1767)
- *Paidiscura pallens* (Blackwall, 1834)
- *Pholcomma gibbum* (Westring, 1851)
- *Robertus arundineti* (O. P.-Cambridge, 1871)
- *Robertus heydemanni* Wiehle, 1965
- *Robertus insignis* O. P.-Cambridge, 1907
- *Robertus lividus* (Blackwall, 1836)
- *Robertus lyrifer* Holm, 1939
- *Robertus neglectus* (O. P.-Cambridge, 1871)
- *Robertus scoticus* Jackson, 1914
- *Rugathodes bellicosus* (Simon, 1873)
- *Rugathodes instabilis* (O. P.-Cambridge, 1871)
- *Simitidion simile* (L. Koch, 1836)
- *Steatoda albomaculata* (De Geer, 1778)
- *Steatoda bipunctata* (Linnaeus, 1758)
- *Steatoda castanea* (Clerck, 1757)
- *Steatoda grossa* (C. L. Koch, 1838)
- *Steatoda phalerata* (Panzer, 1801)
- *Theonoe minutissima* (O. P.-Cambridge, 1879)
- *Theridion conigerum* Simon, 1914
- *Theridion familiare* O. P.-Cambridge, 1871
- *Theridion hemerobium* Simon, 1914
- *Theridion impressum* L. Koch, 1881
- *Theridion mystaceum* L. Koch, 1870
- *Theridion pictum* (Walckenaer, 1802)
- *Theridion pinastri* L. Koch, 1872
- *Theridion sisyphium* (Clerck, 1757)
- *Theridion tinctum* (Walckenaer, 1802)
- *Theridion varians* Hahn, 1833
- *Thymoites bellissimus* (L. Koch, 1879)

*Theridion sisyphium* female**THERIDIOSOMATIDAE**

- *Theridiosoma gemmosum* (L. Koch, 1877)

LINYPHIIDAE

- *Abacoproeces saltuum* (L. Koch, 1872)
- *Acartauchenius scurrilis* (O. P.-Cambridge, 1872)
- *Agyneta cauta* (O. P.-Cambridge, 1902)
- *Agyneta conigera* (O. P.-Cambridge, 1863)
- *Agyneta decora* (O. P.-Cambridge, 1871)
- *Agyneta olivacea* (Emerton, 1882)
- *Agyneta ramosa* Jackson, 1912
- *Agyneta subtilis* (O. P.-Cambridge, 1863)
- *Agyneta suecica* Holm, 1950
- *Allomengea scopigera* (Grube, 1859)
- *Allomengea vidua* (L. Koch, 1879)
- *Anguliphantes angulipalpis* (Westring, 1851)
- *Aphileta misera* (O. P.-Cambridge, 1882)
- *Araeoncus crassiceps* (Westring, 1861)
- *Araeoncus humilis* (Blackwall, 1841)
- *Asthenargus paganus* (Simon, 1884)
- *Baryphyma gowerense* (Lockett, 1965)
- *Baryphyma pini* (Holm, 1950)
- *Baryphyma pratense* (Blackwall, 1861)
- *Baryphyma trifrons* (O. P.-Cambridge, 1863)
- *Bathyphantes approximatus* (O. P.-Cambridge, 1871)
- *Bathyphantes gracilis* (Blackwall, 1841)
- *Bathyphantes nigrinus* (Westring, 1851)
- *Bathyphantes parvulus* (Westring, 1851)
- *Bathyphantes reprobus* (Kulczynski, 1916)
- *Bathyphantes setiger* F. O. P.-Cambridge, 1894
- *Bathyphantes similis* Kulczynski, 1894
- *Bolyphantes alticeps* (Sundevall, 1833)
- *Bolyphantes index* (Thorell, 1856)
- *Bolyphantes luteolus* (Blackwall, 1833)
- *Carorita limnaea* (Crosby & Bishop, 1927)
- *Carorita paludosa* Duffey, 1971
- *Centromerita bicolor* (Blackwall, 1833)
- *Centromerita concinna* (Thorell, 1875)
- *Centromerus arcanus* (O. P.-Cambridge, 1873)
- *Centromerus brevivulvatus* Dahl, 1912
- *Centromerus incilium* (L. Koch, 1881)
- *Centromerus levitarsis* (Simon, 1884)
- *Centromerus pabulator* (O. P.-Cambridge, 1875)
- *Centromerus persimilis* (O. P.-Cambridge, 1912)
- *Centromerus prudens* (O. P.-Cambridge, 1873)
- *Centromerus semiater* (L. Koch, 1879)
- *Centromerus sylvaticus* (Blackwall, 1841)
- *Ceratinella brevipes* (Westring, 1851)
- *Ceratinella brevis* (Wider, 1834)
- *Ceratinella scabrosa* (O. P.-Cambridge, 1871)
- *Ceratinella wideri* (Thorell, 1871)
- *Cnephalocotes obscurus* (Blackwall, 1834)
- *Collinsia holmgreni* (Thorell, 1871)
- *Collinsia inerrans* (O. P.-Cambridge, 1885)
- *Collinsia spetsbergensis* (Thorell, 1871)
- *Decipiphantes decipiens* (L. Koch, 1879)
- *Dicymbium nigrum* (Blackwall, 1834)
- *Dicymbium tibiale* (Blackwall, 1836)
- *Diplocentria bidentata* (Emerton, 1882)
- *Diplocentria rectangulata* (Emerton, 1915)
- *Diplocephalus cristatus* (Blackwall, 1833)
- *Diplocephalus dentatus* Tullgren, 1955
- *Diplocephalus latifrons* (O. P.-Cambridge, 1863)
- *Diplocephalus permixtus* (O. P.-Cambridge, 1871)
- *Diplocephalus picinus* (Blackwall, 1841)
- *Diplostyla concolor* (Wider, 1834)
- *Dismodicus bifrons* (Blackwall, 1841)
- *Dismodicus elevatus* (C. L. Koch, 1838)
- *Donacochara speciosa* (Thorell, 1875)
- *Drapetisca socialis* (Sundevall, 1833)
- *Drepanotylus borealis* Holm, 1945
- *Drepanotylus uncatatus* (O. P.-Cambridge, 1873)
- *Entelecara acuminata* (Wider, 1834)
- *Entelecara congenera* (O. P.-Cambridge, 1879)
- *Entelecara depilata* Tullgren, 1955
- *Entelecara erythropus* (Westring, 1851)
- *Entelecara forsslundi* Tullgren, 1955
- *Erigone arctica* (White, 1852)
- *Erigone atra* Blackwall, 1833
- *Erigone dentigera* O. P.-Cambridge, 1874

*Linyphia triangularis* female*Pityohyphantes phrygianus* female

- *Erigone dentipalpis* (Wider, 1834)
- *Erigone longipalpis* (Sundevall, 1830)
- *Erigone psychrophila* Thorell, 1872
- *Erigone svenssoni* Holm, 1975
- *Erigone tirolensis* L. Koch, 1872
- *Erigone welchi* Jackson, 1911
- *Erigonella hiemalis* (Blackwall, 1841)
- *Erigonella ignobilis* (O. P.-Cambridge, 1871)
- *Erigonoplus globipes* (L. Koch, 1872)
- *Estrandia grandaeva* (Keyserling, 1886)
- *Evansia merens* O. P.-Cambridge, 1900
- *Flagelliphantes bergstromi* (Schenkel, 1931)
- *Floronia bucculenta* (Clerck, 1757)
- *Glyphesis cottonae* (La Touche, 1945)
- *Gnathonarium dentatum* (Wider, 1834)
- *Gonatium paradoxum* (L. Koch, 1869)
- *Gonatium rubellum* (Blackwall, 1841)
- *Gonatium rubens* (Blackwall, 1833)
- *Gongylidiellum latebricola* (O. P.-Cambridge, 1871)
- *Gongylidiellum murcidum* Simon, 1884
- *Gongylidium rufipes* (Linnaeus, 1758)
- *Halorates reprobus* (O. P.-Cambridge, 1879)
- *Helophora insignis* (Blackwall, 1841)
- *Hilaira excisa* (O. P.-Cambridge, 1871)
- *Hilaira frigida* (Thorell, 1872)
- *Hilaira hermosa* (Thorell, 1875)
- *Hilaira nubigena* Hull, 1911
- *Hilaira pervicax* Hull, 1908
- *Hilaira tatraica* Kulczynski, 1915
- *Horcotes strandi* (Sytshevskaya, 1935)
- *Hybauchenidium ferrumequinum* (Grube, 1861)
- *Hylyphantes graminicola* (Sundevall, 1830)
- *Hypomma bituberculatum* (Wider, 1834)
- *Hypomma cornutum* (Blackwall, 1833)
- *Hypomma fulvum* (Bösenberg, 1902)
- *Hypselistes jacksoni* (O. P.-Cambridge, 1902)
- *Hypselistes paludicola* Tullgren, 1955
- *Improphantes complicatus* (Emerton, 1882)
- *Improphantes decolor* (Westring, 1861)
- *Improphantes holmi* (Kronstedt, 1975)
- *Incestophantes kochiellus* (Strand, 1900)
- *Islandiana falsifica* (Keyserling, 1886)
- *Jacksonella falconeri* (Jackson, 1908)
- *Kaestneria dorsalis* (Wider, 1834)
- *Kaestneria pullata* (O. P.-Cambridge, 1863)
- *Labulla thoracica* (Wider, 1834)
- *Lasiargus hirsutus* (Menge, 1869)
- *Lepthyphantes abiskoensis* Holm, 1945
- *Lepthyphantes antroniensis* Schenkel, 1933
- *Lepthyphantes cornutus* Schenkel, 1927
- *Lepthyphantes crucifer* (Menge, 1866)
- *Lepthyphantes ericaeus* (Blackwall, 1853)
- *Lepthyphantes expunctus* (O. P.-Cambridge, 1875)
- *Lepthyphantes insignis* O. P.-Cambridge, 1913
- *Lepthyphantes keyserlingi* (Ausserer, 1867)
- *Lepthyphantes leprosus* (Ohlert, 1865)
- *Lepthyphantes minutus* (Blackwall, 1833)
- *Lepthyphantes obscurus* (Blackwall, 1841)
- *Lepthyphantes pallidus* (O. P.-Cambridge, 1871)
- *Lepthyphantes punctulatus* Holm, 1939
- *Mughiphantes suffusus* (Strand, 1901)
- *Mughiphantes whympersi* (O. P.-Cambridge, 1894)
- *Leptorhoptrum robustum* (Westring, 1851)
- *Leptothrix hardyi* (Blackwall, 1850)
- *Lessertia denticelis* (Simon, 1884)
- *Linyphia hortensis* Sundevall, 1830
- *Linyphia tenuipalpis* Simon, 1884
- *Linyphia triangularis* (Clerck, 1757)
- *Lophomma punctatum* (Blackwall, 1841)
- *Macrargus boreus* Holm, 1968
- *Macrargus carpenteri* (O. P.-Cambridge, 1894)
- *Macrargus multesimus* (O. P.-Cambridge, 1875)
- *Macrargus rufus* (Wider, 1834)
- *Maro lehtineni* Saaristo, 1971
- *Maro lepidus* Casemir, 1961
- *Maro minutus* O. P.-Cambridge, 1906
- *Maro sublestus* Falconer, 1915

- *Maso sundevalli* (Westring, 1851)
- *Mecynargus borealis* (Jackson, 1930)
- *Mecynargus monticola* (Holm, 1943)
- *Mecynargus morulus* (O. P.-Cambridge, 1873)
- *Mecynargus paetulus* (O. P.-Cambridge 1875)
- *Mecynargus sphagnicola* (Holm, 1939)
- *Megalephyphantes nebulosus* (Sundevall, 1830)
- *Meioneta affinis* (Kulczynski, 1898)
- *Meioneta gulosa* (L. Koch, 1869)
- *Meioneta innotabilis* (O. P.-Cambridge, 1863)
- *Meioneta mollis* (O. P.-Cambridge, 1871)
- *Meioneta mossica* Schikora, 1993
- *Meioneta nigripes* (Simon, 1884)
- *Meioneta rurestris* (C. L. Koch, 1836)
- *Meioneta saxatilis* (Blackwall, 1844)
- *Meioneta similis* (Kulczynski, 1926)
- *Metopobactrus prominulus* (O. P.-Cambridge, 1872)
- *Micrargus apertus* (O. P.-Cambridge, 1871)
- *Micrargus herbigradus* (Blackwall, 1854)
- *Micrargus subaequalis* (Westring, 1851)
- *Microctenonyx subitaneus* (O. P.-Cambridge, 1875)
- *Microlinyphia impigra* (O. P.-Cambridge, 1871)
- *Microlinyphia pusilla* (Sundevall, 1830)
- *Microneta viaria* (Blackwall, 1841)
- *Minicia marginella* (Wider, 1834)
- *Minyriolus pusillus* (Wider, 1834)
- *Mioxena blanda* (Simon, 1884)
- *Moebelia penicillata* (Westring, 1851)
- *Monocephalus castaneipes* (Simon, 1884)
- *Nerienne clathrata* (Sundevall, 1830)
- *Nerienne emphana* (Walckenaer, 1841)
- *Nerienne montana* (Clerck, 1757)
- *Nerienne peltata* (Wider, 1834)
- *Nerienne radiata* (Walckenaer, 1841)
- *Notioscopus sarcinatus* (O. P.-Cambridge, 1872)
- *Oedothorax agrestis* (Blackwall, 1853)
- *Oedothorax apicatus* (Blackwall, 1850)
- *Oedothorax fuscus* (Blackwall, 1834)
- *Oedothorax gibbosus* (Blackwall, 1841)
- *Oedothorax retusus* (Westring, 1851)
- *Oreonetides vaginatus* (Thorell, 1872)
- *Oryphantes angulatus* (O. P.-Cambridge, 1881)
- *Ostearius melanopygius* (O. P.-Cambridge, 1879)
- *Panamomops mengei* Simon, 1926
- *Parapelecopsis nemoralis* (Blackwall, 1841)
- *Pelecopsis elongata* (Wider, 1834)
- *Pelecopsis mengei* (Simon, 1884)
- *Pelecopsis parallela* (Wider, 1834)
- *Pelecopsis radicularis* (L. Koch, 1872)
- *Peponocranium ludicrum* (O.P.-Cambridge, 1861)
- *Pityohyphantes phrygianus* (C. L. Koch, 1836)
- *Pocadicnemis juncea* Locket et Millidge, 1953
- *Pocadicnemis pumila* (Blackwall, 1841)
- *Poeciloneta variegata* (Blackwall, 1841)
- *Porrhomma campbelli* F. O. P.-Cambridge, 1894
- *Porrhomma convexum* (Westring, 1851)
- *Porrhomma egeria* Simon, 1884
- *Porrhomma montanum* Jackson, 1913
- *Porrhomma lativelum* Tretzel, 1956
- *Porrhomma microphthalmum* (O. P.-Cambridge, 1871)
- *Porrhomma pallidum* Jackson, 1913
- *Porrhomma pygmaeum* (Blackwall, 1834)
- *Saaristoa abnormis* (Blackwall, 1841)
- *Saloca diceros* (O. P.-Cambridge, 1871)
- *Satilatlas britteni* (Jackson, 1913)
- *Savignia frontata* Blackwall, 1833
- *Savignia producta* Holm, 1977
- *Scandichrestus tenuis* (Holm, 1943)
- *Scotinotylus alpigenuus* (L. Koch, 1869)
- *Scotinotylus clavatus* (Schenkel, 1927)
- *Scotinotylus evansi* (O. P.-Cambridge, 1894)
- *Semljicola alticola* (Holm, 1950)
- *Semljicola angulatus* (Holm, 1963)
- *Semljicola barbiger* (L. Koch, 1879)
- *Semljicola faustus* (O. P.-Cambridge, 1900)
- *Semljicola lapponicus* (Holm, 1939)
- *Semljicola latus* (Holm, 1939)

- *Silometopus acutus* Holm, 1977
- *Silometopus elegans* (O. P.-Cambridge, 1872)
- *Silometopus incurvatus* (O. P.-Cambridge, 1873)
- *Silometopus reussi* (Thorell, 1871)
- *Sintula corniger* (Blackwall, 1856)
- *Sisicus apertus* (Holm, 1939)
- *Stemonyphantes lineatus* (Linnaeus, 1758)
- *Styloctetor stativus* (Simon, 1881)
- *Tallusia experta* (O. P.-Cambridge, 1871)
- *Tapinocyba biscissa* (O. P.-Cambridge, 1872)
- *Tapinocyba insecta* (L. Koch, 1869)
- *Tapinocyba pallens* (O. P.-Cambridge, 1872)
- *Tapinocyba praecox* (O. P.-Cambridge, 1873)
- *Tapinocyboides pygmaeus* (Menge, 1869)
- *Tapinopa longidens* (Wider, 1834)
- *Taranucnus setosus* (O. P.-Cambridge, 1863)
- *Tenuiphantes alacris* (Blackwall, 1853)
- *Tenuiphantes cristatus* (Menge, 1866)
- *Tenuiphantes flavipes* (Blackwall, 1854)
- *Tenuiphantes mengei* (Kulczynski, 1887)
- *Tenuiphantes nigriventris* (L. Koch, 1879)
- *Tenuiphantes tenebricola* (Wider, 1834)
- *Tenuiphantes tenuis* (Blackwall, 1852)
- *Tenuiphantes zimmermanni* (Bertkau, 1890)
- *Thyreosthenius biovatus* (O. P.-Cambridge, 1875)
- *Thyreosthenius parasiticus* (Westring, 1851)
- *Tibioploides arcuatus* (Tullgren, 1955)
- *Tibioplus diversus* (L. Koch, 1879)
- *Tiso aestivus* (L. Koch, 1872)
- *Tiso vagans* (Blackwall, 1834)
- *Tmeticus affinis* (Blackwall, 1855)
- *Trematocephalus cristatus* (Wider, 1834)
- *Trichoncus hackmani* Millidge, 1956
- *Trichoncus saxicola* (O. P.-Cambridge, 1861)
- *Trichopterna cito* (O. P.-Cambridge, 1872)
- *Trichopterna thorelli* (Westring, 1861)
- *Troxochrota scabra* Kulczynski, 1894
- *Troxochrus nasutus* Schenkel, 1925
- *Troxochrus scabriculus* (Westring, 1851)
- *Typhochrestus digitatus* (O. P.-Cambridge, 1872)
- *Wabasso quaestio* (Chamberlin, 1948)
- *Walckenaeria acuminata* Blackwall, 1833
- *Walckenaeria alticeps* (Denis, 1952)
- *Walckenaeria antica* (Wider, 1834)
- *Walckenaeria atrotibialis* (O. P.-Cambridge, 1878)
- *Walckenaeria capito* (Westring, 1861)
- *Walckenaeria clavicornis* (Emerton, 1882)
- *Walckenaeria cucullata* (C. L. Koch, 1836)
- *Walckenaeria cuspidata* Blackwall, 1833
- *Walckenaeria dysderoides* (Wider, 1834)
- *Walckenaeria furcillata* (Menge, 1869)
- *Walckenaeria incisa* (O. P.-Cambridge, 1871)
- *Walckenaeria karpinskii* (O. P.-Cambridge, 1873)
- *Walckenaeria kochi* (O. P.-Cambridge, 1872)
- *Walckenaeria mitrata* (Menge, 1868)
- *Walckenaeria monoceros* (Wider, 1834)
- *Walckenaeria nodosa* O. P.-Cambridge, 1873
- *Walckenaeria nudipalpis* (Westring, 1851)
- *Walckenaeria obtusa* Blackwall, 1836
- *Walckenaeria picetorum* (Palmgren, 1976)
- *Walckenaeria unicornis* O. P.-Cambridge, 1861
- *Walckenaeria vigilax* (Blackwall, 1853)
- *Zornella cultrigera* (L. Koch, 1879)

TETRAGNATHIDAE

- *Meta menardi* (Latreille, 1804)
- *Metellina mengei* (Blackwall, 1869)
- *Metellina merianae* (Scopoli, 1763)
- *Metellina segmentata* (Clerck, 1757)
- *Pachygnatha clercki* Sundevall, 1823
- *Pachygnatha degeeri* Sundevall, 1830
- *Pachygnatha listeri* Sundevall, 1830
- *Tetragnatha dearmata* Thorell, 1873
- *Tetragnatha extensa* (Linnaeus, 1758)
- *Tetragnatha montana* Simon, 1874
- *Tetragnatha nigrita* Lendl, 1886
- *Tetragnatha obtusa* C. L. Koch, 1837



Tetragnatha extensa
female



Meta menardi female



Meta menardi
egg sac

- *Tetragnatha pinicola* L. Koch, 1870
- *Tetragnatha striata* L. Koch, 1862

ARANEIDAE

- *Aculepeira ceropegia* (Walckenaer, 1802)
- *Aculepeira lapponica* (Holm, 1945)
- *Agalenatea redii* (Scopoli, 1763)
- *Araneus alsine* (Walckenaer, 1802)
- *Araneus angulatus* Clerck, 1757
- *Araneus diadematus* Clerck, 1757
- *Araneus marmoreus* Clerck, 1757
- *Araneus nordmanni* (Thorell, 1870)
- *Araneus quadratus* Clerck, 1757
- *Araneus saevus* (L. Koch, 1872)
- *Araneus sturmi* (Hahn, 1831)
- *Araneus triguttatus* (Fabricius, 1775)
- *Araniella alpica* (L. Koch, 1869)
- *Araniella cucurbitina* (Clerck, 1757)
- *Araniella displicata* (Hentz, 1847)
- *Araniella inconspicua* (Simon, 1874)
- *Araniella opisthographa* (Kulczynski, 1905)
- *Araniella proxima* (Kulczynski, 1895)
- *Argiope bruennichi* (Scopoli, 1772)
- *Cercidia prominens* (Westring, 1851)
- *Cyclosa conica* (Pallas, 1772)
- *Gibbaranea bituberculata* (Walckenaer, 1802)
- *Gibbaranea gibbosa* (Walckenaer, 1802)
- *Gibbaranea omoeda* (Thorell, 1870)
- *Hypsosinga albovittata* (Westring, 1851)
- *Hypsosinga heri* (Hahn, 1831)
- *Hypsosinga pygmaea* (Sundevall, 1831)
- *Hypsosinga sanguinea* (C. L. Koch, 1844)
- *Larinioides cornutus* (Clerck, 1757)
- *Larinioides patagiatus* (Clerck, 1757)
- *Larinioides scolopetarius* (Clerck, 1757)
- *Mangora acalypha* (Walckenaer, 1802)
- *Neoscona adianta* (Walckenaer, 1802)
- *Nuctenea silvicultrix* (C. L. Koch, 1835)
- *Nuctenea umbratica* (Clerck, 1757)
- *Singa hamata* (Clerck, 1757)
- *Singa nitidula* (C. L. Koch, 1845)
- *Zilla diodia* (Walckenaer, 1802)
- *Zygiella atrica* (C. L. Koch, 1845)
- *Zygiella stroemi* (Thorell, 1870)
- *Zygiella x-notata* (Clerck, 1757)

LYCOSIDAE

- *Acantholycosa lignaria* (Clerck, 1757)
- *Acantholycosa norvegica* (Thorell, 1872)
- *Alopecosa aculeata* (Clerck, 1757)
- *Alopecosa barbipes* (Sundevall, 1833)
- *Alopecosa cuneata* (Clerck, 1757)
- *Alopecosa cursor* (Hahn, 1831)
- *Alopecosa fabrilis* (Clerck, 1757)
- *Alopecosa inquilina* (Clerck, 1757)
- *Alopecosa pinetorum* (Thorell, 1856)
- *Alopecosa pulverulenta* (Clerck, 1757)
- *Alopecosa schmidti* (Hahn, 1834)
- *Alopecosa taeniata* (C. L. Koch, 1835)
- *Alopecosa trabalis* (Clerck, 1757)
- *Arctosa alpigena* (Doleschall, 1852)
- *Arctosa cinerea* (Fabricius, 1777)
- *Arctosa figurata* (Simon, 1876)
- *Arctosa leopardus* (Sundevall, 1833)
- *Arctosa lutetiana* (Simon, 1876)
- *Arctosa perita* (Latreille, 1799)
- *Aulonia albimana* (Walckenaer, 1805)
- *Hygrolycosa rubrofasciata* (Ohlert, 1865)
- *Pardosa agrestis* (Westring, 1861)
- *Pardosa agricola* (Thorell, 1856)
- *Pardosa alacris* (C. L. Koch, 1833)
- *Pardosa amentata* (Clerck, 1757)
- *Pardosa atrata* (Thorell, 1873)
- *Pardosa bifasciata* (C. L. Koch, 1834)
- *Pardosa eiseni* (Thorell, 1875)
- *Pardosa fulvipes* (Collett, 1875)
- *Pardosa hyperborea* (Thorell, 1872)
- *Pardosa lapponica* (Thorell, 1872)

*Araneus diadematus* female*Araneus quadratus* female*Acantholycosa lignaria* female with egg sac*Pardosa lugubris* female with egg sac*Pardosa fulvipes* male

- *Pardosa lasciva* L. Koch, 1879
- *Pardosa lugubris* (Walckenaer, 1802)
- *Pardosa monticola* (Clerck, 1757)
- *Pardosa nigriceps* (Thorell, 1856)
- *Pardosa paludicola* (Clerck, 1757)
- *Pardosa palustris* (Linnaeus, 1758)
- *Pardosa plumipes* (Thorell, 1875)
- *Pardosa prativaga* (L. Koch, 1870)
- *Pardosa pullata* (Clerck, 1757)
- *Pardosa riparia* (C. L. Koch, 1833)
- *Pardosa saltans* Töpfer-Hofmann, 2000
- *Pardosa septentrionalis* (Westring, 1861)
- *Pardosa sphagnicola* (Dahl, 1908)
- *Pardosa trailli* (O. P.-Cambridge, 1873)
- *Pirata hygrophilus* Thorell, 1872
- *Pirata insularis* Emerton, 1885
- *Pirata piraticus* (Clerck, 1757)
- *Pirata piscatorius* (Clerck, 1757)
- *Pirata tenuitarsis* Simon, 1876
- *Pirata uliginosus* (Thorell, 1856)
- *Trochosa ruricola* (De Geer, 1778)
- *Trochosa spinipalpis* (F. O. P.-Cambridge, 1895)
- *Trochosa terricola* Thorell, 1856
- *Xerolycosa miniata* (C. L. Koch, 1834)
- *Xerolycosa nemoralis* (Westring, 1861)

PISAURIDAE

- *Dolomedes fimbriatus* (Clerck, 1757)
- *Dolomedes plantarius* (Clerck, 1757)
- *Pisaura mirabilis* (Clerck, 1757)



Dolomedes fimbriatus female



Pisaura mirabilis female

OXYOPIDAE

- *Oxyopes ramosus* (Martini et Goeze, 1778)

AGELENIDAE

- *Agelena labyrinthica* (Clerck, 1757)
- *Tegenaria atrica* C. L. Koch, 1843
- *Tegenaria domestica* (Clerck, 1757)
- *Tegenaria ferruginea* (Panzer, 1804)
- *Textrix denticulata* (Olivier, 1789)



Tegenaria atrica female

CYBAEIDAE

- *Argyroneta aquatica* (Clerck)

HAHNIIDAE

- *Antistea elegans* (Blackwall, 1841)
- *Cryphoeca silvicola* (C. L. Koch, 1834)
- *Hahnia helveola* Simon, 1875
- *Hahnia montana* (Blackwall, 1841)
- *Hahnia nava* (Blackwall, 1841)
- *Hahnia ononidum* Simon, 1875
- *Hahnia pusilla* C. L. Koch 1841

DICTYNYIDAE

- *Archaeodictyna consecuta* (O. P.-Cambridge, 1872)
- *Arctella lapponica* Holm, 1945
- *Argenna patula* (Simon, 1874)
- *Argenna subnigra* (O. P.-Cambridge, 1861)
- *Brommella falcigera* (Balogh, 1935)
- *Cicurina cicur* (Fabricius, 1793)
- *Dictyna alaskae* Chamberlin et Ivie, 1947
- *Dictyna arundinacea* (Linnaeus, 1758)
- *Dictyna latens* (Fabricius, 1775)
- *Dictyna major* Menge, 1869
- *Dictyna pusilla* Thorell, 1856
- *Dictyna 'schmidti'* sensu Lehtinen, 1967
- *Dictyna uncinata* Thorell, 1856
- *Emblyna annulipes* (Blackwall, 1846)
- *Hackmania prominula* (Tullgren, 1948)
- *Lathys humilis* (Blackwall, 1855)

- *Lathys nielseni* (Schenkel, 1932)
- *Mastigusa arietina* (Thorell, 1871)

AMAUROBIDAE

- *Amaurobius fenestralis* (Ström, 1768)
- *Amaurobius ferox* (Walckenaer, 1830)
- *Amaurobius similis* (Blackwall, 1861)
- *Arctobius agelenoides* (Emerton, 1919)
- *Coelotes atropos* (Walckenaer, 1830)

*Amaurobius fenestralis* female**TITANOECIDAE**

- *Titanoeca psammophila* Wunderlich, 1993
- *Titanoeca nivalis* Simon, 1874

MITURGIDAE

- *Cheiracanthium campestre* Lohmander, 1944
- *Cheiracanthium elegans* Thorell, 1875
- *Cheiracanthium erraticum* (Walckenaer, 1802)
- *Cheiracanthium oncognatum* Thorell, 1871
- *Cheiracanthium pennyi* O. P.-Cambridge, 1873
- *Cheiracanthium punctorium* (Villers, 1789)
- *Cheiracanthium virescens* (Sundevall, 1833)

ANYPHAENIDAE

- *Anyphaena accentuata* (Walckenaer, 1802)

LIOCRANIDAE

- *Agraecina striata* (Kulczynski, 1882)
- *Agroeca brunnea* (Blackwall, 1833)
- *Agroeca cuprea* Menge, 1873
- *Agroeca dentigera* Kulczynski, 1913
- *Agroeca lusatica* (L. Koch, 1875)
- *Agroeca proxima* (O. P.-Cambridge, 1871)
- *Apostenus fuscus* Westring, 1851
- *Liocranum rupicola* (Walckenaer, 1830)
- *Phrurolithus festivus* (C. L. Koch, 1835)
- *Phrurolithus minimus* C. L. Koch, 1839
- *Scotina celans* (Blackwall, 1841)
- *Scotina gracilipes* (Blackwall, 1859)
- *Scotina palliardi* (L. Koch, 1881)

CLUBIONIDAE

- *Clubiona brevipes* Blackwall, 1841
- *Clubiona caeruleascens* L. Koch, 1867
- *Clubiona comta* C. L. Koch, 1839
- *Clubiona corticalis* (Walckenaer, 1802)
- *Clubiona diversa* O. P.-Cambridge, 1862
- *Clubiona frisia* Wunderlich et Schuett, 1995
- *Clubiona frutetorum* L. Koch, 1866
- *Clubiona genevensis* L. Koch, 1866
- *Clubiona germanica* Thorell, 1870
- *Clubiona kulczynskii* Lessert, 1905
- *Clubiona lutescens* Westring, 1851
- *Clubiona neglecta* O. P.-Cambridge, 1862
- *Clubiona norvegica* Strand, 1900
- *Clubiona pallidula* (Clerck, 1757)
- *Clubiona phragmitis* C. L. Koch, 1843
- *Clubiona reclusa* O. P.-Cambridge, 1863
- *Clubiona stagnatilis* Kulczynski
- *Clubiona subsultans* Thorell, 1875
- *Clubiona subtilis* L. Koch, 1866
- *Clubiona terrestris* Westring, 1851
- *Clubiona trivialis* C. L. Koch, 1843

*Clubiona phragmitis* male**GNAPHOSIDAE**

- *Callilepis nocturna* (Linnaeus, 1758)
- *Drassodes cupreus* (Blackwall, 1834)
- *Drassodes pubescens* (Thorell, 1856)
- *Drassodes villosus* (Thorell, 1856)
- *Drassyllus lutianus* (L. Koch, 1866)
- *Drassyllus praeficus* (L. Koch, 1866)
- *Drassyllus pumilus* (C. L. Koch, 1839)
- *Drassyllus pusillus* (C. L. Koch, 1833)
- *Echemus angustifrons* (Westring, 1861)
- *Gnaphosa bicolor* (Hahn, 1831)
- *Gnaphosa lapponum* (L. Koch, 1866)
- *Gnaphosa leporina* (L. Koch, 1866)

*Callilepis nocturna* female

- *Gnaphosa lucifuga* (Walckenaer, 1802)
- *Gnaphosa microps* Holm, 1939
- *Gnaphosa montana* (L. Koch, 1866)
- *Gnaphosa muscorum* (L. Koch, 1866)
- *Gnaphosa nigerrima* L. Koch, 1877
- *Gnaphosa orites* Chamberlin, 1922
- *Gnaphosa sticta* Kulczynski, 1908
- *Haplodrassus cognatus* (Westring, 1861)
- *Haplodrassus dalmatensis* (L. Koch, 1866)
- *Haplodrassus moderatus* (Kulczynski, 1897)
- *Haplodrassus signifer* (C. L. Koch, 1839)
- *Haplodrassus silvestris* (Blackwall, 1833)
- *Haplodrassus soerenseni* (Strand, 1900)
- *Haplodrassus umbratilis* (L. Koch, 1866)
- *Micaria aenea* Thorell, 1871
- *Micaria alpina* L. Koch, 1872
- *Micaria formicaria* (Sundevall, 1831)
- *Micaria fulgens* (Walckenaer, 1802)
- *Micaria lenzi* Bösenberg, 1899
- *Micaria nivosa* L. Koch, 1866
- *Micaria pulicaria* (Sundevall, 1831)
- *Micaria silesiaca* L. Koch, 1875
- *Micaria subopaca* Westring, 1861
- *Micaria tripunctata* Holm, 1978
- *Phaeoedus braccatus* (L. Koch, 1866)
- *Poecilochroa variana* (C. L. Koch, 1839)
- *Scotophaeus blackwalli* (Thorell, 1871)
- *Scotophaeus quadripunctatus* (Linnaeus, 1758)
- *Scotophaeus scutulatus* (L. Koch, 1866)
- *Trachyzelotes pedestris* (C. L. Koch, 1837)
- *Zelotes clivicola* (L. Koch, 1870)
- *Zelotes electus* (C. L. Koch, 1839)
- *Zelotes 'exiguus'* sensu Holm 1968
- *Zelotes latreillei* (Simon, 1878)
- *Zelotes longipes* (L. Koch, 1866)
- *Zelotes petrensis* (C. L. Koch, 1839)
- *Zelotes puritanus* Chamberlin, 1922
- *Zelotes subterraneus* (C. L. Koch, 1833)

ZORIDAE

- *Zora armillata* Simon, 1878
- *Zora nemoralis* (Blackwall, 1861)
- *Zora nemoralis palmgreni* Holm, 1945
- *Zora parallela* Simon, 1878
- *Zora silvestris* Kulczynski, 1897
- *Zora spinimana* (Sundevall, 1833)

SPARASSIDAE

- *Micrommata virescens* (Clerck, 1757)

PHILODROMIDAE

- *Philodromus aureolus* (Clerck, 1757)
- *Philodromus cespitum* (Walckenaer, 1802)
- *Philodromus collinus* C. L. Koch, 1835
- *Philodromus dispar* Walckenaer, 1826
- *Philodromus emarginatus* (Schränk, 1803)
- *Philodromus fallax* Sundevall, 1833
- *Philodromus fuscomarginatus* (De Geer, 1778)
- *Philodromus histrio* (Latreille, 1819)
- *Philodromus margaritatus* (Clerck, 1757)
- *Philodromus poecilus* (Thorell, 1872)
- *Philodromus praedatus* O. P.-Cambridge, 1871
- *Philodromus rufus* Walckenaer, 1826
- *Thanatus arcticus* Thorell, 1872
- *Thanatus arenarius* L. Koch, 1872
- *Thanatus atratus* Simon, 1875
- *Thanatus formicinus* (Clerck, 1757)
- *Thanatus striatus* C. L. Koch, 1845
- *Tibellus maritimus* (Menge, 1875)
- *Tibellus oblongus* (Walckenaer, 1802)

*Gnaphosa lucifuga* female*Gnaphosa bicolor* female*Micrommata virescens* female *Micrommata virescens* male*Philodromus collinus* female

THOMISIDAE

- *Coriarachne depressa* (C. L. Koch, 1837)
- *Diaea dorsata* (Fabricius, 1777)
- *Misumena vatia* (Clerck, 1757)
- *Ozyptila arctica* Kulczynski, 1908
- *Ozyptila atomaria* (Panzer, 1801)
- *Ozyptila brevipes* (Hahn, 1826)
- *Ozyptila gertschi* Kurata, 1944
- *Ozyptila nigrita* (Thorell, 1875)
- *Ozyptila praticola* (C. L. Koch, 1837)
- *Ozyptila scabricula* (Westring, 1851)
- *Ozyptila trux* (Blackwall, 1846)
- *Ozyptila westringi* (Thorell, 1873)
- *Pistius truncatus* (Pallas, 1772)
- *Thomisus onustus* Walckenaer, 1806
- *Xysticus albidus* Grese, 1909
- *Xysticus audax* (Schrank, 1803)
- *Xysticus bifasciatus* C. L. Koch, 1837
- *Xysticus chippewa* Gertsch, 1953
- *Xysticus cristatus* (Clerck, 1757)
- *Xysticus erraticus* (Blackwall, 1834)
- *Xysticus kochi* Thorell, 1872
- *Xysticus lanio* C. L. Koch, 1824
- *Xysticus lineatus* (Westring, 1851)
- *Xysticus luctator* L. Koch, 1870
- *Xysticus luctuosus* (Blackwall, 1836)
- *Xysticus obscurus* Collett, 1877
- *Xysticus robustus* (Hahn, 1832)
- *Xysticus sabulosus* (Hahn, 1832)
- *Xysticus ulmi* (Hahn, 1832)

*Xysticus ulmi* female**SALTICIDAE**

- *Aelurillus v-insignitus* (Clerck, 1757)
- *Ballus chalybeius* (Walckenaer, 1802)
- *Dendryphantes hastatus* (Clerck, 1757)
- *Dendryphantes rudis* (Sundevall, 1833)
- *Euophrys frontalis* (Walckenaer, 1802)
- *Evarcha arcuata* (Clerck, 1757)
- *Evarcha falcata* (Clerck, 1757)
- *Evarcha laetabunda* (C. L. Koch, 1848)
- *Heliophanus aeneus* (Hahn, 1831)
- *Heliophanus auratus* C. L. Koch, 1835
- *Heliophanus camtschadalicus* Kulczynski, 1885
- *Heliophanus cupreus* (Walckenaer, 1802)
- *Heliophanus dubius* C. L. Koch, 1835
- *Heliophanus flavipes* (Hahn, 1832)
- *Marpissa muscosa* (Clerck, 1757)
- *Marpissa radiata* (Grube, 1859)
- *Myrmarachne formicaria* (De Geer, 1778)
- *Neon levis* (Simon, 1871)
- *Neon reticulatus* (Blackwall, 1853)
- *Neon robustus* Lohmander, 1945
- *Neon valentulus* Falconer, 1912
- *Pellenes lapponicus* (Sundevall, 1833)
- *Pellenes tripunctatus* (Walckenaer, 1802)
- *Phlegra fasciata* (Hahn, 1826)
- *Pseudeuophrys erratica* (Walckenaer, 1826)
- *Pseudicius encarpatus* (Walckenaer, 1802)
- *Salticus cingulatus* (Panzer, 1797)
- *Salticus scenicus* (Clerck, 1757)
- *Salticus zebraneus* (C. L. Koch, 1837)
- *Sibianor laeae* Logunov, 2001
- *Sitticus caricis* (Westring, 1861)
- *Sitticus distinguendus* (Simon, 1868)
- *Sitticus floricola* (C. L. Koch, 1837)
- *Sitticus inexpectus* Logunov et Kronstedt, 1997
- *Sitticus pubescens* (Fabricius, 1775)
- *Sitticus ranieri* Peckham & Peckham, 1909
- *Sitticus saltator* (O. P.-Cambridge, 1868)
- *Sitticus terebratus* (Clerck, 1757)
- *Sitticus zimmermanni* (Simon, 1877)
- *Synageles hilarulus* (C. L. Koch, 1848)
- *Synageles venator* (Lucas, 1836)
- *Talavera aequipes* (O. P.-Cambridge, 1871)
- *Talavera petrensis* (C. L. Koch, 1837)
- *Talavera thorelli* (Kulczynski, 1891)
- *Talavera westringi* auctt.

*Salticus scenicus* female

[To Torbjörn Kronestedt's Homepage](#)

[/en/spindlar.html](#)

[/en/spindlar.html](#)

Last updated March 12 2001

[Comments](#) about this page.

[Comments](#) about the NRM Website.