

An updated checklist of the Thomisidae (Araneae) of Turkey with zoogeographical remarks

Hakan Demir

Department of Biology, Faculty of Science and Arts, Gazi University, TR-06500 Ankara, Turkey
Corresponding e-mail address: ozyptila@gmail.com

Abstract

This updated checklist of Thomisidae of Turkey is based on published and unpublished records available to the author. At present, 79 species belonging to 14 genera of family Thomisidae have been identified in Turkey.

Keywords: Araneae, Thomisidae, Turkey.

Introduction

Arachnological studies of Turkey began towards the end of the 19th century. They were increased during the 20th century, specially faunistic and taxonomic works. Recently, they continued with an increased speed. In the fauna of Turkey, Thomisidae must be regarded as an insufficiently studied family. The most important papers dealing with the thomisid fauna of the country are those of Kulczyński (1903), Nosek (1905), Roewer (1959), and Simon (1875, 1879, 1884, 1914). Early studies on this family were carried out by Karol (1966a, 1966b, 1966c, 1966d, 1968), who described four new *Xysticus* species from Turkey. However, *X. sislii* Karol, 1966, *X. turcicus* Karol, 1966 and *X. pelini* Karol, 1968 are now synonyms of *X. thessalicus* Simon, 1916 (Wunderlich, 1995), and *X. jezequelii* Karol, 1966 is a synonym of *X. gymnocephalus* Strand, 1915 (Assi, 1986). The previous works were densely concentrated on central Anatolian region, Black Sea region, and the Mediterranean region, i.e. central parts of Turkey.

Turkey has significant ecological properties. It may be the origin country of many spider taxa and it may also be a refugium (an area where conditions have enabled a species or a community of species to survive after extinction in surrounding areas) for living creatures affected by geological and climatic changes more than many countries in the World.

A list of 44 species belonging to 10 genera of family Thomisidae from Turkey was firstly published by Karol (1967). Afterwards, both Turkish and foreign researches made important contributions to the Turkish thomisid fauna. They recorded 55 species belonging to 12 genera from Turkey (Bayram, 2002). According to the checklist of Topçu *et al.* (2005), the Turkish thomisid fauna comprises 56 valid species belonging to 10 genera, most of which were known from a single or just a few localities.

The present checklist is based on all available published and some unpublished records of the distribution of thomisid spiders in Turkey. A total of 79 species from family Thomisidae are recorded in this country. The principal paper is that of Topçu *et al.* (2005), which was based on 41 publications. Now, 21 species are added from recently published 14 papers (i.e. Marusik *et al.*, 2005; Topçu *et al.*, 2006; Logunov, 2006; Logunov & Demir, 2006; Özdemir *et al.*, 2006; Varol *et al.*, 2006; Demir *et al.*, 2006, 2007a, 2007b, 2008, 2008a, 2008b, In press; Bayram *et al.*, 2007).

Material and Methods

The present checklist of the thomisid species of Turkey is mainly based on the data included in “A Checklist of the spiders of Turkey” (Topçu *et al.*, 2005). It is compiled using published records and original data. World distribution of all species follows Platnick (2008). Distribution of species in geographical regions of Turkey is summarized in remarks according to Topçu *et al.* (2005) [MR = Marmara, AR = Aegean, CAR = Central Anatolia, BSR = Black Sea Region, EAR = East Anatolia, SAR = Southeastern Anatolia Region and MER = Mediterranean Regions]. The present zoogeographical characterization is based on the chorotype classification of Anatolian fauna, recently proposed by Vigna Taglianti *et al.* (2000). In this study, as possible as one chorotype description can be identified for each taxon. But this kind of description can not be possible for some taxa, so one or two chorotypes are used for them. The species which are recorded from only one locality in Turkey are characterised by an asterisk (*).

Taxonomy

Family Thomisidae Sundevall, 1833

Genus *Coriarachne* Thorell, 1870

Coriarachne depressa (C.L. Koch, 1837)

Records in Turkey: Aegean and Mediterranean region in Turkey (Bayram, 2002).

Distribution in Turkey: AR and MER. **Chorotype:** Palearctic.

World Distribution: Palearctic.

Genus *Cozyptila* Lehtinen & Marusik, 2005

Cozyptila blackwalli (Simon, 1875)

Records in Turkey: Ankara (Logunov & Demir, 2006).

Distribution in Turkey: CAR *. **Chorotype:** Palearctic.

World Distribution: Palearctic.

Cozyptila guseinovorum Marusik & Kovblyuk, 2005

Records in Turkey: Sakarya, İzmir (Marusik *et al.*, 2005).

Distribution in Turkey: MR and AR. **Chorotype:** Central Asiatic + East European.

World Distribution: Central Asia, Russia, Turkey.

Cozyptila thaleri Marusik & Kovblyuk, 2005

Records in Turkey: Sakarya, Bolu, Konya, Ankara (Logunov & Demir, 2006).

Distribution in Turkey: MR and CAR. **Chorotype:** Eastern European.

World Distribution: Turkey, Ukraine.

Genus *Diaeä* Thorell, 1869

Diaeä livens Simon, 1876

Records in Turkey: Bursa (Bayram *et al.*, 2002); Gaziantep (Özdemir *et al.*, 2006).

Distribution in Turkey: MR, MER and SAR. **Chorotype:** Holarctic.

World Distribution: USA, Central Europe to Azerbaijan.

Genus *Ebrechtella* Dahl, 1907

***Ebrechtella tricuspidata* (Fabricius, 1775)**

Records in Turkey: Çankırı, Ankara, Yozgat (Demir *et al.*, 2007a).

Distribution in Turkey: CAR *. **Chorotype:** Palearctic.

World Distribution: Palearctic.

Genus *Heriaeus* Simon, 1875

***Heriaeus buffoni* (Audouin, 1825)**

Records in Turkey: İstanbul (Kulczyński, 1903).

Distribution in Turkey: MR *. **Chorotype:** North African + South West Asiatic.

World Distribution: North Africa, Israel.

***Heriaeus graminicola* (Doleschall, 1852)**

Records in Turkey: Niğde, Mersin (Topçu *et al.*, 2006).

Distribution in Turkey: MER *. **Chorotype:** European + Central Asiatic.

World Distribution: Europe to Central Asia.

***Heriaeus hirtus* (Latreille, 1819)**

Records in Turkey: İstanbul (Simon, 1875; Pavesi, 1876).

Distribution in Turkey: MR *. **Chorotype:** European.

World Distribution: Europe to Georgia.

***Heriaeus mellotteei* Simon, 1886**

Records in Turkey: Niğde, Mersin, Adana (Topçu *et al.*, 2006); İstanbul (Kulczyński, 1903).

Distribution in Turkey: MR, CAR and MER. **Chorotype:** Palearctic.

World Distribution: Palearctic.

***Heriaeus orientalis* Simon, 1918**

Records in Turkey: İstanbul (Kulczyński, 1903).

Distribution in Turkey: MR *. **Chorotype:** Balkano – Anatolian.

World Distribution: Greece, Turkey, Ukraine.

***Heriaeus pilosus* Nosek, 1905**

Records in Turkey: Sivas, Nevşehir, Kayseri (Nosek, 1905); Erzurum, Tokat (Loerbroks, 1983); Turkey (Reimoser, 1919).

Distribution in Turkey: CAR, EAR and BSR. **Chorotype:** Anatolian endemic.

World Distribution: Turkey.

***Heriaeus setiger* (O.P.-Cambridge, 1872)**

Records in Turkey: Muğla (Dalmas, 1920).

Distribution in Turkey: AR *. **Chorotype:** Palearctic.

World Distribution: Palearctic.

***Heriaeus simoni* Kulczyński, 1903**

Records in Turkey: Hatay, İstanbul (Kulczyński, 1903); İzmir, Bursa, Aydın (Loerbroks, 1983); Turkey (Reimoser, 1919).

Distribution in Turkey: MR and MER. **Chorotype:** Palearctic.

World Distribution: Palearctic.

***Heriaeus spinipalpus* Loerbroks, 1983**

Records in Turkey: Adana, Bitlis, Erzurum, Van (Loerbroks, 1983).

Distribution in Turkey: MER and EAR. **Chorotype:** Mediterranean.

World Distribution: Eastern Mediterranean.

Genus *Misumena* Latreille, 1804

***Misumena vatia* (Clerck, 1757)**

Records in Turkey: İstanbul (Simon, 1875; Pavesi, 1876); Sakarya (Nosek, 1905); Turkey (Reimoser, 1919); Niğde, Mersin (Topçu *et al.*, 2006).

Distribution in Turkey: MER, CAR and MR. **Chorotype:** Holarctic.

World Distribution: Holarctic.

Genus *Monaeses* Thorell, 1869

***Monaeses israeliensis* Levy, 1973**

Records in Turkey: Antalya (Bayram *et al.*, 2007).

Distribution in Turkey: MER *. **Chorotype:** West Asiatic + Central Asiatic.

World Distribution: Greece, Turkey, Israel, Lebanon, Central Asia.

Genus *Ozyptila* Simon, 1864

***Ozyptila ankarensis* Karol, 1966**

Records in Turkey: Ankara (Karol, 1966c).

Distribution in Turkey: CAR *. **Chorotype:** Anatolian endemic.

World Distribution: Turkey.

***Ozyptila atomaria* (Panzer, 1801)**

Records in Turkey: Van (Bayram, 1996a, b), Kırıkkale (Bayram *et al.*, 2005).

Distribution in Turkey: EAR and SAR. **Chorotype:** Palearctic.

World Distribution: Palearctic.

***Ozyptila claveata* (Walckenaer, 1837)**

Records in Turkey: İzmir (Lehtinen, 2002); Gaziantep (Özdemir *et al.*, 2006; Varol *et al.*, 2006); Niğde (Topçu *et al.*, 2006).

Distribution in Turkey: AR and SAR. **Chorotype:** Palearctic.

World Distribution: Palearctic.

***Ozyptila clavidorsa* Roewer, 1959**

Records in Turkey: Gaziantep (Roewer, 1959).

Distribution in Turkey: SAR *. **Chorotype:** Anatolian endemic.

World Distribution: Turkey.

***Ozyptila conostyla* Hippa, Koponen & Oksala, 1986**

Records in Turkey: Yozgat (Hippa *et al.*, 1986).

Distribution in Turkey: CAR *. **Chorotype:** Caucasian.

World Distribution: Turkey to Turkmenistan.

***Ozyptila praticola* (C.L. Koch, 1837)**

Records in Turkey: İstanbul (Roewer, 1959); Van (Bayram, 1996a; Bayram & Varol, 1996); Denizli (Bayram *et al.*, 1998); Kırıkkale (Bayram *et al.*, 2005); Adana (Topçu *et al.*, 2006).

Distribution in Turkey: MR, EAR, AR, CAR and MER. **Chorotype:** Holarctic.

World Distribution: Holarctic.

***Ozyptila rauda* Simon, 1875**

Records in Turkey: Zonguldak (Roewer, 1959).

Distribution in Turkey: BSR (WBR) *. **Chorotype:** Palearctic.

World Distribution: Palearctic.

***Ozyptila sanctuaria* (O.P.-Cambridge, 1871)**

Records in Turkey: Van (Bayram, 1996a); Kırıkkale (Bayram *et al.*, 2005).

Distribution in Turkey: EAR and CAR. **Chorotype:** European.

World Distribution: Europe.

Ozyptila simplex (O.P.-Cambridge, 1862)

Records in Turkey: Turkey (Caporiacco, 1935); Gaziantep (Özdemir *et al.*, 2006); Adana (Topçu *et al.*, 2006).

Distribution in Turkey: EAR and MER. **Chorotype:** Palearctic.

World Distribution: Palearctic.

Ozyptila spirembola Wunderlich, 1995

Records in Turkey: Bolu (Wunderlich, 1995).

Distribution in Turkey: MR *. **Chorotype:** Anatolian endemic.

World Distribution: Turkey.

Ozyptila tricoloripes Strand, 1913

Records in Turkey: Kahramanmaraş, Hatay, (Demir *et al.*, 2008a).

Distribution in Turkey: MER *. **Chorotype:** Turanian.

World Distribution: Israel, Azerbaijan, Turkmenistan.

Genus *Pistius* Simon, 1875

Pistius truncatus (Pallas, 1772)

Records in Turkey: Niğde, Mersin (Topçu *et al.*, 2006); İstanbul (Simon, 1879); Turkey (Karol, 1967).

Distribution in Turkey: CAR, MER and MR. **Chorotype:** Palearctic.

World Distribution: Palearctic.

Genus *Runcinia* Simon, 1875

Runcinia grammica (C.L. Koch, 1837)

Records in Turkey: Sakarya (Nosek, 1905); Mersin, Adana (Topçu *et al.*, 2006); Bahkesir (Karol, 1966d); İstanbul, Hatay (Kulczyński, 1903); Turkey (Karol, 1967; Caporiacco, 1935); Bursa (Kaya & Uğurtaş, 2007).

Distribution in Turkey: MR, MER and AR. **Chorotype:** Palearctic + Ethiopian.

World Distribution: Palearctic, St. Helena, South Africa.

Genus *Synema* Simon, 1864

Synema globosum (Fabricius, 1775)

Records in Turkey: Kocaeli, Sakarya (Nosek, 1905); Hatay, Kilis, Adana, Osmaniye (Demir *et al.*, 2007b); Gaziantep (Özdemir *et al.*, 2006); Niğde, Mersin, Adana (Topçu *et al.*, 2006); Bursa (Karol, 1966d); İstanbul (Simon, 1875; Pavesi, 1876); Hatay, İstanbul (Kulczyński, 1903); Turkey (Strand, 1917; Caporiacco, 1935); Aydın (Giltay, 1932); Osmaniye (Roewer, 1959); Bursa (Kaya & Uğurtaş, 2007).

Distribution in Turkey: MR, MER, SAR, AR and CAR. **Chorotype:** Palearctic.

World Distribution: Palearctic.

Synema plorator (O.P.-Cambridge, 1872)

Records in Turkey: Osmaniye, Kahramanmaraş (Demir *et al.*, 2007b); Bursa (Karol, 1966d).

Distribution in Turkey: MER and MR. **Chorotype:** South West Asiatic + Central Asiatic.

World Distribution: Slovakia to Israel, Central Asia.

Synema utotchkini Marusik & Logunov, 1995

Records in Turkey: Kahramanmaraş (Demir *et al.*, 2007b).

Distribution in Turkey: MER *. **Chorotype:** Caucasian.

World Distribution: Kazakhstan, Kyrgyzstan, Turkey.

Genus *Thomisus* Walckenaer, 1805

***Thomisus citrinellus* Simon, 1875**

Records in Turkey: Black Sea Region (Topçu *et al.*, 2005).

Distribution in Turkey: BSR (MBR) *. **Chorotype:** Afrotropico - Mediterranean.

World Distribution: Mediterranean, Africa, Yemen, Socotra, Seychelles.

***Thomisus onustus* Walckenaer, 1805**

Records in Turkey: Van (Bayram, 1996b; Bayram & Varol, 1996); Denizli (Bayram *et al.*, 1998); Manisa, İzmir, Aydın (Bayram *et al.*, 2000); Sakarya, Sivas, Konya, Kayseri (Nosek, 1905); Kırıkkale (Bayram *et al.*, 2005); Gaziantep (Özdemir *et al.*, 2006); Niğde, Mersin, Adana (Topçu *et al.*, 2006); Ankara (Karol, 1966d); Turkey (Caporiacco, 1935; Ono & Martens, 2005); İstanbul (Simon, 1875; Pavesi, 1876; Kulczyński, 1903; Giltay, 1932); İzmir (Kulczyński, 1903); Hatay, Gaziantep (Roewer, 1959); Bursa (Kaya & Uğurtaş, 2007).

Distribution in Turkey: EAR, AR, MR, SAR, MER and CAR. **Chorotype:** Palearctic.

World Distribution: Palearctic.

***Thomisus zyuzini* Marusik & Logunov, 1990**

Records in Turkey: Osmaniye (Demir *et al.*, 2008a).

Distribution in Turkey: MER *. **Chorotype:** South West Asiatic + Central Asiatic.

World Distribution: Saudi Arabia to Central Asia.

Genus *Tmarus* Simon, 1875

***Tmarus piochardi* (Simon, 1866)**

Records in Turkey: Diyarbakır, Muğla (Bayram *et al.*, 2007).

Distribution in Turkey: SAR and MER. **Chorotype:** Mediterranean.

World Distribution: Mediterranean.

***Tmarus stellio* Simon, 1875**

Records in Turkey: Ankara (Demir *et al.*, 2007a).

Distribution in Turkey: CAR *. **Chorotype:** Palearctic.

World Distribution: Palearctic.

Genus *Xysticus* C.L. Koch, 1835

***Xysticus abditus* Logunov, 2006**

Records in Turkey: Niğde (Logunov, 2006).

Distribution in Turkey: CAR *. **Chorotype:** Balkano – Anatolian.

World Distribution: Bulgaria, Turkey.

***Xysticus acerbus* Thorell, 1872**

Records in Turkey: Turkey (Simon, 1914); İstanbul (Simon, 1875; Pavesi, 1876); Gaziantep (Roewer, 1959).

Distribution in Turkey: MR and SAR. **Chorotype:** European + Central Asiatic.

World Distribution: Europe to Central Asia.

***Xysticus anatolicus* Demir, Aktaş & Topçu, 2008**

Records in Turkey: Kayseri (Demir *et al.*, 2008b).

Distribution in Turkey: CAR *. **Chorotype:** Anatolian endemic.

World Distribution: Turkey.

***Xysticus audax* (Schrank, 1803)**

Records in Turkey: İstanbul, Adana, Osmaniye, Hatay (Roewer, 1959); Turkey (Simon, 1914).

Distribution in Turkey: MR and MER. **Chorotype:** Palearctic.

World Distribution: Palearctic.

Xysticus bacurianensis Mccheidze, 1971

Records in Turkey: Trabzon (Logunov & Demir, 2006).

Distribution in Turkey: BSR *. **Chorotype:** Turanian + East European.

World Distribution: Turkey, Russia, Georgia, Azerbaijan.

Xysticus bifasciatus C.L. Koch, 1837

Records in Turkey: Konya (Bayram & Allahverdi, 1994, 1999); Manisa, İzmir, Aydın (Bayram *et al.*, 2000); Kırıkkale (Bayram *et al.*, 2005); Niğde (Topçu & Demir, 2004; Topçu *et al.*, 2006).

Distribution in Turkey: CAR and AR. **Chorotype:** Palearctic.

World Distribution: Palearctic.

Xysticus bufo (Dufour, 1820)

Records in Turkey: İstanbul (Simon, 1875; Pavesi, 1876); İzmir (Pavesi, 1876).

Distribution in Turkey: MR and AR. **Chorotype:** Mediterranean.

World Distribution: Mediterranean.

Xysticus caperatus Simon, 1875

Records in Turkey: Osmaniye (Demir, *et al.*, In press).

Distribution in Turkey: MER *. **Chorotype:** Mediterranean + East European.

World Distribution: Mediterranean, Russia.

Xysticus cibratus Simon, 1885

Records in Turkey: Turkey (Lehtinen, 2002); Ankara (Karol, 1966d).

Distribution in Turkey: CAR *. **Chorotype:** Mediterranean + Asiatic.

World Distribution: Mediterranean to China, Sudan.

Xysticus cristatus (Clerck, 1757)

Records in Turkey: Van (Bayram, 1994, 1996b, c; Bayram *et al.*, 1999; Bayram & Varol, 1999); Manisa, İzmir, Aydın (Bayram *et al.*, 2000); Denizli (Bayram *et al.*, 1998); Konya (Bayram & Allahverdi, 1999); Kırıkkale (Bayram *et al.*, 2005); Niğde, Mersin (Topçu *et al.*, 2006); Trabzon (Logunov & Demir, 2006).

Distribution in Turkey: EAR, SAR, AR, CAR, MER and BSR. **Chorotype:** Palearctic.

World Distribution: Palearctic.

Xysticus demirsoyi Demir, Topçu & Türkès, 2006

Records in Turkey: Niğde (Demir, *et al.*, 2006).

Distribution in Turkey: CAR *. **Chorotype:** Anatolian endemic.

World Distribution: Turkey.

Xysticus edax (O. P.-Cambridge, 1872)

Records in Turkey: Osmaniye, Kilis, Kahramanmaraş, Hatay (Demir, *et al.*, In press).

Distribution in Turkey: MER *. **Chorotype:** South West Asiatic.

World Distribution: Israel.

Xysticus erraticus (Blackwall, 1834)

Records in Turkey: Van (Bayram, 1996a).

Distribution in Turkey: EAR *. **Chorotype:** West Palearctic.

World Distribution: Europe, Russia.

Xysticus ferrugineus Menge, 1876

Records in Turkey: Niğde (Topçu & Demir, 2004; Topçu *et al.*, 2006).

Distribution in Turkey: CAR *. **Chorotype:** Palearctic.

World Distribution: Palearctic.

Xysticus ferus O.P.-Cambridge, 1876

Records in Turkey: Muğla (Dalmas, 1920).

Distribution in Turkey: MER *. **Chorotype:** East Mediterranean.

World Distribution: Cyprus, Egypt, Israel.

Xysticus gallicus Simon, 1875

Records in Turkey: Turkey (Simon, 1914; Karol, 1967; Ono & Martens, 2005); Kayseri (Nosek, 1905).

Distribution in Turkey: CAR *. **Chorotype:** Palearctic.

World Distribution: Palearctic.

Xysticus graecus C.L. Koch, 1837

Records in Turkey: Ankara (Karol, 1966d), Bursa, İstanbul (Kulczyński, 1903).

Distribution in Turkey: CAR and MR. **Chorotype:** East Mediterranean + East European.

World Distribution: Eastern Mediterranean, Russia.

Xysticus gymnocephalus Strand, 1915

Records in Turkey: Ankara (Karol, 1966a, 1966d).

Distribution in Turkey: CAR *. **Chorotype:** South West Asiatic.

World Distribution: Turkey, Lebanon, Israel.

Xysticus kaznakovi Utochkin, 1968

Records in Turkey: Kahramanmaraş, Osmaniye (Demir, *et al.*, In press).

Distribution in Turkey: MER *. **Chorotype:** Central Asiatic.

World Distribution: Central Asia.

Xysticus kochi Thorell, 1872

Records in Turkey: Sakarya, Kayseri (Nosek, 1905); Turkey (Reimoser, 1919); Konya (Bayram & Allahverdi, 1994, 1999); Denizli (Bayram *et al.*, 1998); Van (Bayram & Varol, 2000); Kırıkkale (Bayram *et al.*, 2005); Niğde, Mersin (Topçu *et al.*, 2006); Ankara, Bursa (Karol, 1966d); İstanbul (Simon, 1875; Pavesi, 1876); Turkey (Caporiacco, 1935); Çankırı, Konya, Yozgat, Isparta, Bolu, Adana, Kayseri, Sinop, Zonguldak, Nevşehir (Logunov & Demir, 2006).

Distribution in Turkey: MR, CAR, AR, SAR, MER and BSR. **Chorotype:** Palearctic.

World Distribution: Europe, Mediterranean to Central Asia.

Xysticus laetus Thorell, 1875

Records in Turkey: Konya (Nosek, 1905); Turkey (Reimoser, 1919); Mersin (Topçu *et al.*, 2006); Nevşehir, Çankırı, Konya, Kırıkkale, Bolu, Yozgat, Niğde, Antalya (Logunov, 2006); Bursa (Kaya & Uğurtas, 2007).

Distribution in Turkey: CAR, MER and MR. **Chorotype:** East Mediterranean + South West Asiatic.

World Distribution: Italy to Central Asia.

Comment: All records of *Xysticus kempeleni* Thorell, 1872 from Turkey belong to *X. laetus* (Logunov, 2006).

Xysticus lalandei (Audouin, 1825)

Records in Turkey: Ankara (Karol, 1966d); Konya, Niğde, Kayseri (Nosek, 1905); Turkey (Strand, 1917); Muğla (Dalmas, 1920).

Distribution in Turkey: CAR and MER. **Chorotype:** North East African - Sindian.

World Distribution: Egypt, Israel.

Xysticus lanio C.L. Koch, 1835

Records in Turkey: İstanbul (Pavesi, 1876); Hatay (Roewer, 1959); Niğde, Mersin (Topçu *et al.*, 2006).

Distribution in Turkey: MR, MER and SAR. **Chorotype:** Palearctic.

World Distribution: Palearctic.

Xysticus lineatus (Westring, 1851)

Records in Turkey: Mersin (Topçu & Demir, 2004; Topçu *et al.*, 2006).

Distribution in Turkey: MER *. **Chorotype:** Palearctic.

World Distribution: Palearctic.

Xysticus luctator L. Koch, 1870

Records in Turkey: Niğde (Nosek, 1905).

Distribution in Turkey: CAR *. **Chorotype:** Palearctic.

World Distribution: Palearctic.

Xysticus luctuosus (Blackwall, 1836)

Records in Turkey: Van (Bayram & Varol, 1999, 2000; Bayram *et al.*, 1999); Konya (Bayram & Allahverdi, 1994); Gaziantep (Özdemir *et al.*, 2006).

Distribution in Turkey: SAR, CAR and EAR. **Chorotype:** Holarctic.

World Distribution: Holarctic.

Xysticus macedonicus Silhavy, 1944

Records in Turkey: Gaziantep (Özdemir *et al.*, 2006).

Distribution in Turkey: SAR *. **Chorotype:** Central European.

World Distribution: Germany, Switzerland, Austria, Macedonia, Turkey.

Xysticus ninnii fusciventris Crome, 1965

Records in Turkey: Sivas (Nosek, 1905); Turkey (Reimoser, 1919); Van (Bayram *et al.*, 1999; Bayram & Varol, 2000); Mersin (Topçu *et al.*, 2006); Sivas, Çankırı, Kayseri, Konya (Logunov & Demir, 2006).

Distribution in Turkey: SAR, MER and CAR. **Chorotype:** Eastern Europe + Central Asiatic.

World Distribution: Eastern Europe to Mongolia.

Xysticus nubilus Simon, 1875

Records in Turkey: İzmir (Simon, 1914).

Distribution in Turkey: AR *. **Chorotype:** Mediterranean.

World Distribution: Mediterranean, Azores, Macronesia.

Xysticus pseudolanio Wunderlich, 1995

Records in Turkey: Bolu, Artvin (Wunderlich, 1995).

Distribution in Turkey: MR and BSR. **Chorotype:** Anatolian endemic.

World Distribution: Turkey.

Xysticus pseudorectilineus (Wunderlich, 1995)

Records in Turkey: Adana (Wunderlich, 1995); Adana, Niğde, Adana, Nevşehir, Osmaniye, Antalya, Kayseri, Mersin, Şanlıurfa, Gaziantep (Demir *et al.*, 2008).

Distribution in Turkey: MER, CAR and EAR. **Chorotype:** Anatolian endemic.

World Distribution: Turkey.

Xysticus rectilineus (O.P.-Cambridge, 1872)

Records in Turkey: Ankara (Karol, 1966d).

Distribution in Turkey: CAR *. **Chorotype:** South West Asiatic.

World Distribution: Syria, Lebanon, Israel.

Xysticus robustus (Hahn, 1832)

Records in Turkey: Van (Bayram, 1996a; Bayram & Varol, 1996; Bayram *et al.*, 1999); Konya (Bayram & Allahverdi, 1994, 1999); Manisa, İzmir, Aydın (Bayram *et al.*, 2000); Kırıkkale (Bayram *et al.*, 2005); Gaziantep (Özdemir *et al.*, 2006); Niğde, Mersin (Topçu *et al.*, 2006).

Distribution in Turkey: SAR, CAR, AR, EAR and MER. **Chorotype:** European + Central Asiatic.

World Distribution: Europe to Central Asia.

Xysticus sabulosus (Hahn, 1832)

Records in Turkey: Van (Bayram *et al.*, 1999); Niğde, Mersin (Topçu *et al.*, 2006); İstanbul (Simon, 1875; Pavesi, 1876).

Distribution in Turkey: SAR, CAR and MR. **Chorotype:** Palearctic.

World Distribution: Palearctic.

Xysticus striatipes L. Koch, 1870

Records in Turkey: Ankara (Karol, 1966d); Mersin (Topçu *et al.*, 2006); Bursa (Kaya & Uğurtaş, 2007).

Distribution in Turkey: CAR, MER and MR. **Chorotype:** Palearctic.

World Distribution: Palearctic.

Xysticus thessalicoides Wunderlich, 1995

Records in Turkey: Antalya, Kastamonu, Bolu, Trabzon (Logunov & Demir, 2006).

Distribution in Turkey: MER, BSR and MR. **Chorotype:** Balkano – Anatolian.

World Distribution: Greece, Crete, Turkey.

Xysticus thessalicus Simon, 1916

Records in Turkey: Ankara (Karol, 1966d); Konya (Karol, 1966b, 1966d, 1968); Manisa (Azarkina & Logunov, 2000); Konya, Mersin, Yozgat (Logunov & Demir, 2006).

Distribution in Turkey: CAR, AR and MER. **Chorotype:** North East Mediterranean.

World Distribution: Balkans, Greece, Turkey, Israel.

Xysticus tristrami (O.P.-Cambridge, 1872)

Records in Turkey: Konya, Niğde, Kayseri (Nosek, 1905); Hatay (Simon, 1884).

Distribution in Turkey: CAR and MER. **Chorotype:** South West Asiatic + Central Asiatic.

World Distribution: Saudi Arabia to Central Asia.

Xysticus ulmi (Hahn, 1831)

Records in Turkey: Konya (Karol, 1966d); Van (Bayram, 1996c; Bayram *et al.*, 1999); Kırıkkale (Bayram *et al.*, 2005); Gaziantep (Özdemir *et al.*, 2006); Mersin (Topçu *et al.*, 2006).

Distribution in Turkey: CAR, SAR, EAR and MER. **Chorotype:** Palearctic.

World Distribution: Palearctic.

Xysticus viduus Kulczyński, 1898

Records in Turkey: Niğde (Topçu & Demir, 2004; Topçu *et al.*, 2006).

Distribution in Turkey: CAR *. **Chorotype:** Palearctic.

World Distribution: Palearctic.

Xysticus xerodermus Strand, 1913

Records in Turkey: Konya, Yozgat (Logunov & Demir, 2006).

Distribution in Turkey: CAR *. **Chorotype:** South West Asiatic.

World Distribution: Turkey, Israel.

Discussion

Turkey is a bridge between Asia and Europe. Its geographical location is very important in Palearctic region. Turkey consists of two general parts as Trace and Anatolia. The European section of Turkey is eastern Thrace. It forms the borders of Turkey with Greece and Bulgaria. The Asian part of the country, Anatolia, consists of a high central plateau with narrow coastal plains. Turkey has a great variability in topography and climate. The fact that Anatolia is surrounded from three sides by sea, its situation in the temperate climatic zone, its geological and geomorphic structure, and topography are main contributing factors affecting diversity of species in terrestrial

ecosystems. This study has increased the number of thomisid spiders in Turkey to 79 species belonging to 14 genera.




Fig. 1. Provinces of Turkey.

Zoogeographically, *Coriarachne depressa*, *Cozyptila blackwalli*, *Ebrechtella tricuspidata*, *Heriaeus mellotteei*, *H. setiger*, *H. simoni*, *Ozyptila atomaria*, *O. claveata*, *O. rauda*, *O. simplex*, *Pistius truncatus*, *Runcinia grammica*, *Synema globosum*, *Thomisus onustus*, *Tmarus stellio*, *Xysticus audax*, *X. bifasciatus*, *X. cristatus*, *X. ferrugineus*, *X. gallicus*, *X. kochi*, *X. lanio*, *X. lineatus*, *X. luctator*, *X. sabulosus*, *X. striatipes*, *X. ulmi* and *X. viduus* have the Palearctic chorotype. *Cozyptila guseinovorum* has the Central Asiatic + East European chorotypes. *Cozyptila thaleri* has the Eastern European chorotype. *Diaealivens*, *Misumena vatia*, *Ozyptila praticola* and *Xysticus luctuosus* have the Holarctic chorotype. *Heriaeus buffoni* has the North African + South West Asiatic chorotypes. *Heriaeus graminicola*, *Xysticus acerbus* and *X. robustus* have the European + Central Asiatic chorotypes. *Heriaeus hirtus* and *Ozyptila sanctuaria* have the European chorotype. *Heriaeus orientalis*, *Xysticus abditus* and *X. thessalicoides* have the Balkano – Anatolian chorotype. *Monaeses israeliensis*, *Synema plorator*, *Thomisus zyuzini* and *Xysticus tristrami* have the South West Asiatic + Central Asiatic chorotypes. *Ozyptila conostyla* and *Synema utotchkini* have the Caucasian chorotype. *Ozyptila tricoloripes* has the Turanian chorotype. *Thomisus citrinellus* has the Afrotropico - Mediterranean chorotype. *Heriaeus spinipalpus*, *Tmarus piocchardi*, *Xysticus bufo* and *X. nubilus* have the Mediterranean chorotype. *Xysticus bacurianensis* has the Turanian + East European chorotypes. *Xysticus cibratus* has the Mediterranean + Asiatic chorotypes. *Xysticus edax*, *X. gymnocephalus*, *X. rectilineus* and *X. xerodermus* has the South West Asiatic chorotype. *Xysticus erraticus* has the West Palearctic chorotype. *Xysticus ferus* has the East Mediterranean chorotype. *Xysticus graecus* has the East Mediterranean + East European chorotypes. *Xysticus caperatus* has the Mediterranean + East European chorotypes. *Xysticus kaznakovi* has the Central Asiatic chorotype. *Xysticus laetus* has the East Mediterranean + South West Asiatic chorotypes. *Xysticus lalandei* has the North East African - Sindian chorotype. *Xysticus macedonicus* has the Central European chorotype. *X. ninnii fusciventris* has the Eastern Europe + Central Asiatic chorotypes. *Xysticus thessalicus* has the North East Mediterranean chorotype. *Heriaeus pilosus*, *Ozyptila ankarensis*, *O. clavidorsa*, *O. spirembola*, *Xysticus anatolicus*, *X. demirsoyi*, *X. pseudolanolio* and *X. pseudorectilineus* have the Anatolian endemic chorotype.

References

- Assi, F. 1986. Note faunistique sur les Thomisidae et les Philodromidae du Liban (Araneae). *Revue Arachnologique*, 7(1): 41-46.
- Azarkina, G.N. & Logunov, D.V. 2000. Separation and distribution of *Xysticus cristatus* (Clerck, 1758) and *X. audax* (Schrank, 1803) in eastern Eurasia, with description of a new species from the mountains of Central Asia (Aranei: Thomisidae). *Arthropoda Selecta*, 9(2): 133-150.
- Bayram, A. 1994. Tarla kenarlarında yer alan ot kümelerinin Arthropod faunası. *Yüzüncü Yıl Üniversitesi Ziraat Fakültesi Dergisi*, 4: 139-149.
- Bayram, A. 1996a. Spider fauna (Ordo: Araneae) of Çarpanak island (Lake Van). *Yüzüncü Yıl University Journal of Education*, 1(2): 59-68.
- Bayram, A. 1996b. Spiders from Akdamar island (Lake Van): Faunistic notes, habitat descriptions and sampling methods. *Gazi Üniversitesi Fen-Edebiyat Fakültesi Fen Bilimleri Dergisi*, 6: 1-10.
- Bayram, A. 1996c. Van'daki örümcek populasyonlarında mevsime bağlı olarak ortaya çıkış. *Gazi Üniversitesi Fen-Edebiyat Fakültesi Fen Bilimleri Dergisi*, 6: 11-23.
- Bayram, A. 2002. Türkiye Örümceklerinin (Araneae) Tür Listesi ve Yayınlıları, 23. (pp.638-657) Bölüm. *Genel Zoocografya ve Türkiye Zoocoğrafyası "Hayvan Coğrafyası"*, In: Ali Demirsoy (Ed.), 5. Baskı, Meteksan Yayınları, 2002, 1007 sh.
- Bayram, A. & Allahverdi, H. 1994. Tarla, Orman ve Step yer örümceklerinin (Ordo: Araneae) ekolojik açıdan incelenmesi. *Yüzüncü Yıl Üniversitesi, Fen Bilimleri Dergisi*, 5(5): 27-39.
- Bayram, A. & Allahverdi, H. 1999. Tarımsal ekosistemlerde örümceklerin habitat tercihleri üzerine. *Centre for Entomological Studies Miscellaneous Papers*, 58: 1-7.
- Bayram, A., Allahverdi, H., Varol, M.İ.. & Pekdemir, H. 1998. Denizli yöresi tütün tarlalarının örümcek faunası. *Centre for Entomological Studies Miscellaneous Papers*, 57: 1-6.
- Bayram, A., Danışman, T., Bolu, H. & Özgen, İ. 2007. Two records new for the Turkish araneofauna: *Tmarus piochardi* (Simon, 1866) and *Monaeses israeliensis* Levy, 1973 (Araneae: Thomisidae). *Munis Entomology & Zoology*, 2(1): 129-136.
- Bayram, A., Danışman, T., Yeşilyurt, F., Çorak, I. & Ünal, M. 2005. Kırıkkale İlinin Araneo-Faunası Üzerine (Arthropoda: Arachnida). *Ekoloji Çevre Dergisi*, 14(56): 1-8.
- Bayram, A., Özdağ, S. & Kaya, R. 2002. New spider [Araneae] records for Turkey: *Hyptiotes paradoxus* (Koch C.L., 1834) [Uloboridae], *Diaeа pictilis* (Banks, 1896) [Thomisidae], *Alopecosa fabrilis* (Clerck, 1757) [Lycosidae] and *Evarcha arcuata* (Clerck, 1757) [Salticidae]. *Israel Journal of Zoology*, 48: 250-251.
- Bayram, A. & Varol, M.İ. 1996. Spider fauna (Ordo: Araneae) of Van Castle and its environment. *Yüzüncü Yıl University Journal of Faculty of Education*, 1(2): 183-193.
- Bayram, A. & Varol, M.İ. 1999. A Study on Spiders (Araneae) in Grass Tussocks in Van Vicinity. *Turkish Journal of Zoology*, 23, 1: 173-175.
- Bayram, A. & Varol, I. 2000. Spiders active on snow in eastern Turkey. *Zoology in the Middle East*, 21: 133-137.
- Bayram, A., Varol, M.İ., Allahverdi, H., Polat, M. & Bulut, M. 1999. Van'da bir korunga tarlasının örümcek faunası. *Ekoloji Çevre Dergisi*, 8(33): 1-4.
- Bayram, A., Varol, M.İ. & Tozan, İ.H. 2000. The Spider (Araneae) fauna of the cotton fields located in the western part of Turkey. *Serket*, 6(4): 105-114.
- Caporiacco, L. di. 1935. Escursione del Prof. Nello Beccari in Anatolia. Aracnidi. *Monitore zoologico italiano*, 46(9): 283-289.

- Dalmas, R. de. 1920. Liste d'araignées de Boudron en Asie Mineure suivie d'une étude des espèces méditerranéennes du genre *Habrocestum*. *Annali del Museo Civico di Storia Naturale di Genova*, 50: 57-69.
- Demir, H., Aktaş, M. & Seyyar, O. 2008. The female of *Xysticus pseudorectilineus* (Wunderlich, 1995) (Araneae: Thomisidae) from Turkey. *Zootaxa*, 1674: 65-68.
- Demir, H., Aktaş, M., Topçu, A. & Seyyar, O. 2007a. A contribution to the crab spider fauna of Turkey (Araneae: Thomisidae). *Serket*, 10(3): 86-90.
- Demir, H., Aktaş, M. & Topçu, A. 2007b. A review of the genus *Synema* Simon, 1864 (Araneae: Thomisidae) in Turkey with a new record, *Synema utotchkini* Marusik & Logunov, 1995. *Serket*, 10(4): 120-122.
- Demir, H., Aktaş, M. & Topçu, A. 2008a. Two thomisid species new to the Turkish spider fauna: *Ozyptila tricoloripes* Strand, 1913 and *Thomisus zyuzini* Marusik & Logunov, 1990 (Araneae: Thomisidae). *Turkish Journal of Arachnology*, 1(1): 44-48.
- Demir, H., Aktaş, M. & Topçu, A. 2008b. *Xysticus anatolicus* n.sp. (Araneae: Thomisidae), a new species from Turkey. *Entomological News*, 119(3): 287-290.
- Demir, H., Aktaş, M. & Topçu, A. (In press). New records of little-known species of *Xysticus* C. L. Koch, 1835 (Araneae: Thomisidae) in Turkey. *Zoology in the Middle East*.
- Demir, H., Topçu, A. & Türkeş, T. 2006. A new species of the genus *Xysticus* C. L. Koch from Turkey (Araneae: Thomisidae). *Zootaxa*, 1364: 45-49.
- Giltay, L. 1932. Arachnides recueillis par M. d'Orchymont au cours de ses voyages aux Balkans et en Asie Mineure en 1929, 1930 et 1931. *Bulletin du Museum royale d'histoire naturelle de Belgique*, 8(22): 1-40.
- Hippa, H., Koponen, S. & Oksala, I. 1986. Revision and classification of the Holarctic species of the *Ozyptila rauda* group (Araneae, Thomisidae). *Annales Zoologici Fennici*, 23: 321-328.
- Karol, S. 1966a. Description d'une araignée nouvelle en Turquie (Araneae, Thomisidae). *Communications de la Faculté des Sciences de l'Université d'Ankara*, 11(C): 1-5.
- Karol, S. 1966b. Sur une nouvelle espèce du genre *Xysticus* (Araneae, Thomisidae) en Turquie. *Communications de la Faculté des Sciences de l'Université d'Ankara*, 11(C): 7-9.
- Karol, S. 1966c. Description d'une nouvelle espèce du genre *Oxyptila* en Turquie (Araneae, Thomisidae). *Communications de la Faculté des Sciences de l'Université d'Ankara*, 11(C): 11-14.
- Karol, S. 1966d. Spiders of Ankara and environs with a description of a new species *Xysticus turcicus* (Araneae, Thomisidae). *Communications de la Faculté des Sciences de l'Université d'Ankara*, 11(C): 15-32.
- Karol, S. 1967. Türkiye Örümcekleri. I. Ön Liste. *Ankara Üniversitesi Fen Fakültesi Yayınları*, 109(7): 1-37.
- Karol, S. 1968. Description de deux espèces nouvelles de Thomisidae (Araneae) de Turquie. *Bulletin du Museum National d'Historie Naturelle*, 39: 908-911.
- Kaya, R.S. & Uğurtaş, İ.H. 2007. A faunistic study on spiders (Araneae) of Terzioğlu Island (Uluabat Lake, Bursa). *Journal of Biological & Environmental Sciences*, 1(1): 31-36.
- Kulczyński, W. 1903. Arachnoidea in Asia Minore et ad Constantinopolim a Dre F. Werner collecta. *Sitzungsberichte der mathematisch-naturwissenschaftlichen Classe der kaiserlichen akademie der Wissenschaften*, 112: 627-680.
- Lehtinen, P.T. 2002. Generic revision of some thomisids related to *Xysticus* C.L.Koch, 1835 and *Ozyptila* Simon, 1864. In Toft, S. & N. Scharff (eds.), *European Arachnology 2000: Proceedings of the 19th European Colloquium of Arachnology*. Aarhus Univ. Press, pp. 315-327.
- Loerbroks, A. 1983. Revision der Krabbenspinnen-Gattung *Heriaeus* Simon (Arachnida: Araneae: Thomisidae). *Verhandlungen des naturwissenschaftlichen Vereins in Hamburg (NF)*, 26: 85-139.

- Logunov, D.V. 2006. Notes on *Xysticus kempeleni* Thorell, 1872 and two closely related spider species (Araneae: Thomisidae). *Acta arachnologica*, 55: 59-66.
- Logunov, D.V. & Demir, H. 2006. Further faunistic notes on *Cozyptila* and *Xysticus* from Turkey (Araneae, Thomisidae). *Arachnologische Mitteilungen*, 31: 40-45.
- Marusik, Y.M., Lehtinen, P.T. & Kovblyuk, M.M. 2005. *Cozyptila*, a new genus of crab spiders (Aranei: Thomisidae: Thomisinae: Coriarachnini) from the western Palaearctic. *Arthropoda Selecta*, 13: 151-163.
- Nosek, A. 1905. Araneiden, Opilionen und Chernetiden. In Penther, A. und E. Zederbauer, Ergebnisse einer naturwissenschaftlichen Reise zum Erdschias-Dagh (Kleinasien). *Annalen des Naturhistorischen Museums in Wien*, 20: 114-154.
- Ono, H. & Martens, J. 2005. Crab spiders of the families Thomisidae and Philodromidae (Arachnida: Araneae) from Iran. *Acta arachn. Tokyo*, 53: 109-124.
- Özdemir, A., Varol, İ., Akan, Z., Kütük, M., Kutbay, F. & Özaslan, M. 2006. The Fauna of Spider (Araneae) in the Nizip and Karkamış – Gaziantep (Turkey). *Biotechnology & Biotechnological Equipment*, 20(1): 74-77.
- Pavesi, P. 1876. Gli arachnidi Turchi. *Atti della Societa Italiana di Scienze Naturali*, 19(1): 1-27.
- Platnick, N.I. 2008. *The world spider catalog*, version 9.0. American Museum of Natural History, online at <http://research.amnh.org/entomology/spiders/catalog/index.html>
- Reimoser, E. 1919. Katalog der echten Spinnen (Araneae) des Paläarktischen Gebietes. *Abhandlungen der zoologisch-botanischen Gesellschaft in Wien*, 10(2): 1-280.
- Roewer, C.F. 1959. Die Araneae, Solifuga und Opiliones der Sammlungen des Herrn Dr. K. Lindberg aus Griechenland, Creta, Anatolien, Iran und Indien. *Göteborgs Kungliga Vetenskaps- och vitterhets-Samhalles Handlingar*, (6B) 8(4): 1-47.
- Simon, E. 1875. Liste d'arachnides de Constantinople et description de deux Opilionides. *Annales de la Société entomologique de France*, 5(5): 96-98.
- Simon, E. 1879. Liste d'arachnides de Constantinople et description d'une espèce nouvelle *Epeira turcica*. *Annales de la Société entomologique de France*, (5) 9(Bull.): 36-37.
- Simon, E. 1884. Etudes arachnologiques. 15e Mémoire. XXII. Arachnides recueillis par M. l'abbé David à Smyrne, à Beyrouth et à Akbès en 1883. *Annales de la Société entomologique de France*, (6) 4: 181-196.
- Simon, E. 1914. *Les arachnides de France. Synopsis générale et catalogue des espèces françaises de l'ordre des Araneae; Ire partie*. Paris, 6: 1-308.
- Strand, E. 1917. Arachnologica varia XIV-XVIII. *Archiv für Naturgeschichte*, 82(A2): 70-76.
- Topçu, A. & Demir, H. 2004. New crab spider (Araneae: Thomisidae) records for Turkey. *Israel Journal of Zoology*, 50: 421-422.
- Topçu, A., Demir, H. & Seyyar, O. 2005. A Checklist of the spiders of Turkey. *Serket*, 9(4): 109-140.
- Topçu, A., Demir, H., Seyyar, O. & Türkeş, T. 2006. The spider fauna of the Gülek Pass (Turkey) and its environs (Araneae). In C. Deltshev & P. Stoev (eds.), European Arachnology 2005. *Acta zoologica bulgarica Suppl.*, 1: 287-295.
- Varol, M.İ., Mart, C., Özaslan, M., Bayram, A., Akan, Z. & Özdemir, A. 2006. Investigation of spider fauna of Hançarık Dam-Nizip (Turkey). *Journal of Biological Sciences*, 6(2): 344-346.
- Vigna Taglianti, A.V., Audisio, P.A., Biondi, M., Bologna, M.A., Carpaneto, G.M., De Biase, A., Fattorini, S., Piattella, E., Sindaco, R., Venchi, A. & Zapparoli, M. 2000. A proposal for a chorotype classification of the Near East fauna, in the framework of the Western Palearctic Region. *Biogeographia*, 20: 31-59.
- Wunderlich, J. 1995. Zur Kenntnis west-paläarktischen Arten der Gattungen *Psammitis* Menge 1875, *Xysticus* C. L. Koch 1835 und *Ozyptila* Simon 1864 (Arachnida: Araneae: Thomisidae). *Beiträge zur Araneologie*, 4: 749-774.