

First Italian reports of *Nemoscolus laurae* (Simon, 1868) and *Zodarion isabellinum* (Simon, 1868) (Araneae Araneidae, Zodariidae) in Sicily and remarkable additions to the Sicilian Araneofauna

Antonino Dentici^{1*}, Antonino Barbera², Angelo Ditta³, Vincenzo Genna⁴ & Salvatore Surdo⁵

¹Via Enrico Cialdini 2, 90124 Palermo, Italy; e-mail: antonino.dentici.1991@gmail.com - Orcid: 0000-0003-0632-1633

²Via Giovanni Prati 28, 91022 Castelvetrano, Trapani, Italy; e-mail: abarberamail@gmail.com

³Via Trieste 16, 91026 Mazara del Vallo, Italy; e-mail: angelo_ditta@libero.it

⁴contrada Bosco 341, 91025 Marsala, Italy; e-mail: genna.vincenzo.1974@gmail.com

⁵Dipartimento di Scienze agrarie, alimentari e forestali, viale delle Scienze, Edificio 4, 90128 Palermo, Italy; e-mail: salvatore.surdo@unipa.it - Orcid:0000-0002-0300-837X

*Corresponding author

ABSTRACT

This paper reports some new data obtained in recent years during surveys on arachnids (Arachnida Araneae) present in Sicily (Italy). Particularly, fifteen new species of spiders are reported for the first time in Sicily: *Nemoscolus laurae* (Simon, 1868); *Emblyna brevidens* (Kulczyński, 1897); *Dysdera lata* Reuss, 1834; *Drassodes luteomicans* (Simon, 1878); *Ceratinella scabrosa* (O. Pickard-Cambridge, 1871); *Entelecara erythropus* (Westring, 1851); *Hilaira excisa* (O. Pickard-Cambridge, 1871); *Parapelecopsis nemoralis* (Blackwall, 1841); *Trichoncus helveticus* Denis, 1965; *Alopecosa cuneata* (Clerck, 1757); *Tibellus macellus* Simon, 1875; *Scytodes univittata* Simon, 1882; *Rhomphaea nasica* (Simon, 1873); *Zodarion isabellinum* (Simon, 1870); *Zodarion pusio* Simon, 1914. Six genera are reported for the first time for Sicily while *Nemoscolus laurae* and *Zodarion isabellinum* are new also for Italian fauna. Additional biological and taxonomic notes are also provided.

KEY WORDS

New record; *Scytodes univittata*; *Entelecara erythropus*; *Dysdera lata*; *Trichoncus helveticus*; *Hilaira excisa*.

Received 16.09.2023; accepted 09.11.2023; published online 30.12.2023

INTRODUCTION

Arachnological studies in Sicily have increased in recent years (Dentici & Galasso, 2021; Dentici, 2022; Dentici et al., 2022; Lenzini, 2022) leading to an increase in the reported species. Currently, the species known for the island amount to 465 -

223 genera and 43 families (Pantini & Isaia, 2019).

We are currently conducting several research on the population of Arachnids in Sicily, of which we summarize some results in the present work: 15 new species of spiders are reported, 2 are new for Italy, and 6 genera are reported for the first time for

Sicily; for each species, the data relating to the collection, are reported.

MATERIAL AND METHODS

Most specimens were collected on sight on plants, under the rocks or in their spiderwebs, some instead using pitfall traps. The specimens were observed, in laboratory, with stereomicroscope for identification, stored in centrifuge tubes of different sizes and fixed in 75% ethanol (Levi, 1977). The samples are stored in A. Dentici's collection (Palermo, Italy) and for each sample all the collection data in Italian language and the collector are indicated.

The maps were created using the QGIS open-source program. We decided to limit the distribution of the *Nemoscolus laurae* and *Zodarion isabellinum* up to Western Palearctic, while for the other species we only showed their presence or absence in the various regions of Italy. Distribution data outside this area was listed in the text. The classification, taxonomic order and nomenclatural arrangement for spiders follow Roberts (1995), Trotta (2004), Pantini & Isaia (2019), World Spider Catalog (2022) and Netwig et al. (2022). Other references have also been used and cited in the text.

RESULTS

Systematics

Familia ARANEIDAE Clerck, 1757
Genus *Nemoscolus* Simon, 1895

Nemoscolus laurae (Simon, 1868)

MATERIAL EXAMINED. Sicily (Italy), Trapani, Marsala, laguna dello Stagnone di Marsala, 37°51'45.8"N, 12°29'04.2"E, 10.VII.2022, 5 ♀♀ and 2 spiderlings, legit V. Genna.

DISTRIBUTION. Western Mediterranean.

REMARKS. The species and the genus are here reported for the first time in Italy. This species was observed for the first time in the Sicilian sampling site as early as August 2020, an occasion in which female specimens with spiderlings were observed (Figs. 1–4). In 2021, also in the month of August, adult fema-

les with spiderlings were observed. In 2022, the year of sampling of the above specimens, the same conditions (adult females with offspring) were observed in mid-June. These observations show a malleability of the reproductive period of the species or its link with temperatures, which in 2022 were hotter than in the previous two years for the same month. The observation of the species on these various occasions has allowed us to notice that the number of young born varies between 40 and 50 individuals (out of a sample of 10 observed cocoons). Another interesting observation concerns the probable "matriphagy". Out of 10 "conical shelters", typical of this species, 4 of these were inhabited only by juveniles, the remaining 6 by females and offspring. The theme of matriphagy is clearly speculative in this case, it has not been observed directly, but it is also undoubtedly frequent in other species of spiders such as *Stegodyphus lineatus* (Latreille, 1817) (Salomon et al., 2005), *Cheiracanthium japonicum* Bösenberg et Strand, 1906 (Toyama, 1999; 2003) or *Amaurobius ferox* (Walckenaer, 1830) (Kim & Horel, 2010).

Familia DICTYNIDAE O. Pickard-Cambridge, 1871
Genus *Emblyna* Chamberlin, 1948

Emblyna brevidens (Kulczyński, 1897)

MATERIAL EXAMINED. Sicily (Italy), Trapani, Mazara del Vallo, 37°40'07.8"N 12°35'01.2"E, 02.V.2023, 1 ♂ and 1 ♀, legit A. Ditta.

DISTRIBUTION. Europe (Fig. 5).

REMARKS. This genus and this species are here reported for the first time in Sicily. Both of these specimens were collected under the same small boulder.

Emblyna brevidens has been ascertained in Italy only in Goggau, municipality of Tarvisio (Udine, Friuli Venezia Giulia) (di Caporiacco, 1927).

Familia DYSDERIDAE C. L. Koch, 1837
Genus *Dysdera* Latreille, 1804

Dysdera lata Reuss, 1834

MATERIAL EXAMINED. Sicily (Italy), Monreale, Caculla, 38°03'03.2"N, 13°14'44.2"E, 16.IV.2017, 1 ♂, legit A. Dentici.



Figures 1–3. Female of *Nemoscolus laurae* from Stagnone di Marsala (Fig. 3) with spiderlings in the peculiar conical cocoon.




Figure 4. Map distribution of *Nemoscolus laurae*.




Figure 5. Map distribution of *Emblyna brevidens* in Italy.

DISTRIBUTION. Mediterranean to Georgia (Fig. 6).

REMARKS. *Dysdera lata* is here reported for the first time in Sicily; this specimen was found while wandering during the night.

Dysdera lata has been ascertained in Italy only in the municipality of San Ferdinando (Reggio Calabria, Calabria) (Pantini et al., 2020a).

Familia GNAPHOSIDAE Pocock, 1898
Genus *Drassodes* Westring, 1851

Drassodes luteomicans (Simon, 1878)

MATERIAL EXAMINED. Sicily (Italy), Trapani, Castellammare del Golfo, Balata di Baida, 38°01'32.6"N 12°47'19.0"E, 25.III.2022, 1 ♂, legit S. Surdo.

DISTRIBUTION. Southern Europe (Fig. 7).

REMARKS. This species is here reported for the first time in Sicily; *Drassodes luteomicans* was collected from its silk case, built under a small stone.

Drassodes luteomicans has been reported in Italy from Friuli-Venezia Giulia (di Caporiacco, 1927), Valle d'Aosta (di Caporiacco, 1928), Lombardia (di Caporiacco, 1941), Sardegna (Pantini et al., 2013; Bazzato et al., 2023).

Familia LINYPHIIDAE Blackwall, 1859
Genus *Ceratinella* Emerton, 1882

Ceratinella scabrosa (O. Pickard-Cambridge, 1871)

MATERIAL EXAMINED. Sicily (Italy), Palermo, Alia, C/da Bordone, 37°47'20"N, 13°43'13"E, 15.V.2022, 1 ♀, legit R. Viviano.

DISTRIBUTION. Europe (Fig. 8), Turkey, Caucasus, Russia (Europe to Far East).

REMARKS. This genus and this species are reported here for the first time in Sicily; it was found among the sediments collected under a rock.

Ceratinella scabrosa has been reported in Italy from Abruzzo (Isaia et al., 2009), Emilia-Romagna (Pantini & Isaia, 2008), Lombardia (Pesarini, 1996; Pesarini, 1997); Isaia et al., 2007), Piemonte (Arnò, 2001; Isaia et al., 2015), Trentino-Alto Adige (Steinberger, 2005; Steinberger, 2007; Steinberger & Zingerle, 2009) and Valle d'Aosta (Negro et al., 2009).

Genus *Entelecara* Simon, 1884

Entelecara erythropus (Westring, 1851)

MATERIAL EXAMINED. Sicily (Italy), Palermo, Monreale, Caculla, 38°03'02.7"N, 13°14'44.2"E, 25.II.2017, 1 ♀, legit A. Dentici; Sicily (Italy), Pa-




Figure 6. Map distribution of *Dysdera lata* in Italy.




Figure 7. Map distribution of *Drassodes luteomicans* in Italy.

lermo, Monreale, 38°02'45.7"N, 13°14'49.0"E, 13.III.2022, 1 ♂, legit A. Dentici.

DISTRIBUTION. Europe (Fig. 9), Russia (Europe to Far East), Kazakhstan, Iran, Japan.

REMARKS. This genus and this species are reported here for the first time in Sicily; this female specimen was found under a small stone, while the male specimen was found wandering over low vegetation.

Entelecara erythropus has been reported in Italy only from Giardini Biennale in the municipality of Venezia, Veneto (Hansen, 2002).

Genus *Hilaira* Simon, 1884

Hilaira excisa (O. Pickard-Cambridge, 1871)

MATERIAL EXAMINED. Sicily (Italy), Palermo, 38°08'31.4"N 13°20'15.4"E, 21.II.2022, 1 ♂, legit A. Ditta.

DISTRIBUTION. Europe (Fig 10).

REMARKS. This species and this genus are reported here for the first time in Sicily; *Hilaira excisa* was collected in an urban green area.

It has been reported in Italy from Friuli-Venezia Giulia (Hansen, 2011), Lombardia (Pantini et al., 2016; Pantini et al., 2020b), Piemonte (Isaia et al.,

2015), Trentino-Alto Adige (Zingerle, 1997; Steinberger, 2007a; Steinberger, 2008; Ballini & Steinberger, 2009; Steinberger, 2010; Ballini et al., 2014; Pantini et al., 2020), Valle d'Aosta (Paschetta et al., 2016) and Veneto (Hansen & Vanin, 2004; Vanin & Turchetto, 2007).

Genus *Parapelecopsis* Wunderlich, 1992

Parapelecopsis nemoralis (Blackwall, 1841)

MATERIAL EXAMINED. Sicily (Italy), Palermo, Carini, 38°10'45" N, 13°08'42" E, 01.V.2018, 1 ♀, legit R. Viviano.

DISTRIBUTION. Europe, Georgia (Fig 11).

REMARKS. This species and this genus are here reported for the first time in Sicily; this specimen was found under rotting wood.

Parapelecopsis nemoralis has been reported in Italy from Friuli-Venezia Giulia (di Caporiacco, 1922) and Toscana (di Caporiacco, 1923).

Genus *Trichoncus* Simon, 1884

Trichoncus helveticus Denis, 1965

MATERIAL EXAMINED. Sicily (Italy), Trapani, Salemi, 23.V.2021, 7 ♀♀, legit A. Ditta; Sicily




Figure 8. Map distribution of *Ceratinella scabrosa* in Italy.




Figure 9. Map distribution of *Entelecara erythropus* in Italy.

(Italy), Caltanissetta, Niscemi, Geloj Wetland, 37°06'01.8"N, 14°20'36.4"E, 15.I.2023, 1 ♂, legit A. Dentici; Sicily (Italy), Trapani, Mazara del Vallo, Borgata Costiera, 20.VI.2023, 2 ♀♀, legit A. Ditta.

DISTRIBUTION. Spain, France, Switzerland, Italy (Fig. 12).

REMARKS. *Trichoncus helveticus* is here reported for the first time in Sicily; these specimens were found under stones.

There are no certain data on the presence of *T. helveticus* in Italy. Recorded from Pesarini (1995), sources not available. Trotta (2005) recalls the record by Pesarini.

Familia LYCOSIDAE Sundevall, 1833
Genus *Alopecosa* Simon, 1885

Alopecosa cuneata (Clerck, 1757)

MATERIAL EXAMINED. Sicily (Italy), Palermo, Monreale, Sagana, 38°02'22.8"N, 13°12'04.8"E, 17.IV.2017, 1 ♀, legit A. Dentici; Sicily (Italy), Palermo, Monreale, Caculla, 38°03'07.9"N, 13°14'45.9"E, 29.IV.2018, 2 ♂♂ and 1 ♀, legit A. Dentici; Sicily (Italy), Trapani, Campobello di Mazara, 37°34'30.4"N 12°40'47.6"E, 30.IV.2023, 1 ♀, legit A. Ditta.

DISTRIBUTION. Europe (Fig. 13), Turkey, Caucasus, Russia (Europe to Far East), Kazakhstan, China.

REMARKS. *Alopecosa cuneata* is here reported for the first time in Sicily; these specimens were collected wandering on low vegetation during the evening, near the banks of the Sant'Elia river.

Alopecosa cuneata has been reported in Italy from Calabria (Ijland & Helsdingen van, 2019), Emilia-Romagna (Canestrini & Pavese, 1868), Friuli-Venezia Giulia (di Caporiacco, 1922; Hansen, 1997; Hansen, 2011), Lazio (Lacasella et al., 2014), Lombardia (Sordelli, 1868; Pavese, 1875; Pavese, 1879; Pantini et al., 2020b), Molise (Trotta, 2020), Piemonte (Lugetti & Tongiorgi, 1969; Lambiase et al., 2007; Isaia et al., 2007; Paschetta et al., 2013; Caprio et al., 2015; Isaia et al., 2015), Toscana (Lugetti & Tongiorgi, 1969; Cianferoni et al., 2010), Trentino-Alto Adige (Noflatscher, 1988; Noflatscher, 1990; Zingerle, 1997; Zingerle, 1999; Zingerle, 2000a; Zingerle, 2000b; Steinberger, 2007a; Steinberger, 2007b; Steinberger, 2008; Ballini et al., 2011; Ballini et al., 2014; Ballini et al., 2015; Fontana et al., 2020; Pantini et al., 2020b; Petri et al., 2021; Plunger et al., 2022), Valle d'Aosta (Isaia et al., 2008; De Angelis & Fantoni, 2008; Negro et al., 2009; Negro et al., 2013) and Veneto (Ninni, 1869; Denis, 1963; Zingerle, 1998; Zin-




Figure 10. Map distribution of *Hilaira excisa* in Italy.




Figure 11. Map distribution of *Parapelecopsis nemoralis* in Italy.

gerle, 2000a; Ballarin et al., 2011; Trotta & Cherubini, 2017; Nardi & Marini, 2021).

Familia PHILODROMIDAE Thorell, 1870
Genus *Tibellus* Simon, 1875

***Tibellus macellus* Simon, 1875**

MATERIAL EXAMINED. Sicily (Italy), Ragusa,

Roccazzo, Chiaramonte Gulfi, 24.IV.2021, 1 ♀, legit S. Surdo.

DISTRIBUTION. Europe (Fig. 14), Turkey, Caucasus, Russia (Europe to Far East), Kazakhstan.

REMARKS. *Tibellus macellus* is here reported for the first time in Sicily; this female specimen was found while stationed on low dry vegetation.




Figure 12. Map distribution of *Trichoncus helveticus* in Italy.




Figure 14. Map distribution of *Tibellus macellus* in Italy.




Figure 13. Map distribution of *Alopecosa cuneata* in Italy.




Figure 15. Map distribution of *Scytodes univittata* in Italy.

Tibellus macellus has been reported in Italy from Calabria (Caffi, 1895; Ijland & Helsdingen van, 2016; Ijland & Helsdingen van, 2019), Emilia-Romagna (di Caporiacco, 1926), Friuli-Venezia Giulia (di Caporiacco, 1949; Nardi et al., 2019; Lami et al., 2020), Veneto (di Caporiacco, 1950; Marcuzzi et al., 1971; Hansen & Iaconcig, 1999); Puglia (Ijland et al., 2012; Lucia et al., 2013), Campania (Ijland & Helsdingen van, 2014) and Basilicata (Ijland & Helsdingen van, 2016).

Familia SCYTODIDAE Blackwall, 1864
Genus *Scytodes* Latreille, 1804

Scytodes univittata Simon, 1882

MATERIAL EXAMINED. Sicily (Italy). Palermo, in domestic house, 02.XII.2018, 1 ♀, legit R. Viviano; idem, 24.VIII.2023, 1 ♀; Trapani, in domestic house, 08.V.2023, 1 ♂, legit L. Di Nicola; idem, 15.V.2023, 1 ♂; idem, in domestic house, 15.V.2023, 1 ♂, legit S. Surdo.

DISTRIBUTION. Egypt, Turkmenistan, Kyrgyzstan, India. Introduced to Hawaii, Mexico, Cuba, Venezuela, Brazil, Paraguay, Chile, Canary Islands, Spain (Fig. 15).

REMARKS. *Scytodes univittata* is reported here for the first time in Sicily.




Figure 16. Map distribution of *Rhomphaea nasica* in Italy.

These Sicilian specimens were found inside houses, in the first case specimens of *Scytodes thoracica* (Latreille, 1802) were found in the same house. The species has been reported in Italy from Lazio and Liguria (Trotta, 2017).

Familia THERIDIIDAE Sundevall, 1833
Genus *Rhomphaea* L. Koch, 1872

Rhomphaea nasica (Simon, 1873)

MATERIAL EXAMINED. Sicily (Italy), Palermo, Monreale, Caculla, 38°02'59.7"N, 13°14'36.4"E, 03.II.2018, 1 ♂, legit A. Dentici.

DISTRIBUTION. Canary Islands, Portugal, Spain, France, Italy (Fig. 16), Croatia, Greece, Africa, St. Helena.

REMARKS. This species is here reported for the first time in Sicily; the specimen was found on its intricate net built on the leaves of a lemon tree.

Rhomphaea nasica has been reported in Italy from Toscana (de Dalmas, 1922; Picchi et al., 2016; Picchi, 2020; Picchi et al., 2020); Puglia (di Caporiacco, 1951), Lazio (Brignoli, 1967), Sardegna (Pantini et al., 2013; Bosmans & Colombo, 2015), Basilicata (Ijland & Helsdingen van, 2016), Calabria (Ijland & Helsdingen van, 2016; Pantini & Mazzoleni, 2018; Ijland & Helsdingen van, 2019).



Figure 17. Map distribution of *Zodarion isabellinum*.

Familia ZODARIIDAE Thorell, 1881
Genus *Zodarion* Walckenaer, 1826

Zodarion isabellinum (Simon, 1870) (Fig. 17)

MATERIAL EXAMINED. Sicily (Italy), Trapani, Mazara del Vallo, Capo Granitola, 37°33'37.8"N 12°40'54.2"E, 03.06.VI.2023, 1 ♂, legit A. Ditta.

DISTRIBUTION. Spain, Morocco, Algeria.

REMARKS. This species is here reported for the first time in Sicily and Italy; it was found wandering on the ground, in a plot near the beach.

Identification follows Benhalima & Bosmans (2020).

Zodarion pusio Simon, 1914

MATERIAL EXAMINED. Sicily (Italy), Trapani, Triscina, 03.IV.2022, 1 ♀, legit A. Barbera.

DISTRIBUTION. France, Italy (Fig. 18), Slovenia, Croatia, Bosnia, Herzegovina, Tunisia.

REMARKS. This species is here reported for the first time in Sicily; this specimen was found under a small stone.

Zodarion pusio has been reported in Italy from Toscana (de Dalmas, 1922; Picchi, 2020), Calabria (Kritscher, 1969), Emilia-Romagna (Pesarini, 1991a;

Pesarini, 1993; Benassi & Dal Zotto, 2020), Lazio (Bosmans, 1997), Liguria (Bosmans, 1997; Trotta, 2019), Lombardia (Giordano et al., 2002; Isaia et al., 2007), Veneto (Ballarin et al., 2011; Ballarin & Petri, 2021), Sardegna (Bosmans & Colombo, 2015; Bosmans et al., 2019; Bazzato et al., 2022; Bazzato et al., 2023) and Basilicata (Ijland & Helsdingen van, 2016).

ACKNOWLEDGEMENTS

We want to thank Roberto Viviano (Palermo, Italy), Luigi and Francesco Di Nicola and Danilo Graffeo (Trapani, Italy) for providing us with useful specimens; Ignazio Sparacio (Palermo, Italy), for useful advice and the friendship that always shows us; our families, whose support is priceless and essential.

REFERENCES

Arnò C., 2001. Ragni dell'area protetta "Fascia fluviale del Po": nota preliminare su tre specie nuove per l'Italia e una nuova per il Piemonte (Arachnida, Araneae). *Rivista Piemontese di Storia Naturale*, 22: 155–164.

Ballarin F., Pantini P. & Hansen H., 2011. Catalogo ragionato dei ragni (Arachnida, Araneae) del Veneto. *Memorie del Museo Civico di Storia Naturale di Verona*, 2°serie, sezione biologica, 21: 1–151.

Ballini S., Stauder F. & Steinberger K.H., 2014. Webspinnen (Arachnida: Araneae). In: Schatz & Wilhalm. *Tag der Artenvielfalt 2013 auf den Armentara-Wiesen (Gemeinde Wengen, Südtirol, Italien)*. *Gredleriana*, 14: 311–315.

Ballini S. & Steinberger K.H., 2009. Weberknechte und Webspinnen (Opiliones und Araneae). In: Wilhalm T. (eds.): *GEO-Tag der Artenvielfalt 2008 am Reschenpass (Gemeinde Grau im Vinschgau, Südtirol, Italien)*. *Gredleriana*, 9: 315–319.

Ballini S., Stauder F. & Steinberger K.H., 2011. Weberknechte und Webspinnen (Opiliones und Araneae). In: *GEO-Tag der Artenvielfalt 2010 im Pfelderer Tal (Gemeinde Moos in Passeir, Südtirol, Italien)*. *Gredleriana*, 11: 195–198.

Ballini S., Stauder F. & Steinberger K.H., 2015. Webspinnen (Arachnida: Araneae). In: Schatz & Wilhalm. *Tag der Artenvielfalt 2014 in St. Felix (Gemeinde Unsere Liebe Frau im Walde St. Felix, Südtirol, Italien)*. *Gredleriana*, 15: 168–171.

Bazzato E., Lallai E., Caria M., Schifani E., Cillo D., An-




Figure 18. Map distribution of *Zodarion pusio* in Italy.

- cona C., Alamanni F., Pantini P., Maccherini S., Bacaro G. & Marignani M., 2022. Land-use intensification reduces multi-taxa diversity patterns of Small Woodlots Outside Forests in a Mediterranean area. *Agriculture, Ecosystems and Environment*, 340: 108–149.
<https://doi.org/10.1016/j.agee.2022.108149>
- Bazzato E., Lallai E., Caria M., Schifani E., Cillo D., Ancona C., Pantini P., Maccherini S., Bacaro G. & Marignani M., 2023. Focusing on the role of abiotic and biotic drivers on cross-taxon congruence. *Ecological Indicators*, 151: 110–323.
<https://doi.org/10.1016/j.ecolind.2023.110323>
- Benassi A. & Dal Zotto M., 2020. La pedofauna di un'area periferiale del Fiume Secchia (provincia di Reggio Emilia). *Atti della Società dei Naturalisti e Matematici di Modena*, 151: 71–84.
- Benhalima S. & Bosmans R., 2020. Revision of the genus *Zodariion* Walckenaer, 1833 (part IV). The species of Morocco (Araneae: Zodariidae). *Zootaxa*, 4899: 93–114.
<https://doi.org/10.11646/ZOOTAXA.4899.1.5>
- Bosmans R., 1997. Revision of the genus *Zodariion* Walckenaer, 1833, part II. Western and Central Europe, including Italy (Araneae: Zodariidae). *Bulletin of the British Arachnological Society*, 10: 265–294.
- Bosmans R. & Colombo M., 2015. New species of spiders from Sardinia (Araneae), with ecological notes on *Lipocrea epeiroides* (O. Pickard-Cambridge, 1872) (Araneae:Araneidae). *Arachnology*, 16: 319–332.
<https://doi.org/10.13156/arac.2015.16.9.319>
- Bosmans R., Pantini P., Loverre P. & Addante R., 2019. New species and new records of ant-eating spiders from Mediterranean Europe (Araneae: Zodariidae). *Arachnologische Mitteilungen / Arachnology Letters*, 57: 8–20.
<https://doi.org/10.30963/aramit5703>
- Brignoli P.M., 1967. Notizie sui Theridiidae del Lazio (Araneae). *Fragmenta entomologica*, Roma 4: 177–196.
- Caffi E., 1895. I Ragni di Calabria. *Tipografia S. Alessandro*, Bergamo, 62 pp.
- Canestrini G. & Pavesi P., 1868. Araneidi italiani. *Atti della Società italiana di Scienze Naturali*, Milano, 11: 738–872.
- Caporiacco L. di, 1922. Saggio sulla fauna aracnologica della Carnia e regioni limitrofe. *Memorie della Società Entomologica Italiana*, Genova, 1: 60–111.
- Caporiacco L. di, 1923. Aracnidi dei dintorni di Firenze. *Memorie della Società Entomologica Italiana*, Genova, 2: 177–226.
- Caporiacco L. di, 1926. Aracnidi della Provincia di Forlì. *Memorie della Società Entomologica Italiana*, Genova, 4: 229–258.
- Caporiacco L. di, 1927. Secondo saggio sulla-fauna aracnologica della Carnia e regioni limitrofe. *Memorie della Società Entomologica Italiana*, Genova, 5: 70–130.
- Caporiacco L. di, 1928. Aracnidi. Il Parco nazionale del Gran Paradiso, 2: 87–90.
- Caporiacco L. di, 1941. Aracnidi cavernicoli bresciani. *Le Grotte d'Italia* 2, 4: 3–11.
- Caporiacco L. di, 1949. Aracnidi della Venezia Giulia. *Atti del Museo Civico di Storia Naturale di Trieste*, 17: 29–43.
- Caporiacco L. di, 1950. Gli Aracnidi della Laguna di Venezia. II Nota. *Bollettino della Società Veneziana di Storia Naturale e del Museo Civico di Storia Naturale*, 5: 114–140.
- Caporiacco L. di, 1951. Aracnidi pugliesi raccolti dai Signori Conci, Giordani-Soika, Gridelli, Ruffo e dall'autore. *Memorie di Biogeografia Adriatica*, 2: 63–94.
- Caprio E., Nervo B., Isaia M., Allegro G. & Rolando A., 2015. Organic versus conventional systems in viticulture: Comparative effects on spiders and carabids in vineyards and adjacent forests. *Agricultural Systems*, 136: 61–69.
<https://doi.org/10.1016/j.agsy.2015.02.009>
- Cianferoni F., Mazza G. & Zinetti F., 2010. Ragni (pp. 213–245). In: Petri L. (Ed.). *Biodiversità in Provincia di Prato*. Volume 6 - Insetti e Ragni. Effigi Edizioni, 263 pp.
- De Angelis S. & Fantoni A., 2008. Contributo alla conoscenza della fauna araneologica (Arachnida, Araneae) del Parco Naturale Mont Avic con una nuova segnalazione per la fauna italiana. *Revue valdôtaine d'histoire naturelle*, 61-62: 109–116.
- Dalmas R. de, 1922. Catalogue des araignées récoltées par le Marquis G. Doria dans l'île Giglio (Archipel toscan). *Annali del Museo civico di Storia naturale di Genova*, 50: 79–96.
- Denis J., 1963. Araignées des Dolomites. *Atti dell'Istituto Veneto di Scienze, Lettere ed Arti*, 121: 253–271.
- Dentici A. & Galasso P., 2021. Checklist of spiders and Opiliones (Arachnida Araneae, Opiliones) of Sicilian southeast swamp lakes “Pantano Cuba”, “Pantano Longarini” and “Pantano Bruno” (Italy). *Fifth contribution to knowledge of Sicilian spider fauna*. *Biodiversity Journal*, 21: 695–703.
<https://doi.org/10.31396/Biodiv.Jour.2021.12.3.695.703>
- Dentici A., Barbera A., Ditta A., Surdo S., 2022. On some new reports on the spider fauna of Italy and Sicily (Arachnida Araneae). *Biodiversity Journal*, 13: 399–408.
<https://doi.org/10.31396/Biodiv.Jour.2022.13.2.399.408>
- Dentici A., 2022. Araneofauna (Arachnida Araneae) of Mount Pellegrino (Sicily, Italy). *Fourth contribution*

- to knowledge of the Sicilian spider fauna. *Biodiversity Journal*, 13: 641–650.
<https://doi.org/10.31396/Biodiv.Jour.2022.13.3.641.650>
- Fontana V., Guariento E., Hilpold A., Niedrist G., Steinwandter M., Spitale D., Nascimbene J., Tappeiner U. & Seeber J., 2020. Species richness and beta diversity patterns of multiple taxa along an elevational gradient in pastured grasslands in the European Alps. *Scientific Reports*, 10: 12516.
<https://doi.org/10.1038/s41598-020-69569-9>
- Giordano V., Lazzarini M. & Bogliani G., 2002. Biodiversità ambientale in ambiente urbano. Il caso della città di Pavia. *Fondazione Lombardia per l'ambiente*, 147 pp.
- Hansen H., 1997. Beitrag zur Kenntnis der Spinnenfauna der Umgebung des Karst-Sees von Doberdò, Prov. Gorizia, Italien (Arachnida: Araneae). *Bollettino del Museo Civico di Storia Naturale di Venezia*, 47: 243–250.
- Hansen H. & Iaconcig M., 1999. Contributo alla conoscenza dell'aracnofauna di alcuni biotopi in prossimità della foce del Tagliamento, NE-Italia (Arachnida Araneae). *Bollettino del Museo Civico di Storia Naturale di Venezia*, 49: 99–109.
- Hansen H., 2002. Segnalazioni 21-40 Arachnida Araneae. *Bollettino del Museo Civico di Storia Naturale di Venezia*, 53: 268–272.
- Hansen H. & Vanin S., 2004. Contributo alla conoscenza della fauna araneologica dell'arco alpino sud-orientale. *Bollettino del Museo Civico di Storia Naturale di Venezia*, 55: 87–95.
- Hansen H., 2011. Contributo alla conoscenza dell'araneofauna di alcuni biotopi naturali del Friuli-Venezia Giulia (Arachnida Araneae). *Gortania. Botanica e zoologica*, 32: 115–134.
- Ijland S., Helsdingen van P. J. & Miller J., 2012. On some spiders from Gargano, Apulia, Italy. *Nieuwsbrief SPINED*, 32: 2–20.
- Ijland S. & Helsdingen van P. J., 2014. On some spiders (Arachnida, Araneae) from the surroundings of Castellabate, Italy. *Nieuwsbrief SPINED*, 34: 16–33.
- Ijland S. & Helsdingen van P. J., 2016. On some spiders (Arachnida, Araneae) of Basilicata and Calabria, Italy. *Nieuwsbrief SPINED*, 36: 25–36.
- Ijland S. & Helsdingen van P. J., 2019. Update on the spiders (Arachnida, Araneae) of Calabria, Italy. *Nieuwsbrief SPINED*, 38: 8–21.
- Isaia M., 2005. Check-list delle specie di ragni (Arachnida, Araneae) della Valle d'Aosta con una nuova segnalazione per la fauna italiana. *Revue valdôtaine d'histoire naturelle*, 59: 25–43.
- Isaia M., Pantini P., Beikes S. & Badino G., 2007. Catalogo ragionato dei ragni (Arachnida, Araneae) del Piemonte e della Lombardia. *Memorie dell'Associazione Naturalistica Piemontese*, 9: 9–161.
- Isaia M., Osella G. & Pannunzio G., 2009. Hygropetric and litter-inhabiting spiders (Araneae) from the Abruzzo Apennines (Central Italy). *Arachnologische Mitteilungen*, 37: 15–26.
- Isaia M., Paschetta M. & Chiarle A., 2015. Annotated checklist of the spiders (Arachnida, Araneae) of the Site of Community Importance and Special Area of Conservation Alpi Marittime (NW Italy). *Zoosystema*, 37: 57–114.
<https://doi.org/10.5252/z2015n1a4>
- Isaia M., Negro M. & Rolando A., 2008. Notes on the distribution of *Berlandina nubivaga* with the description of the male (Araneae, Gnaphosidae). *The Journal of Arachnology*, 36: 180–183.
- Kil-Won Kim K. & Horel A., 2010. Matrifagy in the Spider *Amaurobius ferox* (Araneidae, Amaurobiidae): an Example of Mother-Offspring Interactions, in *Ethology*, vol. 104, n. 12: 1021–1037.
<https://doi.org/10.1111/j.1439-0310.1998.tb00050.x>
- Kritscher E., 1969. Ein Beitrag zur Kenntnis der Araneen-Fauna Italiens. *Memorie del Museo Civico di Storia Naturale di Verona*, 16: 271–319.
- Lacasella F., Gratton C., De Felici S., Isaia M., Zapparoli M., Marta S. & Sbordoni V., 2014. Asymmetrical responses of forest and “beyond edge” arthropod communities across a forest-grassland ecotone. *Biodiversity and Conservation*, 24: 447–465.
<https://doi.org/10.1007/s10531-014-0825-0>
- Lami F., Bartomeus I., Nardi D., Beduschi T., Boscutti F., Pantini P., Santoiemma G., Scherber C., Tschardtke T. & Marini L., 2020. Species-habitat networks elucidate landscape effects on habitat specialization of natural enemies and pollinators. *Ecology Letters*, 24: 288–297.
<https://doi.org/10.1111/ele.13642>
- Lambiase S., De Angelis S., Fantoni A., Fasola M. & Pesarini C., 2007. A contribution to the knowledge of the araneae (Arthropoda, Arachnida) of the Maritime Alps, northern Italy. *Atti della Società italiana di Scienze Naturali e del Museo civico di Storia naturale di Milano*, 148: 153–160.
- Lenzini L., Castellucci F., Poso M., Kulczycki A., Simeon E., Greco G., Piccinini A. & Legittimo C.M., 2022. First records of *Anagraphis ochracea* (Araneae, Gnaphosidae) for continental Italy and Sicily with new observations on its myrmecophilous lifestyle. *Arachnologische Mitteilungen*, 64: 83–92.
<https://doi.org/10.30963/aramit6410>
- Lucia M., Loverre P. & Addante R., 2013. Indagine sull'araneofauna della riserva naturale regionale orientata “Laghi di Conversano e Gravina di Monsignore”, pp. 71–76. In: Calabrese G., Pacucci C., Occhialini W., Russo G. (a cura di), *Atti del IX Convegno nazionale di biodiversità*, Vol. 3, Territorio, paesaggio

- e servizi eco-sistemici, 5–7 settembre 2012, Istituto agronomico mediterraneo di Bari, Valenzano (BA).
- Lugetti G. & Tongiorgi P., 1969. Ricerche sul genere *Alopecosa* Simon (Araneae, Lycosidae). Atti della Società Toscana di Scienze Naturali residente in Pisa. Memorie. Serie B, 76: 1–100.
- Marcuzzi G., Dalla Venezia L. & Lorenzoni A.M., 1971. Appunti ecologico-qualitativi sul popolamento animale di alcuni biotopi litorali dell'alto Adriatico. Atti dell'Istituto Veneto di Scienze, Lettere ed Arti, 129: 119–207.
- Nardi D., Lami F., Pantini P. & Marini L., 2019. Using species-habitat networks to inform agricultural landscape management for spiders. *Biological Conservation*, 239: 1–8.
<https://doi.org/10.1016/j.biocon.2019.108275>
- Nardi D. & Marini L., 2021. Role of abandoned grasslands in the conservation of spider communities across heterogeneous mountain landscapes. *Agriculture, Ecosystems and Environment*, 319: 1–9.
<https://doi.org/10.1016/j.agee.2021.107526>
- Negro M., Isaia M., Palestini C. & Rolando A., 2009. The impact of forest ski-pistes on diversity of ground-dwelling arthropods and small mammals in the Alps. *Biodiversity and Conservation*, 18: 2799–2821.
- Negro M., Rolando A., Barni E., Bocola D., Filippa G., Freppaz M., Isaia M., Siniscalco C. & Palestini C., 2013. Differential responses of ground dwelling arthropods to ski-piste restoration by hydroseeding. *Biodiversity and Conservation*, 22: 2607–2634.
<https://doi.org/10.1007/s10531-013-0544-y>
- Noflatscher M.T., 1988. Ein Beitrag zur Spinnenfauna Südtirols: Epigäische an Xerotherm- und Kulturstandorten bei Albeins (Arachnida: Aranei). *Berichte des Naturwissenschaftlich-Medizinischen Vereins in Innsbruck*, 75: 147–170.
- Ninni A.P., 1869. Catalogo degli araneidi trevigiani. *Venezia*, 1–10.
- Noflatscher M.T., 1990. Zweiter Beitrag zur Spinnenfauna Südtirols: Epigäische Spinnen an Xerothermstandorten bei Säben, Guntzschna und Castelfeder (Arachnida: Aranei). *Berichte des Naturwissenschaftlich-Medizinischen Vereins in Innsbruck*, 77: 63–75.
- Pantini P., Isaia M., 2008. New records for the Italian spider fauna (Arachnida, Araneae). *Arthropoda Selecta*, 17: 133–144.
- Pantini P., Isaia M., Mazzoleni F. & Oneto C., 2016. Nuovi dati sui ragni di Lombardia (Arachnida, Araneae). *Rivista del Museo civico di Scienze Naturali “E. Caffi”*, Bergamo, 29: 21–44.
- Pantini P. & Mazzoleni F., 2018. I ragni di Calabria (Arachnida, Araneae). *Rivista del Museo civico di Scienze Naturali “E. Caffi”*, Bergamo, 31: 11–70.
- Pantini P. & Isaia M., 2019. Araneae.it: the online Catalog of Italian spiders with addenda on other Arachnid Orders occurring in Italy (Arachnida: Araneae, Opiliones, Palpigradi, Pseudoscorpionida, Scorpiones, Solifugae). *Fragmenta Entomologica*, 51: 127–152. Online at www.araneae.it, accessed on {15/10/2023}.
- Pantini P., Bonelli D. & Bonacci T., 2020a. I ragni epigei (Arachnida, Araneae) di un ambiente litoraneo della Calabria. *Rivista del Museo civico di Scienze Naturali “E. Caffi”*, Bergamo, 32: 25–32.
- Pantini P., Mazzoleni F., Gobbi M. & Pedrotti L., 2020b. Ragni (Arachnida, Araneae) di interesse biogeografico e conservazionistico nel Parco Nazionale dello Stelvio (Italia). *Rivista del Museo civico di Scienze Naturali “E. Caffi”*, Bergamo, 33: 23–53.
- Pantini P., Sassu A. & Serra G., 2013. Catalogue of the spiders (Arachnida Araneae) of Sardinia. *Biodiversity Journal*, 4: 3–104.
- Paschetta M., La Morgia V., Masante D., Negro M., Rolando A. & Isaia M., 2013. Grazing history influences biodiversity: a case study on ground-dwelling arachnids (Arachnida: Araneae, Opiliones) in the Natural Park of Alpi Marittime (NW Italy). *Journal of Insect Conservation*, 17: 339–356.
<https://doi.org/10.1007/s10841-012-9515-y>
- Paschetta M., Christille C., Marguerettaz F. & Isaia M., 2016. Regional catalogue of the spiders (Arachnida, Araneae) of Aosta Valley (NW Italy). *Zoosystema*, 38: 49–125. <https://doi.org/10.5252/z2016n1a3>
- Pavesi P., 1875. Note araneologiche. II. Aggiunte all'enumerazione dei ragni dei dintorni di Pavia. *Atti della Società italiana di Scienze Naturali, Milano*. 18: 130–132.
- Pavesi P., 1879. Saggio di una fauna aracnologica del Varesotto. *Atti della Società italiana di Scienze Naturali, Milano*, 21: 789–817.
- Pesarini C., 1991. Primo contributo per una fauna araneologica del ferrarese (Arachnida: Araneae). *Quaderni della stazione di ecologia del civico museo di storia naturale di Ferrara*, 4: 5–34.
- Pesarini C., 1993. Note su alcuni Zodarion Walckenaer della fauna italiana, con descrizione di una nuova specie (Araneae, Zodariidae). *Atti della Società italiana di Scienze Naturali e del Museo civico di Storia naturale di Milano*, 133: 5–12.
- Pesarini C., 1994. Arachnida Araneae. In Minelli A., Ruffo S. & La Posta S. (eds.) *Checklist delle specie della fauna italiana*. Calderini, Bologna, 23: 1–42.
- Pesarini C., 1996. Note su alcuni Erigonidae italiani, con descrizione di una nuova specie (Araneae). *Atti della Società italiana di Scienze Naturali e del Museo civico di Storia naturale di Milano*, 135: 413–429.
- Pesarini C., 1997. I Ragni (Arachnida Araneae) del Monte Barro (Italia, Lombardia, Lecco). *Memorie*

- della Società Italiana di Scienze Naturali e del Museo Civico di Storia Naturale di Milano, 27: 251–263.
- Petri I., Bernasconi M., Ballarin F., Pantini P., Armanini M., Caccianiga M., Chirichella R., Lencioni V., Mustoni A. & Gobbi M., 2021. Ragni (Arachnida: Araneae) d’alta quota delle Dolomiti di Brenta. Studi Trentini di Scienze Naturali, 101: 73–81.
- Picchi M.S., Bocci G., Petacchi R. & Entling M.H., 2016. Effects of local and landscape factors on spiders and olive fruit flies. Agriculture, Ecosystems and Environment, 222: 138–147.
<https://doi.org/10.1111/j.agee.2016.01.045>
- Picchi M.S., 2020. Spiders (Araneae) of olive groves and adjacent semi-natural habitats from central Italy. Arachnologische Mitteilungen / Arachnology Letters, 60: 1–11.
<https://doi.org/10.30963/aramit6001>
- Picchi M.S., Bocci G., Petacchi R. & Entling M.H., 2020. Taxonomic and functional differentiation of spiders in habitats in a traditional olive producing landscape in Italy. Europea Journal of Entomology, 117: 18–26.
<https://doi.org/10.14411/eje.2020.002>
- Plunger J., Guariento E., Steinwandter M., Colla F., Rief A. & Seeber J., 2022. Shifts in ground-dwelling predator communities in response to changes in management intensity in Alpine meadows. Soil organisms, 94: 149–161.
<https://doi.org/10.25674/so94iss3id306>
- Rief A. & Ballini S., 2017. Erhebung der Spinnen und Weberknechte (Arachnida: Araneae, Opiliones) in den LTSEER-Untersuchungsflächen in Matsch (Südtirol, Italien) im Rahmen der Forschungswoche 2016. Gredleriana, 17: 173–183.
- Salomon M., Aflalo E. D. & Coll M., 2015. Dramatic histological changes preceding suicidal maternal care in the subsocial spider *Stegodyphus lineatus* (Araneae: Eresidae). The Journal of Arachnology, 43: 77–85. <https://doi.org/10.1636/B14-15.1>.
- Sordelli F., 1868. Sui Ragni Lombardi. Atti della Società italiana di Scienze Naturali, Milano, 11: 459–476.
- Steinberger K.H., 2005. Die Spinnen (Araneae) und Weberknechte (Opiliones) der Etsch-Auen in Südtirol (Italien). Gredleriana, 4: 55–92.
- Steinberger K.H., 2007a. Spinnen und Weberknechte (Arachnida: Araneae, Opiliones) Südtiroler Flusslandschaften - Auwaldfragmente an Eisack und Rienz (Südtirol, Italien). Gredleriana, 7: 171–194.
- Steinberger K.H., 2007b. Weberknechte und Webspinnen (Opiliones und Araneae). In: GEO-Tag der Artenvielfalt 2007 am Fluß des Plattkofels (Seiser Alm, Gemeinde Kastelruth, Südtirol, Italien). Gredleriana, 7: 438–440.
- Steinberger K.H., 2008. Spinnen und Weberknechte im Naturpark Schlern - Rosengarten (Arachnida: Araneae, Opiliones) (Italien, Südtirol). Gredleriana, 8: 255–286.
- Steinberger K. H. & Zingerle V., 2009. Zur Spinnenfauna (Arachnida, Araneae) von -Prader Sand- und -Schludernser Au-, Reste einer naturnahen Flusslandschaft im oberen Etschtal (Vinschgau, Südtirol, Italien). Gredleriana, 9: 213–230.
- Steinberger K.H., 2010. Spiders of arable land in the Eisack and the Puster Valley (South Tyrol, Italy) (Arachnida: Araneae). Gredleriana, 10: 227–238.
- Toyama M., 1999. Adaptive advantages of maternal care and matrophagy in a foliage spider, *Cheiracanthium japonicum* (Araneae: Coubionidae). Journal of Ethology, 17: 33–39.
<https://doi.org/10.1007/BF02769295>.
- Toyama M., 2003. Relationship between reproductive resource allocation and resource capacity in the matrophagous spider, *Cheiracanthium japonicum* (Araneae: Clubionidae). Journal of Ethology, 21: 1–7.
<https://doi.org/10.1007/s10164-002-0067-6>.
- Trotta A., 2005. Introduzione ai ragni italiani (Arachnida Araneae). Memorie della Società Entomologica Italiana, Genova, 83: 3–178.
- Trotta A., 2017. Segnalazioni Faunistiche Italiane. 609 - *Scytodes univittata* Simon, 1882 (Araneae, Scytodidae). Bollettino della Società Entomologica Italiana, Genova, 149: 93.
- Trotta A., 2019. Note on spiders collected in coastal habitats and first Italian record of *Chaerea maritimus* (Arachnida, Araneae). Bulletin of Environmental and Life Sciences, 1: 33–36.
- Trotta A., 2020. Spiders from Molise (Italy): state of knowledge, new faunistic data and taxonomic notes (Arachnida: Araneae). Fragmenta entomologica, Roma, 52: 77–83.
<https://doi.org/10.13133/2284-4880/415>
- Trotta A. & Cherubini A., 2017. Contributo alla conoscenza dei ragni (Arachnida, Araneae) del Veneto. Bollettino del Museo Civico di Storia Naturale di Verona, Botanica Zoologia, 41: 55–86.
- Vanin S. & Turchetto M., 2007. Winter activity of spiders and pseudoscorpions in the South-Eastern Alps (Italy). Italian Journal of Zoology, 74: 31–38.
- Zingerle V., 1997. Epigäische Spinnen und Weberknechte im Naturpark Puez-Geisler (Dolomiten, Südtirol) (Araneae, Opiliones). Berichte des Naturwissenschaftlich-Medizinischen Vereins in Innsbruck, 84: 171–226.
- Zingerle V., 1999. Epigäische Spinnen und Weberknechte im Naturpark Sextner Dolomiten und am Sellajoch (Südtirol, Italien) (Araneae, Opiliones). Berichte des Naturwissenschaftlich-Medizinischen Vereins in Innsbruck, 86: 165–200.
- Zingerle V., 1998. Ragni e Opilioni del Monte Grappa, Dolomiti, Italia (Araneae, Opiliones). Gruppo Natura Bellunese, Convegno “Aspetti naturalistici della provincia di Belluno” - Belluno 30 maggio 1998, 227–245.

Zingerle V., 2000a. Epigäische Spinnen und Weberknechte aus den nördlichen Dolomiten: Valparola-Pass und Weißhorn (SE-Alpen, Italien) (Araneae, Opiliones). *Berichte des Naturwissenschaftlich-Medizinischen Vereins in Innsbruck*, 87: 165–207.

Zingerle V., 2000b. Zoocenosi di ragni e opilioni nelle Dolomiti sudoccidentali (Parco Naturale Panaveggio - Pale di San Martino, Italia) (Araneae, Opiliones). *Studi Trentini di Scienze Naturali. Acta Biologica*, 75: 87–107.